

Guide to production of pocket cards:

Print page 2 in colors onto A5 paper, fold lengthwise and laminate the card.

After randomisation

- Inform the clinical team about the randomisation
- Write an inclusion note and update relevant ICU notes inside the patient chart
- Update relevant medication orders
- Inform the patient's proxies as soon as possible in order to obtain consent for continued participation

HOTLINE
+45 3545 0606
Available 24/7

www.incept.dk

Screen
your patient



app.easyrf.dk

Version 1.0.1, 2026-05-18

Adult patient (≥ 18 years old) acutely admitted to the ICU

- AND -
Eligible for at least one active domain

INCEPT-Albumin

Shock defined as both of the following fulfilled:

- Ongoing infusion of a vasopressor or inotropic agent
- Plasma lactate ≥ 2 mmol/L in any blood sample performed within the last 3 hours

- AND -
Both interventions in this domain are considered clinically appropriate (i.e., giving albumin for fluid resuscitation and substitution, or not giving albumin)

INCEPT-Thromboprophylaxis

Decision by the treating clinician to start thromboprophylaxis with LMWH, or to continue LMWH prophylaxis

- AND -
All interventions in this domain are considered clinically appropriate (i.e., giving LMWH in a fixed low, fixed intermediate, or weight-adjusted dose)