Light-induced superconductivity via dynamical Hubbard U

Funded through DFG Emmy Noether programme (SE 2558/2)

Bloch group meeting, MPQ Munich, January 19, 2021

Max Planck Institute for the Structure and Dynamics of Matter

Michael A. Sentef lab.sentef.org




Optical response of a superconductor

$$j(t) = \int_{-\infty}^{\infty} \mathrm{d}t' \sigma(t - t') E(t')$$

$$\sigma_{\rm Drude}(\omega) = \frac{ie^2(n/m)_{\rm eff}}{\omega + i/\tau}$$

Dissipationless case: $I/\tau \rightarrow 0^+$ **Real part:** delta-function peak at zero frequency **Imaginary part:** I/ω divergence n = superfluid density

Max Planck Institute for the Structure and Dynamics of Matter




The weight of the delta-function peak / divergence in Re/Im σ measures the superfluid density!








Optically driven superconductivity via vibrational excitation



W. Hu *et al.*, Nat. Mater. **13**, 705 (2014) R. Mankowsky *et al.*, Nature **516**, 71 (2014)

Max Planck Institute for the Structure and Dynamics of Matter

mps




M. Mitrano *et al.*, Nature **530**, 461 (2016) A. Cantaluppi *et al.*, Nat. Phys. **14**, 837 (2018)





Driving the apical oxygen phonon in YBCO

~15 µm 200 fs 5 MV/cm




S. Kaiser *et al.*, Phys. Rev. B **89**, 184516 (2014) C. Hunt *et al.*, Phys. Rev. B **94**, 224303 (2016)


Max Planck Institute for the Structure and Dynamics of Matter

mpsd

1) Transient Plasma Resonance




2) Divergent imaginary conductivity






Driving the apical oxygen phonon in YBCO

~15 µm 200 fs 5 MV/cm



S. Kaiser et al., Phys. Rev. B 89, 184516 (2014) C. Hunt *et al.*, Phys. Rev. B **94**, 224303 (2016)


mpso






Optically driven superconductivity in K₃C₆₀

 $K_{3}C_{60}$ T_C = 20 K





mpsd




M. Mitrano *et al.*, Nature **530**, 461 (2016) A. Cantaluppi et al., Nat. Phys. 14, 837 (2018)


Max Planck Institute for the Structure and Dynamics of Matter




Optically driven superconductivity in K₃C₆₀

Equilibrium $T < T_c$








Max Planck Institute for the Structure and Dynamics of Matter



What about other materials? Can we get a handle on the microscopic mechanism?



Building block of many organic charge transfer salts


Max Planck Institute for the Structure and Dynamics of Matter

M. Buzzi et al., PRX 10, 031028 (2020)

Typically electron donor



Photo-molecular high-temperature superconductivity in an organic kappa salt



Müller et al., Crystals (2018)

Cation Layer (ET molecules)

X Anion Layer


Cation Layer (ET molecules)

X Anion Layer

Cation Layer (ET molecules)

Max Planck Institute for the Structure and Dynamics of Matter

M. Buzzi et al., PRX 10, 031028 (2020)



Effectively a 2D U-t-t' Hubbard model on an anisotropic triangular lattice












Photo-molecular high-temperature superconductivity in an organic kappa salt




kappa-(ET)₂X phase diagram

D. Faltermeier *et al.*, Phys. Rev. B **76**, 165113 (2007)

Nernst effect indicating superconducting fluctuations above Tc: M.-S. Nam et al., Nature **449**, 584 (2007)

Max Planck Institute for the Structure and Dynamics of Matter

M. Buzzi et al., PRX 10, 031028 (2020)



"kappa" arrangement

ET molecules paired in dimers


Half filled conduction band








Photo-molecular high-temperature superconductivity in an organic kappa salt



M. Buzzi et al., PRX 10, 031028 (2020)


Driving with a specific molecular vibration



Max Planck Institute for the Structure and Dynamics of Matter



Photo-molecular high-temperature superconductivity in an organic kappa salt



(a)

2D (meV)

mps



M. Buzzi et al., PRX 10, 031028 (2020)









Photo-molecular high-temperature superconductivity in an organic kappa salt

It only works for specific molecular vibrations



Max Planck Institute for the Structure and Dynamics of Matter


mps

M. Buzzi et al., PRX 10, 031028 (2020)



Photo-molecular high-temperature superconductivity in an organic kappa salt

What is special about these vibrations?



M. Buzzi et al., PRX 10, 031028 (2020)

The C=C bond stretching modes change Hubbard U!


Max Planck Institute for the Structure and Dynamics of Matter




Photo-molecular high-temperature superconductivity in an organic kappa salt

What is special about these vibrations?

(a) C=C stretching mode (1470 cm⁻¹)





Max Planck Institute for the Structure and Dynamics of Matter


M. Buzzi et al., PRX 10, 031028 (2020)

tensor net calculations for dynamically driven U-t-t' Hubbard diamond chain



Photo-molecular high-temperature superconductivity in an organic kappa salt

Theory follow-up: J. Tindall et al., PRL 125, 137001 (2020)



M. Buzzi et al., PRX 10, 031028 (2020)

$$\begin{split} H(t) &= -\tau(t) \sum_{ij \in \langle \mathbf{n}.\mathbf{n} \rangle, \sigma} (c^{\dagger}_{\sigma,i} c_{\sigma,j} + \mathbf{h}.\mathbf{c}) - \\ \tau' \sum_{ij \in \langle \mathbf{vert} \rangle, \sigma} (c^{\dagger}_{\sigma,i} c_{\sigma,j} + \mathbf{h}.\mathbf{c}) + U(t) \sum_{i} n_{i,\uparrow} n_{i,\downarrow}, \end{split}$$

Approximate eta SU(2) symmetry (exact for τ '=0, bipartite lattice):

$$[H(t),\eta^z] \equiv 0, \qquad [H(t),\eta^+\eta^-] \propto \tau', \qquad (3)$$

where

$$\eta^{+} = \sum_{i=1}^{L} f(i) c_{i,\uparrow}^{\dagger} c_{i,\downarrow}^{\dagger}, \qquad \eta^{z} = \sum_{i} (n_{\uparrow,i} + n_{\downarrow,i} - 1), \quad (4)$$


put reference to Yang & Zhang, SO4 paper

Max Planck Institute for the Structure and Dynamics of Matter




Photo-molecular high-temperature superconductivity in an organic kappa salt

Theory follow-up: J.Tindall et al., PRL 125, 137001 (2020)



Rate of change of total $\langle \eta^+ \eta^- \rangle$ tends to zero for T' to zero



Enhancement of long-range pairing persists up to a critical value of τ '=0.7 τ

Max Planck Institute for the Structure and Dynamics of Matter


mp

M. Buzzi et al., PRX 10, 031028 (2020)

"heating-induced order": driving heats the system maximizing its entropy subject to symmetry constraints



What is the big picture?



tuning parameter (doping, pressure, chemical substitution)

Max Planck Institute for the Structure and Dynamics of Matter

can this be tested for a 2D Hubbard model with ultracold fermions?

what are ideal conditions for light-induced SC?

- superconducting instability at low T
- temperature typically 3-5 times equilibrium Tc
- intermediate to strong correlations fluctuation regime about
- driving approximately preserves a symmetry related to SC





ove	Т	С



Acknowledgments



MPSD Hamburg Nicolas Tancogne-Dejean, Michele Buzzi, Daniele Nicoletti, Andrea Cavalleri



Max Planck Institute for the Structure and Dynamics of Matter



Oxford University Joey Tindall, Frank Schlawin, Dieter Jaksch

Thank you for your attention!



