

Place in Site Master File #9

Pocket cards

At the following two pages you will find front and back of a pocket card for clinicians screening and randomising patients in the SUP-ICU trial. There are four identical front pages and four identical back pages.

How to make the cards:

- 1. Write the personal username and password on the back side of the card
- 2. Print the two pages
- 3. Laminate the pages back to back
- 4. Cut out the cards

Inclusion criteria

Acute admission to the ICU

Aged 18 years or above

AND

One or more of the following:

(continuous infusion of vasopressor/inotrope OR MAP < 70 mmHg OR systolic BP < 90 mmHg OR lactate > 4 mmol/l)

Renal replacement therapy

Invasive mechanical ventilation

(expected duration > 24 hours; If in doubt answer YES)

Acute or chronic coagulopathy

(platelets < 50 OR INR > 1.5 OR PT > 20 seconds)

Anticoagulant treatment

(does <u>not</u> include prophylactic doses of heparin/LMWH or acetylsalicylic acid - aspirin)

History of liver disease

(portal hypertension OR variceal bleeding OR hepatic ncephalopathy OR cirrhosis proven by biopsy, CT or ultrasound)

Hotline: +45 3545 7450 **TURN OVER**

Inclusion criteria

Acute admission to the ICU



Aged 18 years or above

AND

One or more of the following:

Shock

(continuous infusion of vasopressor/inotrope OR MAP < 70 mmHg OR systolic BP < 90 mmHg OR lactate > 4 mmol/l)

Renal replacement therapy

(acute OR chronic)

Invasive mechanical ventilation

(expected duration > 24 hours; If in doubt answer YES)

• Acute or chronic coagulopathy (platelets < 50 OR INR > 1.5 OR PT > 20 seconds)

Anticoagulant treatment

(does <u>not</u> include prophylactic doses of heparin/LMWH or acetylsalicylic acid - aspirin)

History of liver disease

(portal hypertension OR variceal bleeding OR hepatic encephalopathy OR cirrhosis proven by biopsy, CT or ultrasound)

Hotline: +45 3545 7450 **TURN OVER**

Inclusion criteria

Acute admission to the ICU

Aged 18 years or above

AND

One or more of the following:

Shock

(continuous infusion of vasopressor/inotrope OR MAP < 70 mmHg OR systolic BP < 90 mmHg OR lactate > 4 mmol/l)

Renal replacement therapy

• Invasive mechanical ventilation

(expected duration > 24 hours; If in doubt answer YES)

Acute or chronic coagulopathy

(platelets < 50 OR INR > 1.5 OR PT > 20 seconds)

• Anticoagulant treatment

(does <u>not</u> include prophylactic doses of heparin/LMWH or acetylsalicylic acid - aspirin)

• History of liver disease

(portal hypertension OR variceal bleeding OR hepatic encephalopathy OR cirrhosis proven by biopsy, CT or ultrasound)

Hotline: +45 3545 7450 **TURN OVER**

Inclusion criteria

Acute admission to the ICU



Aged 18 years or above

AND

One or more of the following:

Shock

(continuous infusion of vasopressor/inotrope OR MAP < 70 mmHg OR systolic BP < 90 mmHg OR lactate > 4 mmol/l)

Renal replacement therapy

(acute OR chronic)

Invasive mechanical ventilation

(expected duration > 24 hours; If in doubt answer YES)

• Acute or chronic coagulopathy (platelets < 50 OR INR > 1.5 OR PT > 20 seconds)

• Anticoagulant treatment

(does <u>not</u> include prophylactic doses of heparin/LMWH or acetylsalicylic acid - aspirin)

History of liver disease

(portal hypertension OR variceal bleeding OR hepatic encephalopathy OR cirrhosis proven by biopsy, CT or ultrasound)

Hotline: +45 3545 7450 **TURN OVER**



1. Check inclusion criteria:

See front page

2. Go to www.sup-icu.com

Follow the link (eCRF) and start screening Check exclusion criteria when completing screening

3. Prior to randomisation:

- A negative pregnancy test (blood or urine) must be present in all fertile women
- 4. Follow procedure to randomise patient and to obtain a vial identifier number
- 5. The identifier number of the vial allocated to the patient will appear on the screen
- 6. Prescribe the trial medication to the patient once daily
- 7. Administer the medication to the patient

Your personal login:

User: Password:

1. Check inclusion criteria:

See front page

2. Go to www.sup-icu.com

Follow the link (eCRF) and start screening Check exclusion criteria when completing screening

- 3. Prior to randomisation:
 - A negative pregnancy test (blood or urine) must be present in all fertile women
- 4. Follow procedure to randomise patient and to obtain a vial identifier number
- 5. The identifier number of the vial allocated to the patient will appear on the screen
- 6. Prescribe the trial medication to the patient once daily
- 7. Administer the medication to the patient

Your personal login:

User: Password: 1. Check inclusion criteria:

See front page

2. Go to www.sup-icu.com

Follow the link (eCRF) and start screening Check exclusion criteria when completing screening

- 3. Prior to randomisation:
 - A negative pregnancy test (blood or urine) must be present in all fertile women
- 4. Follow procedure to randomise patient and to obtain a vial identifier number
- 5. The identifier number of the vial allocated to the patient will appear on the screen
- 6. Prescribe the trial medication to the patient once daily
- 7. Administer the medication to the patient

Your personal login:

User: Password:

1. Check inclusion criteria:

See front page

2. Go to www.sup-icu.com

Follow the link (eCRF) and start screening Check exclusion criteria when completing screening

- 3. Prior to randomisation:
 - A negative pregnancy test (blood or urine) must be present in all fertile women
- 4. Follow procedure to randomise patient and to obtain a vial identifier number
- 5. The identifier number of the vial allocated to the patient will appear on the screen
- 6. Prescribe the trial medication to the patient once daily
- 7. Administer the medication to the patient

Your personal login:

User: Password: