

BASELINE FORM

General Patient Information

- B1 Sex? Male Female [\[info\]](#)
- B2 Hospital admission date? (dd-mm-yyyy) [\[info\]](#)
- B3 ICU admission date? (dd-mm-yyyy) [\[info\]](#)
- B4 ICU admission time? (hh:mm, 24 hours format) [\[info\]](#)
- B5 ICU admission was directly from the operating or recovery room after surgery? Yes No
- B6 Patient height? cm feet / inches (cm) (feet) (inches) [\[info\]](#)
Choose unit first

Respiratory Support

- B7 Type of closed respiratory support system? Invasive MV NIV or CPAP [\[info\]](#)
- B7a TV_{insp}? (mL) [\[info\]](#)
- B7b PEEP? (cmH₂O) [\[info\]](#)
- B7c P_{peak}? (cmH₂O) [\[info\]](#)
- B7d EPAP or CPAP pressure? (cmH₂O) [\[info\]](#)

Arterial Blood Gas before randomisation

- B8 PaO₂ in the last ABG before randomisation? kPa mmHg [\[info\]](#) Not Available
- B9 SaO₂ in the last ABG before randomisation? (%) [\[info\]](#) Not Available
- B10 p-lactate in the last ABG before randomisation? (mmol/L) [\[info\]](#) Not Available
- B11 FiO₂ at the time of the last ABG? [\[info\]](#)

Acute Illness

- B12 Pneumonia? Yes No [\[info\]](#)
- B13 Multiple trauma? Yes No [\[info\]](#)
- B14 Stroke (haemorrhagic or ischaemic)? Yes No [\[info\]](#)
- B15 Traumatic brain injury? Yes No [\[info\]](#)
- B16 Myocardial infarction? Yes No [\[info\]](#)
- B17 Intestinal ischaemia? Yes No [\[info\]](#)
- B18 ARDS at randomisation? Yes No [\[info\]](#)

SOFA (Sequential Organ Failure Assessment) Score

B19 Lowest Glasgow coma score in the 24 hours prior to randomisation?

 [info]

If sedated, estimate the last score before sedation. If unknown write 15

B20 Lowest MAP in 24 hours prior to randomisation?

 (mmHg) [info]

B21 Use of dobutamine, milrinone and/or levosimendan within 24 hours before the randomisation?

 Yes No [info]

B22 Use of any continuous infusion of vasopressors in the 24 hours prior to randomisation?

 Yes No [info]

B22a Highest dose of dopamine in the 24 hours prior to randomisation?

 ($\mu\text{g}/\text{kg}/\text{min}$) [info]

If dopamine has not been used write 0

B22b Highest dose of norepinephrine (noradrenalin) in the 24 hours prior to randomisation?

 ($\mu\text{g}/\text{kg}/\text{min}$) [info]

If norepinephrine has not been used write 0

B22c Highest dose of epinephrine (adrenalin) in the 24 hours prior to randomisation?

 ($\mu\text{g}/\text{kg}/\text{min}$) [info]

If epinephrine has not been used write 0

B23 Highest concentration of bilirubin in 24 hours prior to randomisation?

 ($\mu\text{mol}/\text{L}$) [info] Not Available

B24 Lowest concentration of platelets in 24 hours prior to randomisation?

 ($10^9/\text{l}$) [info] Not Available

B25 Urinary output in the 24 hours prior to randomisation?

 (ml) [info] Not Available

If urine volume is measured for a short period MULTIPLY TO GET TOTAL OUTPUT IN 24 hours!

B26 Highest creatinine in the 24 hours prior to randomisation?

 ($\mu\text{mol}/\text{l}$) [info] Not Available

Chronic Co-morbidities

B27 History of ischaemic heart disease?

 Yes No [info]

B28 Chronic heart failure?

 Yes No [info]

B29 Active metastatic cancer?

 Yes No [info]

B30 Chronic dialysis?

 Yes No [info]

B30a Habitual creatinine level > 110 $\mu\text{mol}/\text{L}$?

 Yes No [info]

[Return to top](#)

Save

Exit (no save)

