



Review paper

The terrestrial and freshwater invertebrate biodiversity of the archipelagoes of the Barents Sea; Svalbard, Franz Josef Land and Novaya Zemlya



S.J. Coulson ^{a,*}, P. Convey ^b, K. Aakra ^c, L. Aarvik ^d, M.L. Ávila-Jiménez ^a, A. Babenko ^e, E.M. Biersma ^b, S. Boström ^f, J.E. Brittain ^d, A.M. Carlsson ^{a,g}, K. Christoffersen ^h, W.H. De Smet ⁱ, T. Ekrem ^j, A. Fjellberg ^k, L. Füreder ^l, D. Gustafsson ^m, D.J. Gwiazdowicz ⁿ, L.O. Hansen ^d, M. Holmstrup ^o, M. Hullé ^p, Ł. Kaczmarek ^q, M. Kolicka ^q, V. Kuklin ^r, H.-K. Lakka ^s, N. Lebedeva ^t, O. Makarova ^e, K. Maraldo ^u, E. Melekhina ^v, F. Ødegaard ^w, H.E. Pilskog ^{a,x}, J.C. Simon ^p, B. Sohlenius ^f, T. Solhøy ^y, G. Søli ^d, E. Stur ^j, A. Tanasevitch ^z, A. Taskaeva ^v, G. Velle ^{aa}, K. Zawierucha ^q, K. Zmudczyńska-Skarbek ^{ab}

^a Department of Arctic Biology, University Centre in Svalbard, P.O. Box 156, 9171 Longyearbyen, Svalbard, Norway

^b British Antarctic Survey, High Cross, Madingley Road, Cambridge CB3 0ET, UK

^c Midt-Troms Museum, Pb. 1080, Meiereiveien 11, 9050 Storsteinnes, Norway

^d University of Oslo, Natural History Museum, Department of Zoology, P.O. Box 1172 Blindern, NO-0318 Oslo, Norway

^e Institute of Ecology and Evolution, Russian Academy of Sciences, Leninski pr. 33, Moscow 119071, Russia

^f Swedish Museum of Natural History, P.O. Box 50007, SE-104 05 Stockholm, Sweden

^g Lancaster Environment Centre, Lancaster University, Lancaster LA1 4YQ, UK

^h Freshwater Biological Laboratory & Polar Science Center, University of Copenhagen, Universitetsparken 4, DK-2100 Copenhagen Ø, Denmark

ⁱ University of Antwerp, Campus Drei Eiken, ECOBE Department of Biology, Universiteitsplein 1, B-2610 Wilrijk, Belgium

^j Department of Natural History, NTNU University Museum, NO-7491 Trondheim, Norway

^k Mageroveien 168, 3145 Tjøme, Norway

^l Faculty for Biology, Technikerstraße 15, Universität Innsbruck, Innrain, 52, A-6020 Innsbruck, Austria

^m Department of Biology, University of Utah, 257 South 1400 East, Salt Lake City, UT 84112, USA

ⁿ Poznań University of Life Sciences, Department of Forest Protection, Wojska Polskiego 71, 60-625 Poznań, Poland

^o Department of Bioscience, Aarhus University, Vejlsøvej 25, 8600 Silkeborg, Denmark

^p UMR 1349 INRA/Agrocampus, Ouest/Université Rennes 1, Institut de Génétique, Environnement et Protection des Plantes (IGEPP), Domaine de la Motte, 35653 Le Rheu Cedex, France

^q Department of Animal Taxonomy and Ecology, Adam Mickiewicz University, Umultowska 89, 61-614 Poznań, Poland

^r Murmansk Marine Biological Institute, Russian Academy of Sciences, Vladimirskaya St. 17, 183010 Murmansk, Russia

^s Department of Environmental Sciences, University of Helsinki, Niemenkatu 73, 15140 Lahti, Finland

^t Southern Scientific Centre, Russian Academy of Sciences and Azov Branch Kola Scientific Centre, Russian Academy of Sciences, Chekhova 41, Rostov-on-Don 344006, Russia

^u Aarhus University, Department of Agroecology, Blichers Allé, DK-8230 Tjele, Denmark

^v Institute of Biology of Komi Scientific Centre of the Ural Branch of the Russian Academy of Sciences, Kommunisticheskaja, 28, Syktyvkar, Russia

^w Norwegian Institute for Nature Research, P.O. Box 5685 Sluppen, NO-7485 Trondheim, Norway

^x Department of Ecology and Natural Resource Management, Norwegian University of Life Sciences, P.O. Box 5003, NO-1432 Aas, Norway

^y EECRG, Institute for Biology, University of Bergen, P.O. Box 7820, N-5020 Bergen, Norway

^z Centre for Forest Ecology and Production, Russian Academy of Sciences, Profsoyuznaya Str. 84/32, Moscow 117997, Russia

^{aa} Uni Environment, Uni Research, Thormøhlensgate 49b, 5006 Bergen, Norway

^{ab} Department of Vertebrate Ecology and Zoology, University of Gdańsk, Wita Stwosza 59, 80-308 Gdańsk, Poland

* Corresponding author. Tel.: +47 79 02 33 34.

E-mail addresses: Steve.coulson@uni.no, sjcoulson1@gmail.com (S.J. Coulson), pcon@bas.ac.uk (P. Convey), kjetil.aakra@mamu.no (K. Aakra), leif.aarvik@nhm.uio.no (L. Aarvik), mlavilaj@gmail.com (M.L. Ávila-Jiménez), lsd@mail.ru (A. Babenko), elisebiersma@gmail.com (E.M. Biersma), Sven.Bostrom@nrm.se (S. Boström), j.e.brittain@nhm.uio.no (J.E. Brittain), anjamorven@gmail.com (A.M. Carlsson), kchristoffersen@bio.ku.dk (K. Christoffersen), willem.desmet@ua.ac.be (W.H. De Smet), torbjorn.ekrem@ntnu.no (T. Ekrem), arnecoll@gmail.com (A. Fjellberg), Leopold.Fuereder@uibk.ac.at (L. Füreder), kotatsu.no.leo@gmail.com (D. Gustafsson), dagwiazd@au.poznan.pl (D.J. Gwiazdowicz), l.o.hansen@nhm.uio.no (L.O. Hansen), mho@dmu.dk (M. Holmstrup), maurice.hulle@rennes.inra.fr (M. Hullé), kaczmar@amu.edu.pl (Ł. Kaczmarek), malgorzata.kolicka@gmail.com (M. Kolicka), VV_Kuklin@mail.ru (V. Kuklin), hanna-kaisa.lakka@helsinki.fi (H.-K. Lakka), bird_happy@mail.ru (N. Lebedeva), ol_makarova@mail.ru (O. Makarova), kmaraldo@hotmail.com (K. Maraldo), melekhina@ib.komisc.ru (E. Melekhina), frode.odegaard@nina.no (F. Ødegaard), hanne.pilskog@umb.no (H.E. Pilskog), Jean-Christophe.Simon@rennes.inra.fr (J.C. Simon), bjorn.sohlenius@nrm.se (B. Sohlenius), geir.solli@nhm.uio.no (G. Søli), Elisabeth.Stur@ntnu.no (E. Stur), tanasevitch@gmail.com (A. Tanasevitch), taskaeva@ib.komisc.ru (A. Taskaeva), gaute.velle@uni.no (G. Velle), k.p.zawierucha@gmail.com (K. Zawierucha), bioz mud@ug.edu.pl (K. Zmudczyńska-Skarbek).

ARTICLE INFO

Article history:

Received 10 May 2013

Received in revised form

3 September 2013

Accepted 2 October 2013

Available online 16 October 2013

Keywords:

Novaya Zemlya

Frans Josef Land

Spitsbergen

Spitzbergen

Biodiversity

Colonization

Isolation

High Arctic

ABSTRACT

Arctic terrestrial ecosystems are generally considered to be species poor, fragile and often isolated. Nonetheless, their intricate complexity, especially that of the invertebrate component, is beginning to emerge. Attention has become focused on the Arctic both due to the importance of this rapidly changing region for the Earth and also the inherent interest of an extreme and unique environment. The three archipelagoes considered here, Svalbard, Franz Josef Land and Novaya Zemlya, delineate the Barents Sea to the west, north and east. This is a region of convergence for Palearctic and Nearctic faunas re-colonising the Arctic following the retreat of the ice after the Last Glacial Maximum (LGM). Despite the harsh Arctic environment and the short period since deglaciation, the archipelagoes of the Barents Sea are inhabited by diverse invertebrate communities. But there is an obvious imbalance in our knowledge of many taxa of each archipelago, and in our knowledge of many taxa. Research effort in Svalbard is increasing rapidly while there are still few reports, particularly in the western literature, from Franz Josef Land and Novaya Zemlya. Nevertheless, there appears to be a surprising degree of dissimilarity between the invertebrate faunas, possibly reflecting colonization history. We provide a baseline synthesis of the terrestrial and freshwater invertebrate fauna of the Barents Sea archipelagoes, highlighting the taxa present, the characteristic elements of fauna and the complexity of their biogeography. In doing so, we provide a background from which to assess responses to environmental change for a region under increasing international attention from scientific, industrial and political communities as well as non-governmental organizations and the general public.

© 2013 Elsevier Ltd. All rights reserved.

1. Introduction

Arctic terrestrial ecosystems are often considered to be species poor and fragile. The high latitude archipelagoes of the Barents Sea are also isolated due to their geographic separation from Eurasia. Nonetheless, their intricate complexity, especially that of the invertebrate component of their communities, is beginning to emerge. The known terrestrial and freshwater invertebrate fauna of the Svalbard archipelago currently contains over 1000 named species (Coulson and Refseth, 2004; Coulson, 2007a, 2013b).

Investigations of poorly sampled regions within the islands along with studies of genetic diversity, including identification and quantification of cryptic speciation, are likely to lead to considerable increases in invertebrate diversity estimates (Ávila-Jiménez, 2011). The existing species inventories also suffer from taxonomic limitations, in particular relating to unidentified synonymies and misidentifications (Coulson, 2007a; Ávila-Jiménez et al., 2011; Bayartogtokh et al., 2011) and detailed knowledge of the distributions and biogeography of the majority of invertebrate species remains limited. Even in comparatively well-known regions such as western Svalbard, the publication of new species records for the archipelago is frequent, and new taxa continue to be formally described (e.g. Pilskog, 2011; Chaubet et al., 2013; Gwiazdowicz et al., 2012a, 2012b; Kaczmarek et al., 2012b). Just as with the uncertainties applying to Svalbard, the diversity of the Russian archipelagoes of Franz Josef Land and Novaya Zemlya remains understudied, while much of the information that is available is not readily accessible in the western (English language) literature.

It is clear that the invertebrate community plays a central role in many key ecosystem processes, such as nutrient cycling, energy flow, decomposition, herbivory, pollination and parasitism (Petersen and Luxton, 1982; Speight et al., 1999; Bardgett, 2005; Evensen et al., 2005; Ott et al., 2012). However, the relationship between species (alpha) diversity and ecosystem function often remains unclear despite considerable debate around the importance, or otherwise, of 'functional redundancy' in maintaining ecosystem stability (Brussard et al., 2007). Polar (Arctic and Antarctic) ecosystems are considered to be

particularly valuable for studies addressing such fundamental questions of ecosystem function, providing examples across a wide range of levels of assemblage structure (Hodkinson et al., 2003, 2004; Adams et al., 2006; Post et al., 2009). In the context of these ecosystems, the relatively high species-level biodiversity of the terrestrial and freshwater ecosystems of the High Arctic (in comparison, for instance, with those of Antarctic regions; Convey, 2007, 2013) may provide them with a robustness and stability to the characteristically large annual variation in climate and hence also provide resilience to environmental change. Nonetheless, despite this possibly inherent resilience to natural environmental variability, these High Arctic systems may be particularly vulnerable to human disturbance (Jónsdóttir, 2005) predominantly due to lengthy recovery and regeneration times.

Attention has recently become focused on the Arctic due both to the importance of this rapidly changing region and to the inherent interest of an extreme and unique environment. Perhaps nowhere is this more evident than in Svalbard with the establishment of the Kongsfjorden International Research Base (KIRB) at Ny-Ålesund. Nevertheless, despite close to 600 published articles concerning the invertebrate fauna of Svalbard (Coulson, 2007a, 2013a, 2013b), research has largely been fragmented and individual, with little attempt at large scale coordination. Hence there is a disparity in our knowledge between the charismatic and the less studied taxa. The recent publication of species inventories (e.g. Coulson, 2007a; Ávila-Jiménez et al., 2011) have highlighted the Svalbard archipelago as having perhaps the most complete inventory of the invertebrate fauna of any Arctic region (Hodkinson, 2013). Nonetheless, an overall synthesis is lacking, either for Svalbard itself, or for the archipelagoes of the wider Barents Sea region. Now is a particularly opportune moment to provide such a synthesis, with a recent consideration of the Arctic invertebrate fauna calling for the establishment of an inventory of Arctic species as a high priority (Hodkinson, 2013). Moreover, the quantity of invertebrate studies is increasing rapidly, as is the importance of Svalbard as a High Arctic research platform, including the current agenda within Norway to establish the eastern regions of Svalbard as a "reference area for research" (Ministry of Justice and the Police, 2009) and the planned

Svalbard Integrated Arctic Earth Observing System (SIOS) initiative, which forms part of the European Strategy Forum on Research Infrastructures (ESFRI) programme (European Commission, 2012). Currently, there is no overall context into which to set these international initiatives.

This article was catalysed by the expertise brought together for an international workshop on the Terrestrial and Freshwater Invertebrate Fauna of Svalbard held at the University Centre in Svalbard (UNIS) in 2011. We summarize the current state of knowledge of the invertebrate faunas of these archipelagoes, including biodiversity, dispersal, colonization and responses to environmental change. Of the three archipelagoes, by far the most detailed studies of the invertebrate fauna are available for Svalbard. Hence, while we focus primarily on this archipelago, we exploit the opportunity to include, wherever possible, the less well described archipelagoes of Franz Josef Land and Novaya Zemlya.

2. The archipelagoes

The three island groups ringing the Barents Sea consist of Svalbard, Franz Josef Land and Novaya Zemlya (Fig. 1) and comprise a natural geographic unit. This is a region of convergence for the Palearctic and Nearctic biota re-colonising following the retreat of the ice. Svalbard is defined as the land area lying within the coordinates of 10° and 35°E and 74° and 81°N, and consists of five main islands, Spitsbergen, Nordaustlandet, Edgeøya and Barentsøya, and the ‘outlier’ Bjørnøya (Bear Island; Fig. 2). It has a land area of approximately 63,000 km² of which 60% is today permanently covered by ice and snow (Hisdal, 1985). The archipelago is under Norwegian sovereignty but governed by the terms of the “Svalbard Treaty” (Treaty of Spitsbergen, 1920). Novaya Zemlya lies to the north of the Nenetsia Russian coast and is comprised of two principle islands separated by the Matochkin Shar strait, and numerous lesser islands, lying between 70° and 77°N and 51° to 69°E (Fig. 3). The main islands stretch almost 900 km along a north–east axis and is up to 145 km wide (Aleksandrova, 1977) with an area of 81,280 km² of which 27% is currently glaciated (Zeeberg, 2002). During the Cold War, Novaya Zemlya was used as a nuclear test site with the result that for many years it has been a closed military region and thus difficult for biologists to visit (Zeeberg and Forman, 2001). Franz Josef Land lies to the north-east of Svalbard between 79°73' and 81°93'N and 37° and 65°50'E. It consists of approximately 190 largely ice-covered islands forming a total area of 12,334 km², 85% of which is glaciated (Aleksandrova, 1977; Zeeberg and Forman, 2001). As with Novaya Zemlya, Franz




Fig. 2. The Svalbard archipelago with the locations discussed in the text indicated: 1 Ny-Ålesund; 2 Longyearbyen; 3 Barentsburg.

Josef Land was a closed military area for much of the Twentieth Century and access today still requires permission from the Russian authorities, including the Federal Service of National Security and Administration of Reserves and Protected Areas.

The three archipelagoes all have an Arctic climate. The most northerly, Franz Josef Land, has the most extreme climate with mean July (mid-summer) temperature varying between -1.2 and +1.6 °C depending on the specific island considered (Aleksandrova, 1977). Cloudy skies occur approximately 90% of the time, reducing solar heating of the ground (Aleksandrova, 1983). Annual precipitation amounts to 300 mm, most falling as snow (Aleksandrova, 1983).

In Svalbard the annual mean air temperature recorded at the official meteorological station at the airport in Longyearbyen in the west of the archipelago (Fig. 2) is -4.6 °C (mean summer temperature +5.2 °C), with 191 mm annual precipitation for the period 1981–2010 (Førland et al., 2011). Precipitation is particularly variable across this archipelago, decreasing rapidly from the west coast towards the interior. Barentsburg and Isfjord Radio, approximately 50–80 km to the west of Longyearbyen and on the west coast, receive 525 and 480 mm respectively per year (Norwegian Meteorological Institute, 2013). Air temperature is also heavily influenced by the surrounding ocean and in particular the dominant local current systems. To the west, a northwards branch of the North Atlantic Drift carries relatively warm water (> +3 °C; Skogseth et al., 2005), past the archipelago. The east coast, however, is influenced by the cold water of the East Spitsbergen Current carrying polar water south at between 0.5° and -1.0 °C (Skogseth et al., 2005). Hence air temperatures in the north and east of Svalbard are generally lower than in the west. Throughout the archipelago, soils may be snow-covered and frozen for nine months of the year (Coulson et al., 1995).

The latitudinal span of Novaya Zemlya results in a considerable climatic gradient (Zeeberg and Forman, 2001). Annual mean temperature decreases from -5.4 °C on the south-west coast to -10.3 °C at the northern extremity. While winters (December, January) are cold, averaging around -15 °C, the summers are relatively mild with July/August mean air temperature around +6 °C. Annual precipitation also varies, decreasing south to north from 386 mm to 283 mm. However, as with Svalbard, the




Fig. 1. The location of the three archipelagoes surrounding the Barents Sea: Svalbard, Franz Josef Land and Novaya Zemlya.




Fig. 3. Novaya Zemlya with locations discussed in the text indicated: 1 Ivanov Bay; 2 Archangelskaya Bay; 3 Bezymiannaya Bay.

climate of Novaya Zemlya is heavily influenced by the surrounding marine environment, with advected warm North Atlantic water on the west coast while the east coast adjoins the cold Kara Sea which is ice-bound during the winter.

A particular feature of the climate of the High Arctic is the extreme variation in photoperiod. For the settlement of Longyearbyen on Spitsbergen, Svalbard, the sun does not rise above the horizon between October 26 and February 16 (113 days). Conversely, during the period of the midnight sun, from April 19 until August 23 (127 days), the sun remains constantly above the horizon. However, although the sun may be permanently above the horizon from mid-April, the ground is not released from snow and ice until later in the season. For Svalbard this may be mid-June (Coulson, 2013a) and the growing season in vegetated regions, if measured from the approximate period the ground begins to clear of snow until the end of the midnight sun, may be less than 70 days. Some photosynthesis will continue to be possible longer into the autumn but the vascular plants may start to senesce from late July to mid-August (Cooper et al., 2011). For Franz Josef Land the period of the midnight sun is approximately from April 15 until August 24 with polar night extending from October 19 until February 21. With a north–south axis the photoperiod of the islands of the Novaya Zemlya archipelago varies considerably. In the south the period of the midnight sun is only from May 21–July 22 while in the north this period is extended, beginning around April 25 and ending August 17. The polar night is similarly shorter in the south commencing on November 22 with the sun returning on January 20 while in the north the period lasts from October 29 to February 13.

Environmental change is particularly rapid in the Arctic land areas and air temperatures are increasing more rapidly than global means, an example of the 'polar amplification' of the global process (ACIA, 2005; IPCC, 2007). The causes of this fast change are unclear but may be a consequence of general background warming, reduced sea ice cover and changes in oceanic and atmospheric circulation (Serreze et al., 2011). Annual temperatures in Svalbard over the period 1981–2010 have increased by 2.1 °C over the 1961–1990 mean while winter and summer means have increased by 3.4 and 1 °C respectively (Førland et al., 2011). These increases are likely to be linked with variations in atmospheric circulations, with increased frequency of southerly and south-west winds (Hanssen-Bauer and Førland, 1998). Overall annual precipitation has increased marginally with a slight trend towards wetter summers and dryer winters (Førland et al., 2011) also linked to the changes in

atmospheric circulation patterns (Hanssen-Bauer and Førland, 1998). By the end of the current century the average winter temperatures may be up to 10 °C greater than the present normal. Currently, air temperatures fall below –28 °C on approximately three to four days per year. Projections suggest that winter warming by 2050 may result in air temperatures declining to only –23 °C at a similar frequency (Førland et al., 2011). Similar detailed analyses for Franz Josef Land and Novaya Zemlya are not available but it is likely that these will experience similar overall general trends in temperatures and precipitation. However, current scenarios include poor sea ice representation, and recent loss of sea ice may have enhanced regional warming at the same time weakening the accuracy of these projections (Førland et al., 2011).

The history of the Last Glacial Maximum (LGM) in the Barents Sea region is complex but it is clear that Svalbard, Franz Josef Land and much of Novaya Zemlya were largely covered by a dynamic ice sheet (Gataullin et al., 2001; Ingólfsson and Landvik, 2013) becoming exposed progressively as the ice began to retreat. At approximately 14,800 cal yr BP ocean warming commenced at the continental margin off western Svalbard and the western Barents Sea (Hald et al., 1996). The ice sheet started to recede from the marginal coastline of Spitsbergen around 15,800–14,800 cal yr BP (13,000–12,500 ^{14}C yr BP), whereas the central fjord region became ice-free around 11,500–10,800 cal yr BP (Lehman and Forman, 1992; Mangerud et al., 1992). Towards the south, Bjørnøya was deglaciated at around 11,500 cal yr BP (Wohlfarth et al., 1995) and towards the east, Edgeøya, Barentsøya and Franz Josef Land were fully deglaciated at around 11,200 cal yr BP (Landvik et al., 1995; Lubinski et al., 1999). The early Holocene summer temperatures of Spitsbergen were about 2 °C warmer than today (Birks, 1991) causing local cirque glaciers to retreat or disappear in western Svalbard (Svendsen and Mangerud, 1997). These glaciers reappeared from about 4000–3000 cal yr BP during the mid-Holocene cooling and generally advanced towards the Little Ice Age. The environmental conditions have been close to those prevailing today during the last 2500–2000 years with the coldest period occurring during the Little Ice Age (Birks, 1991; Velle et al., 2011). For much of the Holocene, temperatures on Franz Josef Land were 1–4 °C warmer than today with retracted glaciers and snowfields (Lubinski et al., 1999; Forman et al., 2000). Reindeer (*Rangifer tarandus* L., 1758) have been absent in historical time in Franz Josef Land, but antlers dated to 6400–1300 cal yr BP suggest a viable population has existed previously and was possibly driven to extinction during a distinct glacial advance around 1000 cal yr BP (Forman et al., 2000).

Recent studies suggest that large areas of the Amsterdamøya plateau in the north-west of Svalbard remained ice free during the LGM (Landvik et al., 2003) providing possible glacial refugia for invertebrates, and that other regions were also periodically exposed during this period (Ingólfsson and Landvik, 2013). There is, hence, the possibility that some invertebrates survived the LGM *in situ*, but evidence is currently lacking and the predominant view remains that the present fauna is the result of recent immigration since the retreat of the ice. Similarly, it is likely that few, if any, plants survived *in situ* during the LGM (Alsos et al., 2007) although a number of recent studies have hinted at the possible existence of refugia (Westergaard et al., 2011), and current thinking is that flora and fauna of Svalbard is the result of recent immigration.

The relatively short period since deglaciation, combined with the Arctic climate and continuing periglacial soil processes, have strongly influenced habitats and ecosystems. As seen across the Arctic, the environment is characteristically highly heterogeneous with, for example, dry stony ridges, periglacial features, areas of late snow melt, heath or wet moss all in close proximity (Thomas et al., 2008). Large areas have been recently reworked by glacial action

and possess continuous underlying permafrost influencing the soil hydrology. On a regional basis, northern areas consist largely of polar desert characterized by low precipitation and a short snow-free growing season. Vascular plant cover is often limited, restricted to less than 15% in both Svalbard and Franz Josef Land (Aleksandrova, 1983; Jónsdóttir, 2005; Cooper, 2011). Vascular plant diversity totals 74 species in Franz Josef Land (Tkach et al., 2008), 173 in Svalbard (Elven and Elvebakk, 1996) and 216 in Novaya Zemlya (Tkach et al., 2008). Bryophyta (mosses, liverworts and hornworts) form an important component of the environment in the Arctic (Turetsky et al., 2012). In Svalbard there are currently 373 accepted species (Frsvoll and Elvebakk, 1996) while lichens are more speciose, 597 species being recorded (Elvebakk and Hertel, 1996). Recent inventories of the bryophytes or lichens of Novaya Zemlya and Franz Josef Land are not available. Along the west coast of Svalbard and the southern areas of Novaya Zemlya areas of dwarf shrub tundra or heath may develop. Bare soil in all three archipelagoes often possesses a “biological crust” of cyanobacteria, bacteria, algae and lichens.

On a landscape scale the habitat is comprised of a heterogeneous mosaic (Jónsdóttir, 2005). The ridge tops, blown free of winter snow, or areas kept clear of snow by wind eddies, occasionally experience winter temperatures below -30°C while organic soils protected under deeper snow face temperatures no lower than -10°C and often considerably higher (Coulson et al., 1995). Melting snow and permafrost may also provide a constant cold water source throughout the summer resulting in chronically cold, wet and boggy areas in direct proximity to drier polar desert vegetation. The shallow active layer in the permafrost exaggerates this effect by hindering drainage. Soils may also vary considerably in depth and form over short distances. Generally the soils are thin, rarely more than a few centimetres thick, and overlie moraine debris, patterned ground or bedrock. In wetter areas, moss may develop into thick carpets or turfs some tens of centimetres deep, efficiently insulating the ground beneath against insolation (Coulson et al., 1993a). Under bird cliffs significant allochthonous nutrient input may occur. Under little auk (*Alle alle*) colonies in Svalbard, circa 60 tonnes dry matter guano per km^2 may be deposited each season (Stempniewicz et al., 2006). In such nutrient enriched areas, organic soils of over 10 cm depth may also accumulate illustrating the impact of nutrient flow from the marine environment to the often nutrient limited terrestrial habitat (Odasz, 1994). These ornithogenic soils and their associated vegetation (Odasz, 1994; Zmudczyńska et al., 2009; Zwolicki et al., 2013) form a characteristic element of the High Arctic environment (Jónsdóttir, 2005; Zmudczyńska et al., 2012) and one that may be especially vulnerable to the introduction of non-native species (Coulson et al., 2013a).

The physical and chemical properties of Arctic inland waters vary greatly including glacier-fed rivers, snow-melt streams, cold oligotrophic lakes and shallow temporary or permanent ponds. Running freshwaters are characterised by a dominance of glacial meltwater inputs, typically in large braided river systems with high sediment loads, highly irregular flows (even cessation after the main period of snow melt), and very low temperatures even in summer. However, in coastal, glacier-free areas, there are snow-melt and spring-fed streams, as well as lake outflows (Füreder and Brittain, 2006), where conditions can be more favourable, although even here many snowmelt streams dry up in summer. There are also warm springs in two areas in the western part of Spitsbergen that have been the subject of chemical and microbiological studies (Hammer et al., 2005; Jamtveit et al., 2006; Lauritzen and Bottrell, 1994). In Svalbard, river flow may initiate in late June to early July. Ice break-up however occurs later, from mid-July until late-August (Svenning and Gullestad, 2002). The

lakes and ponds in the archipelagoes of the Barents Sea are typically found in coastal, lowland areas as in most other Arctic regions (Bøyum and Kjensmo, 1978; Pienitz et al., 2008; Rautio et al., 2011). Temporary thaw ponds, permanent shallow ponds and small lakes are numerous and, because of the low water depth (usually less than 2 m) or small catchments, these water bodies tend to freeze solid during winter while shallower ones can dry out completely during summer.

Shallow ponds are often hotspots of biodiversity and production for micro-organisms, plants and animals in most Arctic regions (Smol and Douglas, 2007), although containing no fish populations. Nutrient input from grazing geese may be significant (Van Geest et al., 2007). Larger and deeper lakes are also present, although are not as numerous as, for example, in West Greenland and Alaska. Lakes with a water depth of more than 3 m are more stable, not freezing solid or drying out, and can host a permanent fish population. However, the environmental conditions for organisms in High Arctic lakes are different from other northern climatic zones as the ice-free period is very short (typically 1–3 months) (Svenning et al., 2007; Vincent et al., 2008), water temperatures and nutrient concentrations are constantly low and the intensity of ultraviolet radiation is often high compared to more temperate regions. Furthermore, there are physical barriers restricting colonisation such as ice caps or remoteness. As a consequence, the biodiversity of freshwater organisms in still waters in Svalbard and other isolated islands is expected to be low even compared to other High Arctic regions such as West Greenland and Alaska (Gislason, 2005; Samchysyna et al., 2008). Arctic rivers, ponds and lakes have a biocomplexity that resembles that of temperate regions, including phototropic biota (algae and macrophytes), invertebrates (insects, crustaceans and rotifers) and fish, although with much fewer taxa and thus with a simpler food web structure than temperate lakes (Christoffersen et al., 2008).

Set against this environmental background, we here provide a synthesis of the known invertebrate fauna of the terrestrial and limnic environments of the three archipelagoes enclosing the Barents Sea, as a baseline for future ecological studies. Examination of complex ecological linkages is beyond the scope of this review. Nonetheless, we attempt to set each taxonomic group in context and discuss the biodiversity of the islands. In particular, we address the history of research and knowledge development, highlighting gaps in our understanding (which varies considerably between the archipelagoes).

3. The invertebrate fauna

3.1. Rotifera

Studies on the rotifer fauna of Svalbard commenced in the second half of the Nineteenth Century, when von Goes (1862) reported two bdelloid ‘*Callidina*’ species and Ehrenberg (1874) reported *Callidina* (now *Pleuretra*) *alpium* (Ehrenberg, 1853) from moss collected in Spitsbergen. Further early records of the rotifer fauna of terrestrial mosses from Spitsbergen, mainly bdelloids, were provided by Bryce (1897, 1922), Murray (1908) and Summerhayes and Elton (1923). Early planktonic rotifer reports were restricted to monogononts, mostly from Spitsbergen (Svalbard) (Richard, 1898; Olofsson, 1918). In the second half of the Twentieth Century, studies focused on monogononts from the plankton and/or periphyton of Barentsøya (Pejler, 1974; De Smet, 1993), Bjørnøya (De Smet, 1988), Edgeøya (De Smet et al., 1988), Hopen (De Smet, 1990), Nordaustlandet (Thomasson, 1958) and Spitsbergen (Thomasson, 1961; Amréen, 1964a, b, c; Vestby, 1983; De Smet et al., 1987; Kubíček and Terek, 1991; Jørgensen and Eie, 1993; De Smet, 1995; Janiec, 1996; Janiec and Salwicka, 1996). Amréen

(1964a, b) carried out long-term population studies of *Keratella quadrata* (Müller, 1786) and *Polyarthra dolichoptera* (Idelson, 1925) in ponds on Spitsbergen, finding temporal morphological variation in *K. quadrata* and thereby demonstrating that the phenomenon was not limited to low altitudes and latitudes as was previously thought. Interest in bdelloids has recently been revived by Kaya et al. (2010) studying representatives from terrestrial mosses from different localities in Svalbard. Limited physiological studies are available, excepting Opaliński and Klekowski (1989, 1992), who measured oxygen consumption in *Macrotrachela musculosa* (Milne, 1886) and *Trichotria truncata* (Whitelegge, 1889) obtained from Spitsbergen tundra. These studies demonstrated relative temperature independence in the range of 2–6 °C for *M. musculosa* suggesting metabolic cold adaptation. Limited older literature, and no recent studies, are available for Novaya Zemlya (Murray, 1908; Idelson, 1925; Økland, 1928; Gorbunow, 1929; Retowski, 1935) and Franz Josef Land (Murray, 1908; Retowski, 1935).

Of the two major divisions of Rotifera, the Bdelloidea have been largely neglected because of difficulties with identification. Their diversity is underestimated since most studies use animals recovered from rehydrated moss samples, precluding recovery of species lacking, or with poor, capacity to form dormant anhydrobiotic stages. Moreover, as is likely to be the case in many invertebrate groups, recent molecular biological studies have demonstrated that cryptic diversity is high in bdelloids (Fontaneto et al., 2007).

A total of 68 formally identified bdelloid morphospecies have been recorded from the Barents Sea archipelagoes, with around 15% of the current global diversity of Bdelloidea (460 morphospecies distributed over 20 genera; Segers, 2008) being present in Svalbard. These include the majority (85%) of the bdelloids known from the Arctic region (De Smet unpubl.). Virtually all the species reported from these archipelagoes are widespread or cosmopolitan, with *Pleuretra hystrix* Bartos, 1950 being the only Arctic-Alpine endemic. However, the discovery of more endemics may be expected as generalists exhibit the highest cryptic diversity (Fontaneto et al., 2009). Data for Svalbard are only available from the islands of Edgeøya, Prins Karls Forland and Spitsbergen. The known Svalbard fauna comprises 67 morphospecies. Only three and two morphospecies respectively have been reported from Franz Josef Land and Novaya Zemlya. All morphospecies recorded in the Barents Sea archipelagoes occur in limno-terrestrial habitats (mosses, lichens) with 15 also reported from freshwater habitats (permanently submerged vegetation, cryoconite holes).

In this group, older reports are biased in favour of the loricates, a group that includes species with a rigid body wall that fix well and are amenable to microscopic study. Species with a soft integument, the illoricates, contract on fixation and become unrecognizable. Furthermore, re-examination of historical samples (Olofsson, 1918), has shown that loricate diversity per sample was on average 2–4 times higher than in the original publication (De Smet unpubl.). Interpretation of older data may also be compromised due to taxonomic inconsistencies. For example, several monogononts show large phenotypic plasticity, while some taxa originally considered to exhibit wide morphological variation are now recognized to consist of several species. Given these reservations it is impossible to differentiate, for instance, the currently recognised species *Keratella hiemalis* Carlin, 1943, *K. quadrata* (Müller, 1786) and *Keratella testudo* (Ehrenberg, 1832) in earlier reports of '*Anuraea* (*Keratella*) *aculeata*' and its forms in the absence of preserved material. Many monogononts have, again, been shown also to be complexes of cryptic species (e.g. Suatoni et al., 2006).

To date, 163 limno-terrestrial and aquatic monogonont morphospecies have been reported from the Barents Sea archipelagoes, with 134 species from Svalbard, 20 from Franz Josef Land and 71 from Novaya Zemlya. Unequal sampling effort across the different

islands and habitats within the archipelagoes clearly hampers comparison of their rotifer biodiversity. The global diversity of non-marine Monogononta totals approximately 1500 species (Segers, 2008), of which 11% occur in the Barents Sea archipelagoes. In the Arctic region as a whole 327 species are known (De Smet unpubl.) of which 50% have been reported from these archipelagoes. Only 16 species occur occasionally in aerophytic moss with the most frequently found being *Encentrum incisum* Wulfert, 1936, *Lecane arcuata* (Bryce, 1891) and *Lepadella patella* (Müller, 1786). As with the bdelloids, the majority of the monogonont species are cosmopolitan or widespread, although a small proportion show more restricted distributions: the Arctic endemic *Notholca latistyla* (Olofsson, 1918) occurs in all three archipelagoes; *Trichocerca longistyla* (Olofsson, 1918), described from Spitsbergen, is also known from Novaya Zemlya and Swedish Lapland; *Encentrum boreale* Harring and Myers, 1928, *Encentrum dieteri* (De Smet, 1995), *Encentrum murrayi* Bryce, 1922 are currently thought to be endemic to Spitsbergen, and the sub-species *Synchaeta lakowitziana arctica* De Smet, 1988 is restricted to Bjørnøya.

3.2. Gastrotricha

The phylum Gastrotricha is a group of aquatic (freshwater and marine) microinvertebrates. They are a common and important component of the benthic, epibenthic and epiphytic communities in all types of freshwater, brackish water and marine habitats (Balsamo et al., 2008; Todaro and Hummon, 2012; Todaro et al., 2012) and, as a group, considered cosmopolitan (Balsamo et al., 2008).

Arctic Gastrotricha are extremely poorly known. No comprehensive studies have been conducted in the Svalbard archipelago. Scourfield (1897) but De Smet et al. (1987) recorded the genus *Chaetonotus* from Spitsbergen and De Smet (1993) noted that Gastrotricha compose 1–18% of the invertebrate taxa obtained from submerged moss samples from Barentsøya. The taxon has never been studied on Franz Joseph Land or Novaya Zemlya.

In the light of our poor knowledge of Gastrotricha from the Barents Sea region, future studies are likely to find many more species in habitats such as cryoconite holes, raised bogs, water bodies, moist soil, fjords and marine interstitial zones (Valdecasas et al., 2006; Todaro and Hummon, 2012).

3.3. Helminthofauna

3.3.1. Free-living terrestrial and freshwater Nematoda

Despite widespread recognition of the almost ubiquitous presence of nematodes in soil faunas globally and their particular importance in soils of some Antarctic ecosystems where most other invertebrates are poorly or not represented (Freckman and Virginia, 1997; Adams et al., 2006; Maslen and Convey, 2006), this group has received limited attention in the archipelagoes of the Barents Sea and there are no records from Franz Josef Land. The first record of terrestrial nematodes from Svalbard is that of Aurivillius (1883a) who described the new species *Aphelenchus nivalis* (Aurivillius, 1883) found in algae on the snow. Menzel (1920) recorded four species, *A. nivalis*, *Dorylaimus* sp., *Acobeloides biutschlii* (de Man, 1884) Thorne, 1925 and *Plectus cirratus* Bastian, 1865. To date, the only extensive collection of terrestrial nematodes in Svalbard (specifically from Spitsbergen) was carried out by van Rossem in 1965. These samples contained about 75 taxa of which 15 were described as new species (Loof, 1971). Samples collected in the area around Ny-Ålesund by Rudbäck in 1985 were examined in part by Boström (1987, 1988, 1989) resulting in the description of one new species but otherwise mainly corroborating the findings of Loof (1971). Although a few other records are available (e.g., Klekowski

and Opaliński, 1986; Janiec, 1996), the majority of information available on the terrestrial nematode fauna of Svalbard remains that provided by Loof (1971). Checklists of terrestrial and freshwater nematode species found in Svalbard include 95 taxa (Coulson and Refseth, 2004).

The first recorded collections of terrestrial nematodes from Novaya Zemlya are those of Stapfer in 1907 (Steiner, 1916), which included 27 species from 13 genera. More recently, Gagarin (1997a, b, c, 1999, 2000) has described many new species from these islands. In total Gagarin (2001) lists 63 species of terrestrial and freshwater nematodes for the archipelago, although 18 of the species recorded by Steiner (1916) are not included among them. There are 24 species in common between Svalbard and Novaya Zemlya, all taxa which are more or less cosmopolitan.

Free-living terrestrial and freshwater nematodes have been largely omitted from soil ecology studies conducted in Svalbard and hence almost nothing is known concerning their abundance, biomass or ecological or functional importance. In 1994, B. Sohlenius collected samples in Adventdalen and Gluudneset (Kongsfjorden) confirming the presence of high diversities and population densities. The mean population density was 78 nematodes per gram soil dry mass in Adventdalen and 119 nematodes per gram dry mass at Gluudneset (B. Sohlenius unpublished data), values are similar to reports from other Arctic areas. Between 24 and 27 taxa of nematodes were identified. At both sites, the genera *Eudorylaimus*, *Plectus* and *Teratocephalus* were found in all samples examined and were amongst the most abundant taxa. In most samples, Adenophorea bacterial feeders and dorylaims were most abundant. Only very few representatives of obligate plant parasitic nematodes were found. The fauna found thus closely resembles that of other cold areas both in the Arctic (Kuzmin, 1976; Procter, 1977; Sohlenius et al., 1997; Ruess et al., 1999a) and in the sub- and maritime Antarctic (Loof, 1975; Andrassy, 1998; Convey and Wynn-Williams, 2002; Maslen and Convey, 2006).

3.3.2. Animal parasitic taxa

The most detailed investigations of parasitic nematodes in Svalbard are from terrestrial mammals where five species have been identified. Studies have focussed on the parasitic nematodes of the Svalbard reindeer (*R. tarandus platyrhynchus* Vrolik, 1829) and are reviewed by Halvorsen and Bye (1999). The abomasal nematode community consists of three polymorphic species of the order Strongylida, where two dimorphic and one trimorphic species have been identified with major and minor morphotypes. Additionally *Nematodirus* eggs have also been found in faecal samples. The major morphs, *Ostertagia gruehneri* Skrjabin, 1929 and *Marshallagia marshalli* (Ransom, 1907), represent 95% of the parasite population in adult reindeer of both sexes. *O. gruehneri* is host specific to reindeer whilst *M. marshalli* has a wide host and geographical distribution, infecting both bovid and cervid species. It is typically a parasite of cold deserts (Halvorsen, 1986; Halvorsen and Bye, 1999; Irvine et al., 2000). The adult *O. gruehneri* load can reach up to 8000 worms per adult reindeer, while that of *M. marshalli* can exceed 15,000 (Irvine et al., 2001). These nematodes have a direct life cycle in which transmission of the infective stage to the host occurs during grazing. Larvae hatching from the deposited eggs develop into T3 infective stages and infect the next host the following season (Carlsson et al., 2012, 2013). Experimental work has implicated the parasite as a significant factor in regulating population dynamics of Svalbard reindeer through negative effects on fecundity (Irvine et al., 2000; Albon et al., 2002; Stien et al., 2002). As is common for most gut nematodes, *O. gruehneri* is transmitted in the summer when conditions are favourable for survival and development of the free-living stages. Faecal egg densities in the summer vary between 124 and 241 eggs

per gram fresh weight (van der Wal et al., 2000) but no eggs are produced during the winter period (Irvine et al., 2001). Surprisingly, *M. marshalli* is transmitted during the cold period from October to April, which is also when peak egg output occurs at around 8 eggs per gram faecal material (Irvine et al., 2000, 2001, Carlsson et al., 2012, 2013).

Nematodes of the genus *Trichinella* are common throughout the world with the species *Trichinella nativa* Britov and Boev, 1972 being the most common in the Arctic with the polar bear (*Ursus maritimus* Phipps, 1774) as the main reservoir. A recent seroprevalence survey found a higher prevalence of this parasite in the Svalbard region (78%) than in the Barents Sea (east of longitude 30°E) (51%) (Asbakke et al., 2010). Ascaridoid nematodes, likely to be predominantly *Toxascaris leonine* (Linstow, 1902), have been found at a prevalence of 33% in the Arctic fox (*Vulpes lagopus*) (Stien et al., 2010). This is a common parasite of Arctic foxes and has a direct life cycle although it may also use rodents as a paratenic host. Other parasite species found in Arctic foxes from Spitsbergen include cestodes (*Echinococcus multilocularis* Leuckart, 1863, *Taenia crassiceps* (Zeder, 1800), *Taenia polycantha* (Leucart, 1856), *Taenia krabbei* Moniez 1879 and *Diphyllobothrium* sp.) and Ancaanthocephala (Stien et al., 2010). The taeniid tapeworm *E. multilocularis* is sylvatic with foxes comprising the definitive host and the vole *Microtus levis* (initially described as *Microtus rossiaeemeridionalis*) the secondary host. The vole transmitted cestodes, *E. multilocularis*, *T. crassiceps* and *T. polycantha*, decrease in prevalence in the fox population with increasing distance from the intermediate host population (Stien et al., 2010) which is extremely restricted in Svalbard and centered on the abandoned coal mine at Grumont, Isfjord (Henttonen et al., 2001). The local conditions here enable the survival of the vole, but it is thought unlikely to be able to expand its range (Fuglei et al., 2008). *E. multilocularis* is known from Novaya Zemlya (Davidson et al., 2012) but is unlikely to be present in Franz Josef Land due to the lack of intermediate host.

Helminth parasites of the Svalbard reindeer include *Moniezia benedina* Moniez, 1872 and *Taenia ovis krabbei* (Moniez, 1879) Verster, 1969 (Bye, 1985). *M. benedina* is present in around 43% of Svalbard reindeer, a similar level of infection as observed in Greenland (Bye, 1985). *M. benedina* forms a link with the soil microarthropod fauna as oribatid mites comprise the intermediate host. *Taenia ovis krabbei* appears to have large population cycles, with infection rates between 1981 and 1982 decreasing from 61% to 29% (Bye, 1985).

The fauna of parasitic nematodes identified in the seabirds of the Barents Sea archipelagoes consists of predominantly widespread species (Kuklin and Kuklina, 2005). For some (*Anisakis* sp. and *Hysterothylacium aduncum* (Rudolphi, 1802)), birds are not primary hosts but the nematodes may enter together with ingested fish. The first records of parasitic helminths from seabirds in the Barents Sea region were obtained from material collected off the western coast of Svalbard during the Swedish Zoological Expedition of 1900 (Odhner, 1905; Zschokke, 1903). Since then, there have been few studies of the avian helminthofauna of Svalbard (Kuklin et al., 2004; Kuklin and Kuklina, 2005). Markov (1941) published on the helminthofauna of Novaya Zemlya (from Bezymyannaya Bay, on the South Island) (Fig. 3) while Kuklin surveyed the helminth fauna of seabirds from Archangelskaya Bay (North Island) (Kuklin, 2000, 2001). In 1926, Skryabin published an examination of the helminthological collections of the Sedov expeditions to the North Pole (1912–1914) and it is likely that the majority of this material was collected from Franz Josef Land. More recent studies were performed in Franz Josef Land in 1990–93 (Galaktionov and Marasaev, 1992; Galaktionov, 1996).

Throughout the archipelagoes of the Barents Sea, parasitological studies exist from 11 species of seabirds (Markov, 1941;

Galaktionov, 1996; Kuklin, 2001; Kuklin et al., 2004). From these, 47 species of parasitic worm species comprising 10 trematodes, 23 cestodes, 10 nematodes and four acanthocephalans have been identified. A characteristic feature of the helminthofauna of seabirds in Arctic regions, noted for North Island of Novaya Zemlya and in Franz Josef Land (Galaktionov, 1996; Kuklin, 2001), is the extremely low species diversity of the trematode fauna. This is likely due to the lack of intermediate hosts, predominantly littoral molluscs, in Arctic ecosystems (Dunton, 1992) and the extreme climatic conditions preventing completion of the life cycle; primarily by restricting free-swimming larval stages (Baer, 1962; Galaktionov and Bustness, 1999).

Typical of the cestodes from seabirds in the northern archipelagoes is their broad range of host species, for example, *Microsomacanthus diorchis* (Fuhrmann, 1913) (otherwise specific for anatides) and *Arctotaenia tetrabothrioides* (Loenberg, 1890) (previously found only in waders) are recorded parasitizing glaucous gulls (*Larus hyperboreus* Gunnerus, 1767) on Spitsbergen and *Microsomacanthus ductilis* (Linton, 1927) (a widespread parasite of gulls) is found in common eiders (*Somateria mollissima* (L. 1758)) and Brünnich's guillemots (*Uria lomvia* (L. 1758)) in Franz Josef Land (Galaktionov, 1996; Kuklin et al., 2004). This ability is likely to enhance their persistence at the northern boundary of their distribution.

3.4. Oligochaeta

Enchytraeid worms are engaged both directly and indirectly in decomposition processes and nutrient mineralization in the soil (Williams and Griffiths, 1989). Records of Enchytraeidae from Svalbard are to date limited to Spitsbergen and other regions of Svalbard are poorly investigated. Early records from Svalbard include those of Michaelsen (1900), Ude (1902) and Stephenson (1922, 1924, 1925). During the 1990s several locations on Spitsbergen were intensively sampled for enchytraeids (Adventdalen, Bjørndalen, Grumant and Ny-Ålesund), recording 13 species of which two (*Mesenchytraeus argentatus* Nurminen, 1973; *Bryodrilus parvus* Nurminen, 1970) were new to Spitsbergen (Birkemoe and Dozsa-Farkas, 1994; Sømme and Birkemoe, 1997; Birkemoe et al., 2000). In total, 42 species of Enchytraeidae from nine genera have been recorded from Spitsbergen (Nurminen, 1965; Birkemoe and Dozsa-Farkas, 1994; Sømme and Birkemoe, 1997; Birkemoe et al., 2000; Coulson et al., 2013a). Even with the limited sampling available, their diversity in Spitsbergen is high compared to other High Arctic locations, for example north-eastern Greenland and the Arctic archipelagoes of Canada where only 12 and 18 species have so far have been reported respectively (Christensen and Dozsa-Farkas, 2006; Sørensen et al., 2006). All the recorded genera in Spitsbergen are Holarctic, but the common and widely distributed genus *Achaeta* has so far not been recorded in Svalbard or at any other High Arctic location. It is also noteworthy that *Cognettia sphagnetorum* (Vejdovsky, 1878) has only been recorded once from a single location on Spitsbergen despite this species being abundant in cold and wet environments such as heathland, tundra and boreal forest throughout the sub-Arctic (Nurminen, 1966, 1967; Maraldo and Holmstrup, 2010). In general, members of the enchytraeid fauna of Spitsbergen are also found in northern Europe, and it has been suggested that the entire Oligochaeta fauna is of recent origin (Nurminen, 1965; Christensen and Dozsa-Farkas, 2006). No data are available from Franz Josef Land and Novaya Zemlya.

Nurminen (1965) reported the observation of a single damaged and undeterminable lumbricid on Spitsbergen, while Coulson et al. (2013a,b) recently recorded two species, *Dendrodrilus rubidus* (Savigny, 1826) and *Dendrobaena hortensis* (Michaelsen, 1890), in anthropogenic soils below the abandoned cowsheds in

Barentsburg. These latter species appear to have been introduced to Svalbard with imported soils for the greenhouse or fodder and have not been recorded beyond the unusual manure-augmented soils in the town. Lumbricidae have also been observed in Novaya Zemlya where *Dendrobaena octaedra* (Savigny, 1826) is recorded (Stöp-Bowitz, 1969).

3.5. Tardigrada

The Tardigrada is a relatively small group of micrometazoans that contains more than 1167 described species (Degma et al., 2013; Vicente and Bertolani, 2013). Tardigrades are known from almost all ecosystems, from polar and high altitude regions to the tropics on land, and to the abyssal depths in the sea. Terrestrial species are most often encountered in mosses, lichens and liverworts but they can be found also in leaf litter and soil. Freshwater and marine species can be found in sediment, on aquatic plants and sometimes in the pelagic zone. A particular feature of tardigrades is their high tolerance to unfavourable environmental conditions, including desiccation, freezing and radiation stresses, in some cases being able to tolerate exposure to levels of these stresses (such as being submerged in liquid nitrogen, liquid helium or the vacuum of space) that lie well beyond the extreme values ever naturally experienced. They have the ability to enter different types of anabiotic states (anabiosis) in response to these stressors, but they can also survive some extremes in an active state (Weinicz et al., 2011).

Although terrestrial and freshwater Tardigrada have been studied in Arctic regions since the early Twentieth Century only fragmentary and mostly faunistic data are available. The most frequently studied Arctic regions are the Svalbard archipelago and Greenland, but some studies have also addressed Arctic regions of Canada, Jan Mayen, Franz Josef Land and Novaya Zemlya (McInnes, 1994), and Alaska (Johansson et al., 2013). Around 200 terrestrial and freshwater tardigrade species have been recorded from Arctic regions (Pugh and McInnes, 1998).

The first record of terrestrial tardigrades in Svalbard is that of Scourfield (1897) describing the new species *Testechniscus spitsbergensis* (Scourfield, 1897), while Richard (1898) reported the first freshwater tardigrade from Spitsbergen, *Dactylobiotus macronyx* (Dujardin, 1851) (according to Kaczmarek et al. (2008, 2012a) the taxonomic position of this species is uncertain). Increasingly intensive studies were conducted during the Twentieth Century. Early papers of Murray (1907) and Richters (1903, 1904, 1911), were followed by studies from a number of authors (Marcus, 1928; Węglarska, 1965; Binda et al., 1980; Pilato et al., 1982; Dastych, 1983, 1985; Klekowski and Opaliński, 1986, 1989; Pilato and Binda, 1987; De Smet et al., 1987, 1988; Van Rompu and De Smet, 1988, 1991, 1994; De Smet and Van Rompu, 1994; Maucci, 1996; Pugh and McInnes, 1998; Łagisz, 1999; Tumanov, 2006; Smykla et al., 2011; Bernardová and Košnar, 2012; Kaczmarek et al., 2012b; Zawierucha et al., 2013). Most of these studies were limited to reports and descriptions of new species, and only Węglarska (1965), Dastych (1985), Maucci (1996); Pugh and McInnes (1998) and Kaczmarek et al. (2012b) undertook more comprehensive studies, including discussion of ecology, origin of the Arctic Tardigrada, and remarks on taxonomy and zoogeography. The majority of studies have concentrated on the largest island in the archipelago, Spitsbergen, and only De Smet et al. (1988) and Van Rompu and De Smet (1988, 1991, 1994) studied freshwater tardigrades on other islands in the archipelago, including Barentsøya, Bjørnøya, Edgeøya and Hopen (Fig. 2). Across all these studies, 92 tardigrade taxa have been reported although some older reports have not been verified based on modern taxonomy (e.g., Bertolani and Rebecchi, 1993; Claxton, 1998; Michalczyk and Kaczmarek, 2006; Fontoura and Pilato, 2007; Bertolani et al.,

2011; Kaczmarek et al., 2011, 2012b; Michalczuk et al., 2012a,b). Among the species known from this region, 17 were described as new to science and four are currently considered endemic. It is clear that Svalbard has been studied very selectively and a comprehensive study of the entire archipelago is still required.

The tardigrades of Franz Josef Land have been reported only by Murray (1907) and Richters (1911). Murray (1907) reported 21 taxa (19 species and two *varietas*) of which, based on modern taxonomy, 17 species are currently valid. Richters (1911) reported a total of seven taxa (six currently valid species). Therefore, in total, only 19 species are currently known from Franz Josef Land.

Older studies of the tardigrades of Novaya Zemlya are again limited to Murray (1907) and Richters (1911), who reported a total of eight species. Bisarov (1996) published the first modern studies of Tardigrada from Novaya Zemlya, reporting 42 species. Bisarov (1999) then reviewed the available knowledge of Novaya Zemlya tardigrades and also described three new species. Based on all published papers, 81 taxa (68 valid species) are currently known from this archipelago, including one marine taxon, eight marked as "cf.", "gr." or "aff." (uncertain identification) species and four taxa identified only to the genus level.

3.6. Chelicerata

3.6.1. Acari

3.6.1.1. Mesostigmata. The first records of mesostigmatid mites from Svalbard are those of Trouessart (1895), who reported *Urosetius acuminatus* (C.L. Koch, 1847) and *Laelaps* sp. In early publications classifying the natural communities of Svalbard, Summerhayes and Elton (1923, 1928) recorded *Haemogamasus ambulans* Thorell, 1872. Thor (1930) described two genera (*Arctoseius*, *Vitzthumia*) and four species new to science from Svalbard. Unfortunately, the type material has not survived (Winston, 1999) and the original photographic documentation included in the study is inadequate for verification and revision of these species. The status of the type species of the genus *Arctoseius*, *A. laterincus* Thor, 1930, is therefore unclear as this species has not been observed since its initial description although nine other species of *Arctoseius* are now known from the archipelago (Ávila-Jiménez et al., 2011). Lindquist and Makarova (2011) considered that, although the genus *Arctoseius* was established on a presumed monotypy, the type series could include specimens of two (or several) morphologically similar species.

More recent studies have included further descriptions of new species or redescription (Hirschmann, 1966; Petrova and Makarova, 1991; Gwiazdowicz and Rakowski, 2009; Gwiazdowicz et al., 2011a, b; Lindquist and Makarova, 2011), faunistic records (Makarova, 1999, 2000a, 2000c, 2011; Gwiazdowicz and Gulvik, 2008; Gwiazdowicz et al., 2009, 2012a, 2012b; Coulson et al., 2011), and the ecology of the group, especially in soil communities (Byzova et al., 1995; Gwiazdowicz and Coulson, 2011), the specific parasitic complex associated with the introduced vole, *M. levis* (Krumpál et al., 1991) and phoretic associations with Diptera (Gwiazdowicz and Coulson, 2010a).

Twenty-nine species of mesostigmatid mites are currently known from Svalbard, with two apparently restricted to Bjørnøya (Summerhayes and Elton, 1923, 1928; Ávila-Jiménez et al., 2011; Gwiazdowicz et al., 2012a, 2012b; Makarova, 2013b; Coulson et al., 2013b). This diversity is comparable with that of other High Arctic sites such as Ellesmere Island and northern Taymyr (Makarova, 2013a). The majority of these species are characteristic of polar areas, but many (44%) also have European or Holarctic temperate, boreal or polyzonal distributions. Four vertebrate parasitic species are present, usually associated with bird nests or small mammals (Krumpál et al., 1991), and one ectoparasite of birds

(Gwiazdowicz et al., 2012a). Phoresy is also known, for example *Thinoseius spinosus* (Willmann, 1939). This species, usually found on the Holarctic seashore and dispersing on various species of Diptera (Makarova and Böcher, 2009), has been found on the calliphorid fly *Protophormia terraenovae* (Robineau-Desvoidy, 1830) (Gwiazdowicz and Coulson, 2010a).

Along the western coasts of the Svalbard archipelago, which experience a milder climate, a relatively high mesostigmatid diversity is present but, in contrast, in polar desert landscapes only five gamasid species were recorded by Ávila-Jiménez et al. (2011). Population densities on this milder coast of Spitsbergen vary widely between habitats from 20 to 4200 individuals m⁻², with the maximum density recorded being found in mossy vegetation near a colony of little auks (*A. alle*) (Seniczak and Plichta, 1978; Byzova et al., 1995). High density (1000–1840 individuals m⁻²) and species diversity have also been observed at other locations with rich vegetation cover (Byzova et al., 1995; Ávila-Jiménez et al., 2011). Poorly vegetated areas such as saline meadows generally contain fewer species and lower densities (Gwiazdowicz and Coulson, 2011).

There are no detailed investigations of gamasid mites in the Novaya Zemlya archipelago. The first information, based on material of large-scale Arctic expeditions, was published in the late Nineteenth and early Twentieth Centuries (Koch, 1879; Trägårdh, 1904, 1928) and cited only five species. A further nine species were identified during the revision of High Arctic *Arctoseius* species from the collections of V.I. Bulavintsev (Makarova, 2000b, 2000c; Lindquist and Makarova, 2011). Thirteen additional species have been found in samples collected by G.V. Khakhin and S.V. Gor'yachkin. The total number of species of Mesostigmata from Novaya Zemlya now numbers 27, similar number to the diversity on Svalbard (Ávila-Jiménez et al., 2011). Considering the long latitudinal gradient, providing a range of environmental conditions, and the current lack of acarological studies, this number is likely to increase. Eleven species of gamasid are common to both Novaya Zemlya and Svalbard (Makarova, 2009; 2013b). Unlike Svalbard, the South Island of the Novaya Zemlya archipelago was mainly free of ice during the LGM (Velichko, 2002), retaining shrub vegetation (Serebryanny et al., 1998). This, as well as subsequent immigration, may explain the presence of bumble bees, lemmings, and their associated gamasid mite fauna (members of genera *Laelaps*, *Parasitellus*, *Melichares*), in Novaya Zemlya. With the exception of *L. hilaris*, associated with the introduced vole in the derelict mining town of Grumant (Krumpál et al., 1991), these genera are absent in Svalbard (Ávila-Jiménez et al., 2011). In both archipelagoes a third of the gamasid species belong to the genus *Arctoseius*, most of which (61–74%) have Arctic or alpine ranges.

Six species of gamasid mites are recorded from Franz Josef Land (Bulavintsev and Babenko, 1983; Makarova, 1999, 2000c, 2013b), five of which belong to the genus *Arctoseius* and one to *Zercon* (*Z. michaeli* Halaškova, 1977).

3.6.1.2. Ixodida. The bird tick *Ixodes uriae* (White, 1852) is common on sea birds breeding on Bjørnøya but has only recently begun to be observed in large numbers in colonies on Spitsbergen (Coulson et al., 2009). It is unclear why the tick populations in the northern regions of Svalbard are becoming more apparent but a recent study has implicated warmer winters (Descamps, 2013) since the tick overwinters in rock crevices at the nesting sites of its host. *I. uriae* is very widely distributed, circumpolar and bipolar, but recorded only from marine birds and their breeding sites. The species is reported from 52 bird species, the main hosts being auks, tube-nosed sea birds, cormorants, sea gulls and penguins. In the north Atlantic, ticks are most common on guillemots (*Uria aalge* (Pontoppidan, 1763), *U. lomvia*), black guillemot (*Cephus grille* (L.,

1758)), razorbill (*Alca torda* (L. 1758)), puffin (*Fratercula arctica* (L. 1758)) and herring gull (*Larus argentatus* Pontoppidan, 1763) (Mehl and Traavik, 1983).

3.6.1.3. Oribatida. The Oribatida is a suborder of the Sarcoptiformes (Krantz and Walter, 2009). They are often the dominant arthropod group in soil-litter systems, including those of the High Arctic and maritime Antarctic (Block and Convey, 1995; Norton and Behan-Pelletier, 2009). Early records of oribatids from Svalbard date back to Thorell (1871), who described four species new to science of which three, *Diapterobates notatus* (as *Oribata notata*), *Ameronothrus lineatus* (as *Eremaeus lineatus*) and *Hermannia reticulata* are common throughout the archipelago. Thorell also described *Camisia borealis* from the islands, a species which is thought today to be within the variability of *Camisia horrida* (Hermann, 1804) (Seniczak et al., 2006). Following on from Thorell, various reports discussing Oribatida from Svalbard appeared (e.g. Trouessart, 1895; Trägårdh, 1904; Hull, 1922; Summerhayes and Elton, 1923, 1928; Thor, 1930, 1934; Hammer, 1946). Additional reports during the past 50 years (e.g. Forsslund, 1957, 1964; Block, 1966; Karppinen, 1967; Niedbał, 1971; Solhøy, 1976; Seniczak and Plichta, 1978; Byzova et al., 1995) have resulted in a current inventory of 81 species of oribatid mites belonging to 17 superfamilies and 25 families from Svalbard (Bayartogtokh et al., 2011). However, these authors did not include several known representatives of the genera *Brachychthonius*, *Spatiodamaeus*, *Achipteria* (mentioned in Lebedeva et al., 2006); *Gymnodamaeus* and *Microtritria* (in Seniczak and Plichta, 1978) or *Berniniella* sp. (in Coulson, 2007a). With inclusion of these taxa the checklist of oribatid mites of Svalbard includes 87 species from 17 superfamilies and 27 families. However, taxonomic confusion remains a significant problem with the current inventory. For example, the genus *Camisia* requires revision based on modern taxonomic methodologies (Bayartogtokh et al., 2011). For others, the species status is currently being debated, for example Bayartogtokh et al. (2011) regards *Moritzoppia neerlandica* (Oudemans, 1900) and *Oppia translamellata* Willmann, 1923 as the same species (*neerlandica*) while Weigmann (2006) regards them as separate species. Such confusion is mirrored in other species and genera of oribatid mites. Often the specimens originally described or identified no longer exist. A new inventory based on fresh material lodged in appropriate museums is urgently required.

The density of oribatid mites in the Arctic tundra of Svalbard is quite high, often between 9168 and 81,400 individuals m^{-2} (Seniczak and Plichta, 1978; Byzova et al., 1995), comparable with values recorded in the northern tundra of the European part of Russia (Melekhina and Zinovjeva, 2012). These values are also comparable with studies in the maritime Antarctic, where oribatid mites are one of the dominant groups of the terrestrial invertebrate fauna (e.g. Block and Convey, 1995; Convey and Smith, 1997).

Recent work on the oribatids of Svalbard has focused on ornithogenic substrates (Lebedeva and Krivolutsky, 2003; Lebedeva et al., 2006; Pilskog, 2011) and has implicated phoresy with migrating birds as a possible dispersal pathway for soil mites from the mainland to remote Arctic islands and archipelagos (Lebedeva and Lebedev, 2008).

Oribatid mite research commenced in the Russian Arctic in the late Nineteenth to early Twentieth Centuries. The first information concerning the oribatid mites of Novaya Zemlya were published by Koch (1879) who identified and described mites that Nordenskiöld collected during the Swedish Arctic expedition of 1875. L. Koch named seven species of oribatid mites for Novaya Zemlya. He described three species new to science, *Ceratoppia sphaerica* (L. Koch, 1879) (as *Oppia sphaerica*), *Oromurcia lucens* (L. Koch, 1879) (as *Oribata lucens*) and *Platynothrus punctatus* (C. L. Koch, 1839), (as

Nothrus punctatus). Furthermore, he described as new to science the species *Oribata crassipes*. Later Trägårdh (1904) identified this species as the variable species *Notaspis exilis* Nicolet 1855, now transferred to the genus *Zygoribatula*. L. Koch also recorded *A. lineatus* (Thorell, 1871) (as *E. lineatus*), *C. borealis* (Thorell, 1871) (as *Nothrus borealis* (Thorell, 1871)) and *D. notatus* (Thorell, 1871) (as *O. notata*) from Novaya Zemlya. Further information on the oribatid mite of Novaya Zemlya appeared in Trägårdh (1901, 1904, 1928). Based on museum collections of Nordenskiöld's samples, Trägårdh (1904) noted nine species from Novaya Zemlya. However, three of these (*Ameronothrus nigrofemoratus* L. Koch, 1879; *H. reticulata* Thorell, 1871 and *Hermannia scabra* L. (Koch, 1879) Nordenskiöld were collected from the island of Vaigach (Fig. 3) which is not formally part of the Novaya Zemlya archipelago (Koch, 1879). Intensive studies of soil oribatid mites on the islands and archipelagoes of the Russian sector of the Arctic were carried out during 1989–2003. Krivolutsky and Kalyakin (1993) found 23 species of oribatid mites in Novaya Zemlya. Krivolutsky et al. (2003) presented a summary checklist of oribatid mites from the Russian Arctic reporting 58 taxa of oribatid mites, of which 52 were identified to species and six identified to genus from 27 families in Novaya Zemlya. Currently, 64 oribatid mites taxa, of which 58 are identified to species, representing 28 families are known from Novaya Zemlya.

Less is known for Franz Josef Land than from Svalbard or Novaya Zemlya. In his monograph, Trägårdh (1904) recorded two species of oribatid mite from Franz Josef Land: *D. notatus* and *Oribata fischeri* Michael (the current taxonomic status of the latter is unclear). Krivolutsky and Kalyakin (1993) recorded one species of oribatid mite (*Fuscozetes sellnicki* Hammer, 1952) from Franz Josef Land. The 15 taxa now known include nine identified to species and six identified to genus level representing 13 families of oribatid mites (Krivolutsky et al., 2003). Further investigations in Novaya Zemlya and Franz Josef Land will undoubtedly increase the species inventories of these archipelagos.

In the three archipelagos the greatest number of species belong to the families Brachychthoniidae, Camisiidae, Oppidae, Suctobelbidae and Ceratozetidae, as is also seen in the mite communities of the European mainland tundra of the Arctic (Melekhina, 2011). Thirty nine species of oribatid mites are common to both Svalbard and Novaya Zemlya (representing 48% of the 81 species of Svalbard and 67% of the 58 species of Novaya Zemlya). The oribatid mite fauna of Svalbard shows only a low similarity to the fauna of the continental tundra. Of the 81 species of oribatid mites listed from Svalbard by Bayartogtokh et al. (2011), only 36 (44%) were found in the tundra of the Kola Peninsula (Karppinen and Krivolutsky, 1982), although caution must be applied in interpreting these figures given the taxonomic challenges described earlier in this section. Most of the oribatid mites in the three archipelagos are Holarctic and cosmopolitan in distribution. Only a few are restricted to the Arctic, for example *Ceratozetes spitsbergensis* (Thor, 1934), *Svalbardia paludicola* (Thor, 1930), *Autogneta kaisilai* (Karppinen, 1967), *Oribatella arctica* (Thor, 1930), *Iugoribates gracilis* (Sellnick, 1944), and *Trichoribates setiger* (Trägårdh, 1910) from Svalbard, while only two species found in Novaya Zemlya are truly Arctic, *S. paludicola* and *O. arctica*.

3.6.1.4. Other taxa of Acari. Coulson and Refseth (2004) present 32 species names of Trombidiformes (Actinedida) from Svalbard. However, there are no recent published studies of this fauna and the concerns about taxonomic uncertainty expressed for the Oribatida must also be considered here. No information is available from Franz Josef Land and Novaya Zemlya concerning other taxa of Acari.

3.6.2. Araneae

Spiders are major invertebrate predators in virtually all terrestrial ecosystems (with the exception of Antarctica) (Oedekoven and Joern, 2000; Platnick, 2012). They have filled a large spectrum of niches and recent research suggests they may have an important control function on their prey populations. Spiders possess good dispersal abilities and are amongst the first colonisers of new ground revealed by retreating glaciers in Svalbard (Hodkinson et al., 2001). In common with other groups of animals and plants, their diversity generally decreases with latitude and tropical faunas are by far the most diverse. However, one important family, the Linyphiidae (dwarf spiders and sheet-weavers) second only to the jumping spiders (Salticidae) in terms of species numbers (Platnick, 2012), reaches its highest species diversity in the northern region of the Northern Hemisphere (Van Helsdingen, 1984) and attains dominant levels furthest north. The Linyphiidae is also the only family of Araneae represented in the sub-Antarctic islands (Pugh, 1994).

The spider fauna of the Svalbard archipelago is comparatively well known. Holm (1958) provided a review of earlier literature and reported a total of 15 species. Since then only two further species have been reported, *Oreonetides vaginatus* (Thorell, 1872) from the warm spring area in Bockfjorden (Tambis-Lyche, 1967) and *Thanatus formicinus* (Clerck, 1757) from Ny-Ålesund (Aakra and Hauge, 2003). Of this total of 17 species, three are clearly introduced to Svalbard (see Holm, 1958; Aakra and Hauge, 2003) - *Hahnia helveola* Simon, 1875, *Tapinocyba insecta* (L. Koch, 1869) and *T. formicinus*. The 14 naturally occurring species are all Arctic-alpine in distribution and all, except one, belong to the Linyphiidae. The exception, *Micaria constricta* (Emerton, 1882) (previously listed as *M. eltonii* Jackson, 1922; for example by Aakra and Hauge, 2003), belongs to the ground spider family Gnaphosidae. It is so far only known from a few localities around Billefjorden in Spitsbergen. Given the total area of Svalbard, the spider fauna is impoverished, probably a result of both environmental severity and geographic isolation. Most spiders are widely distributed across the archipelago but some have only been found in one or a few localities. Other than *M. constricta*, geographically restricted species include *O. vaginatus*, *Collinsia thulensis* (Jackson, 1924) and *Walckenaeria karpinskii* (O. P. Cambridge, 1873). The most common and widely distributed species, *Collinsia spetsbergensis* (Thorell, 1872), *Erigone arctica palaearctica* Braendegård, 1934, *E. psychrophila* Thorell, 1872, *Hilaria glacialis* (Thorell, 1871) and *Mughiphantes sobrius* (Thorell, 1872), are recorded from all, or most of, the major islands.

The majority of spider species known from Svalbard are also found in northern Fennoscandia and neighbouring parts of Russia, but there are three exceptions, *C. thulensis*, *Hilaira glacialis* (Thorell, 1871) and *M. sobrius* (Thorell, 1872). These are High Arctic species also known from Alaska, Canada and Greenland (*C. thulensis*) and Russia (*H. glacialis* and *M. sobrius*), but not currently from Fennoscandia (see Platnick, 2012). The native species are all found below rocks and in the sparse vegetation cover. One, *O. vaginatus*, may be restricted to warm spring habitats where a more diverse flora and fauna can be found. Although known native diversity in this group is unlikely to increase significantly, there are areas of Svalbard that are insufficiently studied and which may yield new species. As with work on many groups, most investigations have concentrated on the main island, Spitsbergen (see Hauge and Sømme, 1997), and any future studies targeting spider diversity should be focussed on the remaining islands and, in particular, their easternmost parts including Kong Karls Land, Svenskøya and Hopen.

The spider fauna of Novaya Zemlya is also well-studied, comprising 20 species of linyphiids, only eight of which are in common with Svalbard. These shared species are all widespread Arctic species (*Agyneta nigripes* (Simon, 1884), *Collinsia holmgreni*

(Thorell, 1871), *C. spetsbergensis*, *E. arctica palaearctica*, *E. psychrophila*, *E. tirolensis*, *H. glacialis* and *M. sobrius*) (see Tanasevitch, 2012), and are likely to be excellent aerial dispersers. The spider fauna of Novaya Zemlya includes some species near their western limit in Europe and that do not occur on Svalbard, including *Erigone remota* L. Koch, 1869, *Collinsia borea* (L. Koch, 1979), *C. proletaria* (L. Koch, 1879), *Hybauchenidium aquilonare* (Koch, 1879), *Masikia indistincta* (Kulczynski, 1908), *Oreoneta leviceps* (Koch, 1879), *Praestigia groenlandica* Holm, 1967, and *Semljicola arcticus* (see Nentwig et al., 2012). This fauna is clearly strongly influenced by that of the adjacent continental mainland.

In clear contrast with both Svalbard and Novaya Zemlya, only two species of spider have been recorded from Franz Josef Land (Tanasevitch, 2012). These species, *C. spetsbergensis* and *E. psychrophila*, are, as previously mentioned, common and widespread species in the region.

3.7. Hexapoda

3.7.1. Collembola

The first comprehensive collections of Collembola from the European Arctic were those of the Swedish Nordenskiöld expeditions along the north coast of Russia during 1875–1880. The pioneering work of Tullberg (1876) reported 15 species from Novaya Zemlya and five from Svalbard. Prior to that, Boheman (1865) was the first to record a collembolan from Svalbard, “*Podura hyperborea*”, a taxon which has subsequently proved impossible to determine under current taxonomy. Schött (1899) reported four species from Franz Josef Land. Other major works from this initial phase of Arctic exploration include those of Schäffer (1895, 1900), Skorikow (1900) and Lubbock (1898). In the period 1900–1960 the faunistics and biogeography of the Arctic archipelagoes were further elaborated, in particular in the Atlantic sector of the Arctic (Brown, 1936; Carpenter, 1900, 1927; Carpenter and Phillips, 1922; Schött, 1923; Zschokke, 1926; Thor, 1930; Linnaniemi, 1935a, b). Stach (1962) and Valpas (1967) provided good overviews of the Svalbard springtail fauna and Fjellberg (1994) provided the first illustrated identification key to the Collembola species from the Norwegian Arctic islands. A recent inventory of the Svalbard fauna was published by Coulson and Refseth (2004), while Babenko and Fjellberg (2006) provided an extensively referenced catalogue of the Collembola of the whole circumpolar Arctic. From 1960 onwards the focus of research shifted to understanding the ecological functions of soil invertebrates in the Arctic and the physical and genetic mechanisms underlying distributional patterns (Ávila-Jiménez, 2011).

A critical review of published and unpublished species lists from Svalbard results in 68 recognized species including a few probably introduced species. Corresponding numbers from Novaya Zemlya and Franz Josef Land are 53 and 14. Franz Josef Land clearly has a depauperate fauna consisting of mainly circumpolar species. Two of these, *Hypogastrura trybomi* (Schött, 1893) and *Vertagopus brevicaudus* (Carpenter, 1900) are not present in Svalbard although they are known from both the Russian and Canadian sectors of the Arctic. The springtail fauna of Novaya Zemlya has clear affinities to the rich fauna of the northern parts of the Russian mainland. Almost 60% of the species from Novaya Zemlya (33 of the 53 species) are not recorded from Svalbard. These include a large proportion of boreal species which also are not known from Fennoscandia. Similarly, more than 70% of the Svalbard fauna (49 of its 68 species) are not recorded from Novaya Zemlya, illustrating the strong North Atlantic influence on the Svalbard springtail fauna. The proportion of true Arctic (i.e. not recorded from the Fennoscandian mainland) species in Svalbard is low, only 14 of 68 species (21%). Most of these are more or less circumpolar in distribution,

although there is a small but significant group with an eastern Palearctic affinity which appears to show a distribution restricted to the eastern part of Svalbard.

The long history of human presence in Svalbard may have resulted in introduction and subsequent dispersal of new Collembola species. Some of these may have become naturalized to such a degree that their dispersal history is no longer evident. Others may still be present only in their original locations. Recently, five species new to Svalbard were identified in imported soils in the Russian settlement in Barentsburg (Coulson et al., 2013a). One of these, *Deuteraphorura variabilis* (Stach, 1964), is not present in Fennoscandia but is well known as a species associated with human settlements in mainland Europe. This species is also common in several natural northern communities of the European part of Russia, the Karelian coast of the White Sea (Pomorski and Skarzynski, 1995), flood-lands in northern taiga of the Komi Republic (Taskaeva, 2009) and coastal tundra of the same region (Taskaeva and Nakul, 2010). Pomorski and Skarzynski (2001) reported the species as being particularly common in ornithogenic soils of the Karelian coast of the White Sea. Now that it has achieved a foothold on Svalbard, it may have the potential of becoming established as an invasive species in nutrient-enriched soils near seabird colonies. The widespread boreal species *Vertagopus pseudocinereus* Fjellberg, 1975 was originally reported from under bark on imported timber at Ny-Ålesund (Fjellberg, 1975) but is unlikely to become naturalised in Svalbard and has not been recorded since.

Collembola may attain very high population densities. In Svalbard densities of almost 600,000 individuals m⁻² have been reported in enriched moss tundra beneath bird cliffs (Bengtson et al., 1974; Byzova et al., 1995) while in ornithogenic substrates in Novaya Zemlya, Babenko and Bulavintsev (1993) observed densities of 1,200,000 individuals m⁻². With the absence of large detritivores such as earthworms and terrestrial isopods the Collembola may assume a major role in primary decomposition and mineralization of plant material, though their precise contribution is yet to be quantified. The abundance and easy accessibility of surface-active species are exploited by feeding birds such as the purple sandpiper (Bengtson et al., 1975; Leinaas and Ambrose, 1992, 1999).

The very obvious patchiness of habitats and the sharp environmental gradients have been the focus for several studies regarding population dynamics and structure (Birkemoe and Leinaas, 2001; Hertzberg et al., 2000; Coulson et al., 2003a; Ims et al., 2004). Similar characteristics are seen in Antarctic terrestrial habitats (Usher and Booth, 1984, 1986), although Antarctic and even sub-Antarctic collembolan assemblages are much simpler than those of the Arctic with typically only 1–3 species being encountered regularly in any given habitat (e.g. Usher and Booth, 1984; Richard et al., 1994; Greenslade, 1995; Convey and Smith, 1997). Cold adaptation and survival under the harsh environmental stresses has also attracted considerable research (Coulson and Birkemoe, 2000; Coulson et al., 2000; Hodkinson and Bird, 2004). In particular, the initial studies of Holmstrup and Sømm (1998) and Worland et al. (1998) on dehydration and cold hardiness in *Megaphorura arctica* (Tullberg, 1876) (previously *Onychiurus arcticus*) has shed light on the important and previously undescribed survival mechanism of cryoprotective dehydration in Arctic invertebrates (Sørensen and Holmstrup, 2011).

3.7.2. Insecta

3.7.2.1. Phthiraptera. The Phthiraptera (lice) are obligate ectoparasites of birds and mammals. Since they lack a free dispersal stage the Phthiraptera known from any given area are strongly correlated with the available hosts (Clay, 1976; Price et al., 2003). The history of phthirapteran studies on Svalbard is patchy, beginning with Boheman (1865), Giebel (1874), Mjöberg (1910),

Waterston (1922a) and Timmermann (1957), who identified a total of 11 species. The first thorough survey of the Phthiraptera of Svalbard was performed by Hackman and Nyholm (1968) who included 44 species (all from birds). However, many of these were limited to Bjørnøya, were identified to genus level only, or the samples and identifications consisted only of nymphs. Kaisila (1973a) added one species of mammal louse. Mehl et al. (1982) reviewed the species list of avian lice of Svalbard, omitting 19 of Hackman and Nyholm's (1968) records as unidentified or uncertain and adding 11 new records. The number of phthirapteran species recognized from Svalbard currently stands at 37 including two only recorded from Bjørnøya and two subspecies. To this can be added four species recorded by Hackman and Nyholm (1968) that were not determined to species level but which are known from adult individuals that could potentially be reliably determined.

Three suborders of Phthiraptera have been recorded from Svalbard from 22 species of bird and two species of mammal (Kaisila, 1973a; Mehl et al., 1982). The most speciose suborder is the Ischnocera (27 species, two only found on Bjørnøya), while the Amblycera (eight species) and the Anoplura (two species) are less represented. This reflects both the global diversity in each group (Price et al., 2003), and the fact that ischnoceran lice are typically more common on birds than are the amblycerans (e.g. Eveleigh and Threlfall, 1976; Hunter and Colwell, 1994).

The Ischnocera of Svalbard have all been obtained from birds, with most (18 of 27 species) from shorebirds (Charadriiformes). The two most speciose genera on Svalbard are *Saemundssonia* (10 species and two subspecies) and *Quadraceps* (six species), both primarily parasites of shorebirds. Other Ischnoceran genera include *Lunaceps*, *Lagopoecus*, *Perineus* and *Anaticola*.

As with the Ischnocera, the majority of the Amblycera recorded on Svalbard have been obtained from shorebirds (five of eight species). While the genus *Austromenopon* has been recorded from five shorebird species on Svalbard, the quill-boring (Waterston, 1922a) shorebird louse genus *Actornithophilus* has been recorded so far only as nymphs (Hackman and Nyholm, 1968) and the species was omitted from Mehl et al.'s (1982) list. Two amblyceran species have been recorded from the Arctic fulmar (*Fulmarus glacialis* (L., 1761)) and one from two species of geese; barnacle (*Branta leucopsis* (Bechstein, 1803)) and pinkfoot (*Anser brachyrhynchus* Baillon, 1834) (Waterston, 1922a).

Holomenopon and the quill-boring *Actornithophilus* have been implicated in feather loss or "wet-feather" disorder in hosts which may subsequently die from pneumonia (Humphreys, 1975; Taylor, 1981). Hosts infested with these lice may be more likely to die before the parasite can transfer to a new host individual and these louse genera may therefore be missing or rare in the High Arctic. However, more thorough sampling of potential hosts of *Actornithophilus* (shorebirds) and *Holomenopon* (ducks and geese) is required to confirm this.

No Phthiraptera have been recorded from Franz Josef Land. A total of seven have been reported from Novaya Zemlya (Ferris, 1923; Markov, 1937) but there are no recent published records. Of these one is from the Amblycera and the remainder from the Ischnocera. Four of these species have also been recorded from Svalbard.

3.7.2.2. Ephemeroptera, Trichoptera and Plecoptera. No Plecoptera are known from Svalbard or Franz Josef Land, although three species of Plecoptera were recorded from Novaya Zemlya by Morten (1923): *Capnia vidua* (Klapálek, 1904), *C. zaicevi* (Klapálek, 1914) and *Nemoura arctica* Esben-Petersen, 1910. There is only one dubious record of a mayfly (Ephemeroptera) from Svalbard (Jørgensen and Eie, 1993; Coulson and Refseth, 2004; Coulson,

2007a), but *Acentrella laponica* Bengtsson, 1912 has been recorded from Novaya Zemlya (Ulmer, 1925). The circum polar trichopteran, *Apatania zonella* Zetterstedt, 1840 occurs sporadically throughout the western parts of the Svalbard archipelago, as well as on Bjørnøya (Bertram and Lack, 1938) and Novaya Zemlya (Ulmer, 1925). Although mainly found in lakes, *A. zonella* also occurs in and around lake outflows.

3.7.2.3. Hemiptera. Virtually all records of Hemiptera species from the archipelagoes of the Barents Sea are restricted to Svalbard and are exclusively of aphids (Hemiptera: Aphididae). A single published aphid record exists for the South Island (Fig. 3) of the Novaya Zemlya archipelago (*Aphis* (s.l.) sp.) (Økland, 1928). The earliest reports of Svalbard aphids are from Parry's North Pole Expedition (Parry, 1828). However, these reports were of aphid specimens found on pack ice or floating trees and were probably transported by wind, ships or sea currents from distant sources (Elton, 1925a). The first inventory of the aphid fauna from Svalbard (Heikinheimo, 1968) was based on previous published works (Ossiannilsson, 1958) or collections and described "seven or eight species". Two of these were reported as endemic, *Acyrthosiphon calvulus* (Ossiannilsson, 1958) (later revised to *Sitobion calvulum* (Eastop and Blackman, 2005)) and *Acyrthosiphon svalbardicum* Heikinheimo, 1968 (formerly listed as *A. svalbardicus* by Heikinheimo (1968)), one as Arctic (*Pemphigus groenlandicus* (Rübsamer, 1898)), one as boreal (*Cinara abieticola* (Cholodkovsky, 1899)) and four not identified to species level.

In their catalogue of the terrestrial and marine fauna of Svalbard, Coulson and Refseth (2004) listed two resident aphid species (*A. calvulus* and *A. svalbardicum*, and five migrant aphid species (*Aphis borealis* (Curtis, 1828), *Aphis* sp., *Cavariella salicis* (Monell, 1879), *C. abieticola* (Cholodkovsky, 1899) and *P. groenlandicus* Rübsamen, 1898). Finally, Coulson (unpublished data) has located a third resident species in Krossfjord whose identity has not yet been formally confirmed but most likely corresponds to *P. groenlandicus*, a species reported from Iceland, Greenland and the Canadian Arctic (Hille Ris Lambers, 1960; Richards, 1963). Thus, there is clear evidence that at least three aphid species are currently resident on Svalbard: *A. svalbardicum* which appears to feed exclusively on *Dryas octopetala* L. (Strathdee et al., 1993), *S. calvulum* which feeds primarily on *Salix polaris* Wahlenb. but also on *Pedicularis hirsuta* L. (Gillespie et al., 2007) and *Pemphigus* sp. which apparently feeds on roots of *Poa* spp. in Svalbard. Hille Ris Lambers (1952) reports this species feeding on the roots of various Gramineae in Greenland. Other earlier aphid records are unlikely to be resident in Svalbard as they have not been subsequently observed and their host plants generally do not occur. *S. calvulum* is restricted to only few sites on the west coast of Spitsbergen, namely Adventdalen and Colesdalen (Gillespie et al., 2007) and Grøndalen. *A. svalbardicum* is more common along the west coast of Spitsbergen but its spatial distribution is very patchy at the local scale (Strathdee and Bale, 1995; Ávila-Jiménez and Coulson, 2011b); occurrence perhaps being partially determined by winter snow depth modulating the length of the summer growing season (Strathdee et al., 1993; Ávila-Jiménez and Coulson, 2011b). *Pemphigus* sp. feeds on roots and is unlikely to be observed without targeted specialist surveys, and therefore its distribution is likely to be currently underestimated.

Ecological studies on Svalbard aphids commenced in the early 1990s (Strathdee et al., 1993; Gillespie et al., 2007; Hullé et al., 2008; Simon et al., 2008; Ávila-Jiménez and Coulson, 2011b) and have focused on the two resident aphid species, *A. svalbardicum* and *S. calvulum*. These studies have highlighted peculiar traits and life histories thought to result from adaptations and constraints exerted by the harsh conditions of the High Arctic (Table 1). Both species

Table 1

Comparison of fauna and life histories of aphids in Svalbard with those of their temperate counterparts.

Svalbard aphid fauna	Temperate aphid fauna
2 (3) generations a year	12-20 generations per year
apterae; none, or rare, winged forms	Massive production of winged forms (alates)
Cues for wing production unknown	Winged forms induced by crowding
Highly host-specialized species	Larger host spectrum
Obligate holocyclic lifecycle.	Facultative holocyclic life cycle.
Sexual forms produced when 24 h photoperiod	Sexual forms often induced by shortening day length
Biased sex ratios induced by local mate competition	Even sex ratios with rare exceptions

have an extremely reduced life cycle compared to their temperate counterparts. *S. calvulum* displays a two-generation life cycle with a first generation of asexual females hatching from cold-resistant eggs in early June and a second generation of sexual forms that mate and lay eggs before the arrival of frost in early August. *A. svalbardicum* has a similar life cycle but, in some instances, may produce an extra intermediate generation although there are uncertainties whether this is achieved in the field (Strathdee et al., 1993; Hullé et al., 2008). When *A. svalbardicum* displays this three-generation life cycle, the first generation hatching from the overwintering egg produces a mixture of asexual and sexual morphs with the former then generating a third generation exclusively composed of sexual individuals. In field environmental manipulation experiments, the inclusion of the extra generation leads to an order of magnitude increase in the numbers of overwintering eggs (Strathdee et al., 1993, 1995), although the cascade effects of this potential change in primary consumer population density have not been researched there are indications that predator and parasitoid densities may increase (Dollery et al., 2006). In the sexual generations of the two species, the sex ratio is biased towards females as a result of local mate competition (Strathdee et al., 1993; Gillespie et al., 2007). Both species also have reduced dispersal capabilities. *S. calvulum* has no known winged form and its populations occur as small, isolated colonies (Gillespie et al., 2007). Populations of *A. svalbardicum* are also patchily distributed (Strathdee and Bale, 1995) and winged individuals were unknown until the discovery of one alate on Storholmen island (in Kongsfjord close to Ny-Ålesund; Fig. 2) (Hodkinson et al., 2002) and several additional specimens in other areas around Ny-Ålesund (Simon et al., 2008). Whether this apparently recent appearance of small numbers of winged morphs in *A. svalbardicum* results from the recent warming of Svalbard, from other factors that may operate locally and only in certain years, or indeed simply from researchers not previously encountering them, is unclear (Hodkinson et al., 2002; Simon et al., 2008).

Very little is known of the biology of natural enemies of Svalbard aphids. Two newly described parasitoid wasps (Hymenoptera: Braconidae) exploit Svalbard aphids as hosts (Chaubet et al., 2013). *Diaeretellus svalbardicum* Chaubet and Tomanović, 2012 parasitizes exclusively the aphid *A. svalbardicum* and displays a unique case of wing polymorphism with macropterous and micropterous forms in both genders. By contrast, *Aphidius leclanti* Tomanović and Chaubet, 2012 can utilize both aphid species as host. Parasitism rates in field-collected aphids are extremely variable between individuals and collection sites, although can reach up to 50% (Outreman et al., unpublished).

3.7.2.4. Coleoptera. The first report of Coleoptera from Svalbard was of a dead specimen of *Philonthus* collected from under seaweed on a beach by the Swedish polar expedition in 1868 (Holmgren,

1869). In the light of current knowledge of the beetle fauna this specimen is of uncertain origin, although likely originating from ship's ballast (Strand, 1942). In 1882, the first living beetle was reported from Billefjord (Beetlefjord) by Nathorst (1884). Although the material was not collected a new collection was taken in 1898 and *Atheta graminicola* (Gravenhorst, 1806) *Boreophila* (*Atheta*) *subplana* (J. Sahlberg, 1880), and *Isochnus flagellum* (Erichson, 1902) were recorded (Sahlberg, 1901). A review of the Coleoptera from Svalbard was published by Strand (1942), and subsequent additional reports of new species for the archipelago were provided by Strand (1969), Kangas (1967, 1973), Bengtson et al. (1975) and Fjellberg (1983) as well as further information being included in several reviews (Sømme, 1979; Klemetsen et al., 1985; Coulson and Refseth, 2004; Coulson, 2007a).

A total of 19 species of Coleoptera are currently known from Svalbard, including six only recorded from Bjørnøya. However, only 14 of these species have been confirmed to be native to the archipelago. Just *B. subplana*, *A. graminicola* and *I. flagellum* are commonly recorded, whilst most species are found only occasionally. The majority of the species have a wide distribution throughout Arctic regions and none are restricted to Svalbard. Two species, *Coccinella septempunctata* L., 1758 and *Oryzaephilus mercator* (Fauvel, 1889), have only been found inside buildings and are considered to be introduced and, if resident rather than transient, then synanthropic. *Atomaria lewisi* Reitter, 1877 has certainly colonized in recent times and is mainly associated with synanthropic habitats (Ødegaard and Tømmerås, 2000). The single specimen of *Gonioctena* (*Phytodecta*) sp. collected by the Oxford Expedition in 1924 is lost and it is not now possible to confirm its identity although, based on general biogeography, this is most probably *G. arctica* (*affinis*) (Strand, 1942). Only one species of weevil, *I. flagellum* is recorded from Spitsbergen, with the report of *I. foliorum* (*saliceti*) (Coulson and Refseth, 2004) referring to the same species (see Strand, 1942).

In recent times, there have been only two studies that have attempted to search for Coleoptera in Franz Josef Land (Bulavintsev and Babenko, 1983; Bulavintsev, 1999) and, as yet, none have been found. Only a few expeditions have collected Coleoptera from Novaya Zemlya. The Nordenskiöld expedition in 1875 reported nine species (Mäklin, 1881). In 1879 the area was further investigated (Markham, 1881) and in 1897 the Russian entomologist Georgii G. Jacobson spent a summer there. Both expeditions provided new additions to the beetle fauna (Jacobson, 1898; Sahlberg, 1897). By 1910, 16 beetle species were known from Novaya Zemlya of which *Upis ceramboides* (L. 1758) and *Pediacus fuscus* (Erichson, 1845) are considered to be introduced. Poppius (1910) added *Hydroporus acutangulus* (published as *H. sumakowi* Popp.). A major contribution was made by the Norwegian expedition to Novaya Zemlya in 1921 (Münster, 1925). There have been no recent collections or reports of beetles from Novaya Zemlya, excepting Yunakov and Korotyaev's (2007) addition of *Phyllobius pomaceus* (leg. K. Baer) to the species identified from the Russian expedition in 1827.

A number of taxonomic advances have been made since these older collections and publications. The record of *Olophrum boreale* (Paykull, 1792) from Novaya Zemlya (Münster, 1935) is likely to be incorrect. Both Münster (1925) and Poppius (1910) mention the specimen from the island of Vaigatsh published by Mäklin (1881), which may have led to confusion. But, Vaigatsh is not geographically part of Novaya Zemlya (Fig. 3). Finally, according to Poppius (1910) and Münster (1925), *Tachinus apterus* (*Tachinus arcticus*) is found in Novaya Zemlya. *T. arcticus* Motsch, 1860 is now regarded as separate species from *T. apterus* (Ullrich and Campbell, 1974). According to the current distribution of the two species (Ullrich and Campbell, 1974), it is undoubtedly *T. arcticus* occurring in Novaya

Zemlya. Both *Boreostiba frigida* (J. Sahlberg, 1880) and *B. sibirica* (Mäklin, 1880) are recorded from Novaya Zemlya in Mäklin (1881) and Münster (1925). These two species were erroneously synonymised by Löbl and Smetana (2004), but Brundin (1940) showed that these are closely related good species.

In total, and incorporating updated taxonomy, there are 32 species of beetles known from Novaya Zemlya, 28 of which are considered native. Most have a wide distribution in Arctic areas (Münster, 1925), but three are currently reported only from Novaya Zemlya, *Phyllodrepa polaris* (J. Sahlberg, 1897), *Atheta holtedahli* (Münster, 1925) and *Oxypoda oeklandi* (Münster, 1925) (Löbl and Smetana, 2004). Novaya Zemlya has only one species of coleopteran in common with Svalbard, *O. boreale*. But, as previously mentioned, the record *O. boreale* from Novaya Zemlya is probably incorrect.

3.7.2.5. Diptera. Diptera are better adapted to the cold and harsh climate in the Arctic than any other order of insects and comprise an important part of the insect fauna both with regard to species number (e.g. Coulson and Refseth, 2004) and biomass (e.g. Bengtson et al., 1974). Nevertheless, our knowledge of Diptera diversity in the Barents Sea archipelagoes is still insufficient, in particular for the most remote and inaccessible islands such as the Nordaustlandet (Svalbard), Franz Josef Land and Novaya Zemlya.

Within the Barents Sea archipelagoes, the best known and well documented dipteran fauna is that of Svalbard (including Bjørnøya) (Coulson and Refseth, 2004; Coulson, 2007a), including a total of 122 species. Of these, the Chironomidae comprise more than 66 recognised species of which at least four are undescribed (Sæther and Spies, 2012; Ekrem and Stur, unpublished data). Taxonomic confusions endure, for example *Orthocladius mixtus* (Holmgren, 1869) originally described from Svalbard but currently regarded as *nomen dubium*.

Seventeen fly species are known from Bjørnøya, excluding the Chironomidae, which probably are represented by up to 40 species (Ekrem and Stur, unpublished data; Sømme, 1979). Among the non-chironomids four have not been reported from elsewhere in Svalbard including the simuliid *Prosimulium ursinum* (Edwards, 1935) (Edwards, 1935). A similar situation exists for the Chironomidae where certain species are restricted to one or two smaller areas in the Svalbard archipelago. A noteworthy example is *Micropsectra logani* Johannsen, 1928 which is widely distributed in the northern Holarctic and also numerous on Bjørnøya. It is, however, not recorded from the other islands of Svalbard.

The first records of Diptera from Novaya Zemlya are those of Holmgren (1883) collected during Nordenskiöld's expedition. In total, 81 species were recorded, including many new species. Further species were added by the Norwegian Novaya Zemlya Expedition in 1921 (Alexander, 1922; Lenz and Thienemann, 1922; Sack, 1923; Kieffer, 1922, 1923). Since then only scattered records have been published. The most recent list contains 147 species (and subspecies) (Fauna Europaea, 2011) but this is far from complete as several species already reported by Holmgren (1883) are missing (e.g. *Tanytarsus gracilentus* Holmgren, 1883) and additional chironomid taxa have been added (Makarchenko et al., 1998). About 49% of the Diptera species (73 spp.) recorded from Novaya Zemlya are chironomids (Makarchenko et al., 1998; Sæther and Spies, 2012). Due to the region's proximity to the Eurasian continent and its geographic extent, the dipteran fauna of Novaya Zemlya is likely to be the most diverse among the archipelagoes. Nine families recorded here have not been reported from Svalbard, among them 3 families in the superfamily Tipuloidea (Limonidae, Pedicidae, and Tipulidae). The two archipelagoes have only about 30 species of Diptera in common. This disparity probably does reflect true differences, but may in part also be underlain by different taxonomic

traditions between Russian and European dipterists, highlighting the need for taxonomic revision and collaboration.

The Dipteron fauna of Franz Josef Land is very poorly known. [Uspenskiy et al. \(1987\)](#), based on a Russian expedition in 1980–81, mentions five species of Diptera belonging to the Chironomidae and Mycetophilidae (of which the latter probably refers to Sciaridae). Four species are listed in [Fauna Europaea \(2011\)](#), *Hydrobaenus conformis* (Holmgren, 1869), *Ditaeniella grisescens* (Meigen, 1830), *Myenniss octopunctata* (Coquibert, 1798) and *Seioptera vibrans* (L. 1758), of which the latter two are most unlikely to inhabit the islands.

3.7.2.6. Siphonaptera. Two species of flea (Siphonaptera) are present in Svalbard, *Ceratophyllus vagabundus vagabundus* Boheman, 1866 and *Mioctenopsylla arctica arctica* Rothschild, 1922 ([Coulson and Refseth, 2004](#)), both belonging to the Ceratophyllidae. The first record of *C. v. vagabundus* was in 1864 (Boheman, 1865) and the species was later observed in pink-footed geese nests by [Dampf \(1911\)](#). Other studies concerning the fleas of Svalbard include [Thor \(1930\)](#), [Cyprich and Krumpål \(1991\)](#), [Mehl \(1992\)](#), [Coulson et al. \(2009\)](#) and [Pilskog \(2011\)](#). Only one species of Siphonaptera is recorded from Novaya Zemlya, *M. a. arctica*. This species was first described from Novaya Zemlya ([Rothschild, 1922](#)) and later recorded in Svalbard ([Kaisila, 1973a](#); [Coulson et al., 2009](#); [Pilskog, 2011](#)). There appear to be no reports of Siphonaptera from Franz Josef Land.

Ceratophyllus v. vagabundus has a northern Holarctic distribution and is common on members of the bird families Anatidae and Laridae and their predators ([Brinck-Lindroth and Smit, 2007](#)). In Svalbard it is recorded as an ectoparasite of the common eider duck (*S. mollissima*), barnacle goose (*B. leucopsis*), pink-footed goose (*A. brachyrhynchus*) and glaucous gull (*L. hyperboreus*) ([Dampf, 1911](#); [Pilskog, 2011](#)) and has also been recorded in nests of snow bunting (*Plectrophenax nivalis* (L., 1758)) ([Pilskog, 2011](#)). As *C. v. vagabundus* is a generalist that uses hosts belonging to different families of birds ([Tripet et al., 2002](#); [Brinck-Lindroth and Smit, 2007](#)) further studies are likely to increase the list of host species present in Svalbard. The second species, *M. a. arctica*, is also known from northern Norway (including Jan Mayen), Iceland, and Alaska ([Mehl, 1992](#); [Brinck-Lindroth and Smit, 2007](#)). This species currently has two subspecies, *M. a. arctica* and *M. a. hadweni* Ewing, 1927. However, although only *M. a. arctica* is recorded as present in Svalbard, it is possible that the sub-specific division is not valid. *Mioctenopsylla a. arctica* is a host-specific flea only present on black-legged kittiwakes (*Rissa tridactyla* (L., 1758)) in Svalbard and, with the exception of [Coulson et al. \(2009\)](#), all records have been obtained from black-legged kittiwake plumage and nests ([Kaisila, 1973a](#); [Cyprich and Krumpål, 1991](#); [Mehl, 1992](#); [Pilskog, 2011](#)) or in the immediate vicinity of their colonies ([Hågvar, 1971](#)). The finding of adult *M. a. arctica* in nests of common eider duck and glaucous gull in Kongsfjorden in Svalbard by [Coulson et al. \(2009\)](#) was probably a misidentification, as this species was not found by [Pilskog \(2011\)](#) in a more thorough investigation of the common eider duck nests in the same area. The effect the fleas have on the host birds is unknown but high flea infestations may generally reduce breeding success in some species of bird including geese breeding in the Arctic such as Ross's, *Chen rossii* (Cassin, 1861) and lesser snow geese, *Chen caerulescens caerulescens* (L., 1758) ([Harriman and Alisauskas, 2010](#)).

Bird fleas spend most of their lives in the nests of their host where they feed on adult birds and chicks ([Lewis and Stone, 2001](#)). High densities of adult fleas and juvenile stages can be present in bird nests in Svalbard ([Cyprich and Krumpål, 1991](#); [Mehl, 1992](#); [Pilskog, 2011](#)), often being the numerically dominant arthropods in the nests of common eider duck, barnacle goose, black-legged kittiwake and glaucous gull breeding in the Kongsfjord area, Svalbard ([Pilskog, 2011](#)). Although the bird fleas are known to bite

humans ([Mehl, 1992](#)) no fleas have been reported from mammals in Svalbard.

3.7.2.7. Lepidoptera. Twenty-three species of Lepidoptera have been recorded from Svalbard and Novaya Zemlya, seven of which (30%) are considered to be vagrants and not resident in the archipelagoes. No Lepidoptera have been recorded from Franz Josef Land. [Kaisila \(1973b\)](#) summarized the Lepidoptera from Svalbard reporting six species, four of which were considered to be resident. With recent additions ([Sendstad et al., 1976](#); [Laasonen, 1985](#); [Coulson, 2007a](#)) the total observed in Svalbard, including accidental migrants, has risen to 10 species, but with no increase in the number of resident species. The resident species total now is considered to be three; *Plutella polaris* Zeller, 1880 ([Bengtsson and Johansson, 2011](#)) (Plutellidae), *Pyla fusca* (Haworth, 1811) (Pyralidae) ([Coulson et al., 2003b](#)) and *Apamea exulis* (Lefèbvre, 1836) (Noctuidae) ([Rebel, 1925](#); [Alendal et al., 1980](#); [Hodkinson, 2004](#)). [Kaisila \(1973b\)](#) also considered *Plutella xylostella* (L., 1758) as resident. However, while this cosmopolitan and migratory species often disperses in great numbers, and has been recorded on several occasions in the Arctic (and likewise in the Southern Hemisphere ([Convey, 2005](#))), it is unlikely that it can overwinter in the archipelago. The closely related *P. polaris* is a distinct species so far only known from Svalbard ([Bengtsson and Johansson, 2011](#)). It is unclear why this species has not been observed since it was first recorded, but the type material of *P. polaris* is held in the Natural History Museum, London and was studied by [Baraniak \(2007\)](#) who drew wings and male genitalia. The distinct features currently support the specific status of *P. polaris*. Ideally, molecular studies would be required to confirm the relationship between these two species. *A. exulis* has been recorded from Svalbard under three different species names, *A. exulis*, *A. maillardi* and *A. zeta*, and this has caused some confusion. According to current taxonomy, *A. maillardi* and *A. zeta* are both species from mountainous regions in southern and central Europe and do not occur at more northern latitudes ([Zilli et al., 2009](#)). *P. fusca* was recorded from Svalbard in 1974 ([Aagaard et al., 1975](#)) and 2002 ([Coulson et al., 2003b](#)). The old record of *Pempelia dilutella* (Denis and Schiffermüller, 1775) ([Elton, 1925b](#)) probably refers to *P. fusca*. The latter species is clearly able to maintain populations in Arctic environments as it is also present in Greenland, Labrador and Alaska ([Kaisila, 1973b](#)). *P. fusca* is a polyphagous species; *S. polaris* and *S. reticulata* (L.) are indicated as possible food plants in Svalbard ([Coulson et al., 2003b](#)).

Lepidoptera recorded from the Swedish Nordenskiöld expedition to Novaya Zemlya were published by [Aurivillius \(1883b\)](#) and those of the Norwegian expedition in 1921 by [Rebel \(1923\)](#). Of the 15 species recorded from Novaya Zemlya only one species, *P. xylostella*, is considered an immigrant resulting in a resident total of 14. Moreover, *P. xylostella* is the only lepidopteran species that Novaya Zemlya and Svalbard have in common and is also the only species of Lepidoptera recorded from Bjørnøya ([Lack, 1933](#); [Sømme, 1979](#)) but is again unlikely to be resident (although, note the caveat mentioned above with reference to the separation of this species from *P. polaris*). The lepidopteran fauna of Novaya Zemlya is composed mainly of species with broad circumpolar Arctic distributions. However, the record of *Argyrolepte mengelana* (Fernald, 1894) (Tortricidae) in Novaya Zemlya is the only observation of this species so far from the Eurasian continent. This species is otherwise known from Greenland, Canada (North West Territory, Yukon), and Alaska ([Jalava and Miller, 1998](#)) and *Glacies coracina* (Esper, 1796) (Geometridae) is known only from the Palaearctic, and is distributed from Fennoscandia to Japan ([Skou, 1984](#)).

3.7.2.8. Hymenoptera. The Hymenoptera is one of the most speciose orders of insects. The majority of species are parasitoids,

attacking a wide variety of insects and other invertebrates. Where there are possible hosts present there are usually hymenopterans and they may occur even in the harshest climate. Nonetheless, it is notable that no species are associated with the two resident Diptera or microarthropods of the Antarctic Peninsula and that very few species are known from the sub-Antarctic islands, both of which have climates less extreme than those of the Barents Sea archipelagoes (Greenslade, 2006; Gressitt, 1970; Convey, 2013).

A total of 39 species of Hymenoptera are currently recorded from Svalbard (Waterston, 1922b; Yu et al., 2005; Coulson and Refseth, 2004; Coulson, 2007a; Jong, 2011). The majority are parasitoids belonging to the families Ichneumonidae (22 species) and Braconidae (five species) in the suborder Apocrita. In addition, the Symphyta is represented by seven species of Tenthredinidae. Braconids are known to parasitise the two Svalbard endemic aphid species.

Novaya Zemlya has 40 species of hymenopteran recorded, probably reflecting low collecting activity given the archipelago's sizeable land area and the close proximity to the continental mainland. The Swedish Nordenskjöld expedition (Holmgren, 1883) and the Norwegian Novaya Zemlya expedition (Friese, 1923) were of great importance in investigating the hymenopteran fauna of this archipelago. Most of the recorded species again belong to the families Ichneumonidae (20 species) and Braconidae (four species). Overall, there are few hymenopteran species shared between Svalbard and Novaya Zemlya, which may support different underlying immigration patterns. Three species of bumblebee are also present (Holmgren, 1883; Friese, 1923), a family not resident in Svalbard. The honey bee, *Apis mellifera* L., 1758 has been reported from all three archipelagoes (Jong, 2011) as an accidental migrant. No hymenopterans have yet been reported from Franz Josef Land, although since some vascular plants (e.g. *S. polaris*) and associated insects are present (Hanssen and Lid, 1932; Jong, 2011) it is plausible that they may occur.

3.8. Freshwater ecosystems

In polar regions freshwater ecosystems are intimately linked with their catchments. Perhaps here more than anywhere else there is a gradation, or grey area, between truly terrestrial and truly limnetic ecosystems. The underlying permafrost results in considerable surface flow during the spring melt (Pienitz et al., 2008) enhancing linkages and resulting in substantial nutrient input to freshwater systems from the surrounding terrestrial terrain (Van Geest et al., 2007; Rautio et al., 2011) and freshwater habitats are traditionally considered along with the terrestrial in polar regions.

3.8.1. Biodiversity and ecosystem function in ponds and lakes

Investigations of freshwater invertebrates on the major islands of the Barents Sea date back more than a hundred years to pioneers such as Bryce (1897), Scourfield (1897) and Olofsson (1918). Summerhayes and Elton (1923) visited Bjørnøya and Spitsbergen in 1921 and sampled ponds and lakes while Økland (1928) reported on species distribution from a Norwegian expedition to Novaya Zemlya in 1921. More recent investigations in Svalbard have typically been carried out in areas close to established research stations on Spitsbergen in Isfjorden (Colesdalen and Kapp Linné), Kongsfjorden (Ny-Ålesund and Brøggerhalvøya), Hornsund and Mosselbukta (Halvorsen and Gullestad, 1976; Husmann et al., 1978; Jørgensen and Eie, 1993; Janiec, 1996), and Bjørnøya (Koch and Meijering, 1985). The branchiopod fauna of Novaya Zemlya is summarized by Vekhoff (1997). Information on the freshwater crustacean fauna of the Franz Joseph Land archipelago is exceedingly scarce and primarily based on a single report from Scott (1899). Apart from this area there is a fairly good understanding of the biodiversity of some organisms (crustaceans and fish);

however, knowledge of microscopic groups such as protozoans is less developed (e.g. Opravilova, 1989; Beyens and Chardez, 1995; De Jonckheere, 2006). Comparison of different Arctic regions based on crustacean species richness (Gíslason, 2005; Samchyshyna et al., 2008) indicates that glaciation history has played an important role in determining community diversity.

The list of Rotifera (Section 3.1) and crustacean species from the Barents Sea archipelagoes is diverse. All of these are currently thought to be circumpolar and the communities do not differ greatly from sub-Arctic regions in Europe, Russia or North America (Ghilarov, 1967; Samchyshyna et al., 2008). The zooplankton species distribution resembles that of Greenland and Alaska, with dominance by cladoceran over copepod species. Several calanoid copepod species (e.g. *Eurytemora raboti* Richard, 1897; *Limnocalanus marcus* G.O. Sars, 1863) are widely distributed in the lakes of Novaya Zemlya and Svalbard (Olofsson, 1918; Halvorsen and Gullestad, 1976; Vekhoff, 1997).

The large branchiopods living in the Barents Sea region occupy the most extreme aquatic environments in Arctic regions (Vekhoff, 1997). Vekhoff (1997) lists four species of Anostraca (*Polyartemia forcipata* (S. Fischer), *Artemiopsis bungei plovomornini* (Jaschnov, 1925), *Branchinecta paludosa* (Gajl, 1933), and *Branchinectella media* (Schmankewitsch, 1873)) and two species of Spinicaudata, *Caenestheria propinquua* (Sars, 1901) and *C. sahlbergi* (Simon, 1886), in addition to *Lepidurus arcticus* (Pallas, 1793) (Branchiopoda, Notostraca) at Novaya Zemlya. It is notable that the northern-most known occurrence of *B. paludosa* is at Ivanov Bay (77°N) in the Novaya Zemlya archipelago (Fig. 3, Vekhoff, 1997). *L. arcticus* frequently occupies shallow freshwater lakes and ponds with no fish population (Jeppeesen et al., 2001) but may exceptionally co-occur with fish in some deep lakes, in shallow cold lakes or in lakes with refugia from fish at the southern-most edges of its distribution range in sub-Arctic regions of mainland Norway and in Iceland (Primicerio and Klemetsen, 1999; Woods, 2011). *L. arcticus* has been recorded in multiple sites on Spitsbergen, Bjørnøya, Novaya Zemlya and Franz Josef Land (Olofsson, 1918; Janiec, 1996; Vekhoff, 1997 (and references therein); Hessen et al., 2004). The crustacean can utilize different habitats in sub-Arctic and Arctic regions including shallow near-shore habitats in Svalbard (Lakka, 2013) and deeper regions of lakes on mainland Norway (Sømme, 1934). Food web studies in Bjørnøya have shown that environmental contaminants can enter the Arctic aquatic food web and that *L. arcticus*, chironomids and Arctic charr can contain elevated levels of both PCBs and DDT (Evenset et al., 2005). *L. arcticus* seems sensitive to various environmental disturbances and therefore can be used as an indicator species of ongoing environmental change in the Arctic and sub-Arctic (Lakka, 2013).

Bottom-dwelling macroinvertebrate species belonging to Nematoda, Oligochaeta, Ostracoda, Hydracarina, Chironomidae, and Trichoptera have been reported in several studies (Summerhayes and Elton, 1923; Jørgensen and Eie, 1993; Janiec, 1996) but there is no detailed information on the biology of the groups. The chironomid diversity is substantial (Styczynski and Rakusa-Suszczewski, 1963; Hirvenoja, 1967; Section 3.7.2.5).

Five species of cestode are known to parasitize the Arctic char (*Salvelinus alpinus* (L., 1758)) in Svalbard. Two of these, *Eubothrium salvelini* (Schrank, 1790) and *Proteocephalus exiguous* (Swiderski and Subilia, 1978), utilize Arctic char as their final host, whereas *Diphyllothrium ditremum* (Creplin, 1825) employs various fish-eating birds as the definite host which, in Svalbard, is likely to be the red-throated diver (*Gavia stellata* (Pontoppidan, 1763)) (Hammar, 2000). Additional groups known to parasitize Arctic char in Svalbard include one species of nematode (*Philonema oncorhynchii* Kuitunen-Ekbaum, 1933) and a copepod (*Salmoncola*

edwardsii Olsson, 1869; Siphonostomatoida) (Kennedy, 1978; Sobecka and Piasecki, 1993).

Studies of food web structure in lakes and ponds are limited, but a number of recent experimental studies have focused on nutrient addition to lakes and ponds mediated by geese (Van Geest et al., 2007), the role of dissolved organic carbon for microbial communities (Hessen et al., 2004), the implications of UV radiation on plankton growth (Van Donk et al., 2001) and the dynamics of microbial communities (Ellis-Evans et al., 2001; Laybourn-Parry and Marshall, 2003). Such studies are important in order to understand the complexity of Arctic aquatic ecosystems and to be able to predict effects of human activities and environmental change (Prowse et al., 2006). Furthermore, van der Wal and Hessen (2009) have highlighted important analogies between aquatic and terrestrial food webs in the High Arctic, as a result of harsh conditions leading to grazer dominated food web dynamics.

3.8.2. Ecosystem function in streams and rivers

Biodiversity in running waters in Svalbard is low, as is probably also the case in Franz Josef Land, although there is little information on the latter. Freshwater biodiversity is however, higher in Novaya Zemlya due to its proximity to the mainland and its more southerly location. Colonisation by freshwater invertebrate fauna is limited by the isolation of the archipelagoes (Gíslason, 2005). In addition, the short summer season and the cessation of flow in most river systems during the long winter render environmental conditions unsuitable for many taxa.

Despite their wide distribution, there have been few ecological studies of Svalbard streams and rivers compared to terrestrial or even lake systems and almost none from Novaya Zemlya or Franz Josef Land. Studies of hydrological and chemical processes, especially in glacier-fed systems are, however, more common (e.g. Gokhman, 1988; Hagen and Lefauconnier, 1995; Killingtveit et al., 2003; Krawczyk and Pettersson, 2007). The significance of microbial activity for nutrient processes in glacial meltwater has also been highlighted from Svalbard studies (Hodson et al., 2008) and there have been studies of freshwater algae and cyanobacteria in the vicinity of Ny-Ålesund (Kim et al., 2011).

Freshwater invertebrate species records derive from both early expeditions and more recent collecting trips (e.g. Morten, 1923; Ulmer, 1925; Bertram and Lack, 1938), or from studies of the aerial insect fauna (Hodkinson et al., 1996; Coulson et al., 2003b). These records are frequently based on collections of adults, mainly chironomids, making it difficult to assign them to the larval environment – terrestrial, wetlands, lakes or streams. The invertebrate fauna of streams and rivers is dominated by chironomids, especially Diamesinae, although Nematoda, Enchytraeidae and Tardigrada have also been recorded from freshwater habitats in Svalbard (Styczynski and Rakusa-Suszczewski, 1963; Hirvenoja, 1967; Janiec, 1996; Coulson and Refseth, 2004). Planktonic and benthic crustaceans can also be found drifting downstream of lakes (Maiolini et al., 2006).

In recent years there has been an increasing focus towards understanding the influence of hydrological processes on stream fauna (ecohydrology). Studies of the influence of water source on benthic stream communities have been undertaken in Svalbard over the last 10–15 years (Brittain and Milner, 2001) demonstrating the importance of channel stability and water temperature in structuring benthic invertebrate communities (Castella et al., 2001; Lods-Crozet et al., 2001; Milner et al., 2001). These studies have focused on two contrasting rivers in Svalbard in the vicinity of Ny-Ålesund, Bayelva and Londonelva. These rivers have been monitored for discharge, sediment transport and water temperature for over 20 years (Bogen and Bønsnes, 2003; Brittain et al., 2009). Bayelva is a glacier-fed river, whereas Londonelva is fed by rain and snowmelt. This difference in water source gives rise to distinct differences in their

chironomid faunas, with higher densities in Londonelva, a greater proportion of Orthocladiinae and different species of *Diamesa* (Diamesinae) (Lods-Crozet et al., 2007). In general Chironomidae (especially the genus *Diamesa*) dominate in glacial systems whereas in non-glacial systems their relative abundance decreases and the subfamily Orthocladiinae as well as other taxa including Oligochaeta, Copepoda, Acari, Collembola and Tardigrada become more frequent (Fürer and Brittain, 2006). However, most species are similar to the nearby sub-Arctic areas as the coastal regions of the Barents Sea or more temperate areas. Studies in a wider range of streams (Fürer and Brittain, 2006) have shown that species number, abundance and food web complexity follow a gradient with regard to catchment characteristics such extent of glacier cover and the extent of nutrient input from bird cliffs or upstream lakes. Furthermore, a recent study of geothermal streams on Iceland (Woodward et al., 2010) demonstrated that water temperature is a key parameter among the factors directly affecting community structure and trophic interactions.

Invertebrate drift is generally a widespread and important phenomenon in running waters, and this is again the case on Svalbard. Studies during the Arctic summer in a stream near Ny-Ålesund (Maiolini et al., 2006; Marziali et al., 2009) showed that drift rates can be high and that there are distinct diurnal patterns, even in continuous daylight, which are controlled by environmental variables such as water temperature and discharge rate. Drift rates were enhanced by artificial shading of the stream, indicating a strong behavioural component. Invertebrate drift from streams and glacial outlet rivers, contributes a significant source of food for seabirds and waders (Mehlum, 1984). It is clear that freshwaters on Svalbard are an important link for nutrients and biota between terrestrial, estuarine and marine ecosystems.

4. Adaptation to conditions – ecophysiology and life histories

The climates of all three archipelagoes are characterized by low precipitation, subzero temperatures for most of the year, and only a short summer season allowing the growth and reproduction of invertebrates. The low winter air temperatures (monthly means of -10 to -15 °C for at least 6 months, and much lower extreme minima) combined with permafrost and shallow depth of snow pose a significant challenge to the invertebrates, because thermally buffered microhabitats are often not available above or in the soil (Coulson et al., 1995). Clearly, the species occurring in these archipelagoes have appropriate ecophysiological and more general life history adaptations to their harsh conditions, and these have formed a focus of polar invertebrate research generally and that in Svalbard specifically.

Two primary cold tolerance strategies are widely used by Arctic invertebrates. Freeze-tolerant animals have the capacity to survive ice formation in extracellular body fluid compartments whereas freeze-avoiding species possess physiological mechanisms that promote extensive supercooling of body fluids throughout the winter (for reviews of, and an introduction to, the biology of extreme environments and the wider cold tolerance literature see Zachariassen, 1985; Sømme, 1999; Wharton, 2002; Thomas et al., 2008; Ávila-Jiménez et al., 2010; Denlinger and Lee, 2010; Bell, 2012). These two main strategies for survival of extreme conditions ensure that body water is more or less conserved during winter, either trapped as ice (in freeze-tolerant species) or because typical freeze-avoiding species often have a relatively impermeable cuticle that limits evaporative water loss.

Many soil and freshwater invertebrates such as tardigrades, nematodes, enchytraeids, prostigmatid mites and Collembola are often of small size (<5 mm length) and have little resistance to evaporative water loss through their cuticle (Harrisson et al., 1991;

(Convey et al., 2003). At the same time, groups such as nematodes, annelids and tardigrades, which are active within the surface layer of water on soil particles and in moss/peat are also susceptible to inoculative spreading of ice to body fluids when the soil or sediment water that they are in contact with freezes, meaning that freeze-avoidance by supercooling is not possible (e.g. Wharton, 1986, 2002; Convey and Worland, 2000). Thus, such invertebrates have only two options: survive freezing of body fluids or avoid freezing by other means than supercooling (Pedersen and Holmstrup, 2003). Encasement in air spaces in frozen soil or sediment may lead to desiccation of small species with low resistance to water loss, as water inevitably transfers from the liquid state within the animal's body to the ice crystals surrounding it (Danks, 1971; Holmstrup and Westh, 1994). A few invertebrates have taken advantage of this process, developing a third strategy, termed cryoprotective dehydration, driven by differences in water vapour pressure between the unfrozen body fluids and surrounding ice (Worland et al., 1998; Holmstrup et al., 2002; Sørensen and Holmstrup, 2011).

Many Arctic invertebrates, due to the short growing season, show extended development, and often Arctic populations have life cycles of two or more years whereas the same or closely related species in temperate regions have annual life cycles or more than one generation each year (Danks, 1992; Strathdee and Bale, 1998). Thus, Collembola, enchytraeids and Acari from Svalbard may have two-year life cycles or longer (Birkemoe and Sømme, 1998; Birkemoe and Leinaas, 1999; Birkemoe et al., 2000; Søvik, 2004). These life cycles may become closely adapted to, and synchronised with, the local environmental conditions. For example, chironomids may have sufficient life cycle flexibility to permit one or two periods of adult emergence each summer, probably depending on temperature conditions (Hodkinson et al., 1996). One striking example is the Svalbard endemic aphid, *A. svalbardicum* (see Section 3.7.2.3) which has a highly modified programmed life cycle (Strathdee et al., 1993, 1995, Table 1).

5. Paleocommunities – trends of the past

Relatively few Late Quaternary and Holocene palaeozoological studies have been performed in freshwater or terrestrial environments in Svalbard and to our knowledge such studies are lacking in Franz Josef Land and Novaya Zemlya. The oldest terrestrial sub-fossils from Svalbard are recorded from Visdalens (Edgeøya) and dated to $14,700 \pm 500$ cal yr BP (Bennike and Hedenas, 1995), suggesting very early post-glacial colonization or perhaps the presence of glacial refugia (rapidity of colonisation being consistent with local refugia, cf. Convey et al., 2008). The assemblage includes *L. arcticus*, *Candonia* sp. (Crustacea, Podocopida) and a questionable Lepidoptera. Several other taxa are recorded from Visdalens during the early Holocene, including Oribatida, Chironomidae, a questionable Ichneumonidae, *O. boreale*, *Daphnia pulex* (L., 1758) and *Erigone* sp. (Bennike and Hedenas, 1995). The presence of *Lepidurus*, *Daphnia* and *Candonia* suggests that mesotrophic ponds existed in the area. The staphylinid beetle *O. boreale* has also been recorded from Early Holocene lake sediments on Bjørnøya (Wohlfarth et al., 1995) together with the beetles *Agabus bipustulatus* (L., 1767) and *Eucnecosum tenuis* (LeConte, 1863). The only Trichoptera in the palaeoecological record, noted as Limnephilidae indet, was also found in the Early Holocene sediments of Bjørnøya, as well as *Lepidurus* sp. and an unidentified Hymenoptera (Wohlfarth et al., 1995). In addition to the abovementioned studies, rotifer resting eggs and testate amoeba have been retrieved from sediments in Kongressvatn (Grønfjord) on Spitsbergen and Rosenbergdalen on Edgeøya, respectively (Beyens and Chardez, 1987; Guizzoni et al., 2006).

Remains of Chironomidae and Cladocera have received the greatest attention in palaeozoological studies from Svalbard. Unidentified chironomids have been recorded from Bjørnøya (Wohlfarth et al., 1995) and Edgeøya (Bennike and Hedenas, 1995), while studies from Nordaustlandet (Luoto et al., 2011) and from five lakes on Spitsbergen (Brooks and Birks, 2004; Fadnes, 2010; Velle et al., 2011) included detailed identifications and environmental interpretations based on the chironomid assemblages. These records typically include about 10 taxa and show large among-site differences in species assemblages. Most likely, some sites experienced nutrient enrichment from bird guano or proximity to the sea, whereas others were influenced by glacial melt-water. In a survey of chironomid sub-fossils retrieved from the upper 1 cm of sediment (representing about 25 years) from 23 western Svalbard lakes, 18 taxa were found. The abundance and distribution of these taxa were primarily influenced by pH, nutrient concentrations, water temperature and water depth (Brooks and Birks, 2004).

Cladocera sub-fossils have been retrieved from lake sediments in Kongressvatn and in the Hornsund areas of Spitsbergen (Guizzoni et al., 2006; Zawisza and Szeroczyńska, 2011), in Visdalens on Edgeøya (Bennike and Hedenas, 1995), and in Lake Einstaken on Nordaustlandet (Luoto et al., 2011; Nevalainen et al., 2012). The sub-fossil Cladocera assemblages often have a low diversity compared to contemporary assemblages, although this may be the result of physical and chemical processes influencing the preservation of the remains in sediments, such as bottom water freezing during winter (Sywula et al., 1994; Zawisza and Szeroczyńska, 2011).

6. Invertebrate immigration, dispersal and biogeography in the archipelagoes of the Barents Sea

Some areas of the archipelagoes of the Barents Sea were ice free during parts of the last glaciation, including nunataks above 300 m altitude in northwest Svalbard (Landvik et al., 2003), low lying areas along the west coast of Spitsbergen and Prins Karls Forland down at sea level (Andersson et al., 2000; Ingólfsson and Landvik, 2013), and substantial parts of Novaya Zemlya (Mangerud et al., 2008). Nunataks have been proposed to act as refugia for some crustaceans with the ability to survive as relicts due to their hardy resting eggs (Samchyshyna et al., 2008). However, most biota could not survive on nunataks (Brochmann et al., 2003; Schneeweiss and Schönswitter, 2011) due to the prevailing polar desert conditions in the ice free areas (Andersson et al., 2000). These harsh conditions and the general observation that a relatively limited number of species currently occur on nunataks is consistent with the *tabula rasa* hypothesis; that is, that few if any plants or animals survived in Svalbard during the LGM and that the communities observed today are the result of recent immigration after the retreat of the ice. For example, molecular studies have indicated that plant diversity in the Arctic is the result of glaciation cycles combined with subsequent dispersal barriers (Eidesen et al., 2013). Furthermore, species richness is often found to be lower in areas that are known to have been covered by ice sheets during the last glaciation, suggesting that dispersal limitation has been a key factor structuring many contemporary communities in the Arctic (Samchyshyna et al., 2008; Strecker et al., 2008; Ávila-Jiménez and Coulson, 2011a). However, local microclimatic and microhabitat conditions vary widely on small spatial scales, as do species distributions, and survival in small but particularly benign ice-free refugia at either low or higher altitudes cannot automatically be discounted (Landvik et al., 2003; Paus et al., 2006; Skrede et al., 2006; Westergaard et al., 2011). Notwithstanding this, the contemporary invertebrate fauna is currently thought to be primarily the result of recent immigration and colonization processes. Pugh and

McInnes (1998) suggested that the biogeography of Tardigrada in the Arctic can be explained by colonization from a Nearctic source following the retreat of the ice. Similarly, the community structure of Collembola throughout the Arctic appears to be the result of colonization from numerous source populations outside of the Arctic with subsequent dispersal within the Arctic (Ávila-Jiménez and Coulson, 2011a, Fig. 4) and Arctic plant communities are considered to have been selected for species with high dispersability by the repeated cycle of glaciation in the Arctic (Alsos et al., 2007). Parts of the South Island, Novaya Zemlya (Fig. 3), were ice-free, with shrub vegetation surviving throughout the last glaciation (Serebryanny et al., 1998; Velichko, 2002; Mangerud et al., 2008), providing source populations for the colonization of other islands in the archipelago as the ice retreated. With the existence of widespread plant refugia in Novaya Zemlya, and the putative presence of plant refugia and/or deglaciated areas in Svalbard, it is highly likely that invertebrate faunas also existed in these refugia. Studies from Antarctica have demonstrated that even in the most climatically extreme and isolated ice-free areas there is a viable, if limited, terrestrial fauna (Convey, 2013). But, although a glacial refugium has been proposed for certain freshwater species such as the *D. pulex* complex in the Canadian High Arctic archipelago (Weider and Hobæk, 2000), no evidence of *in situ* faunal survival has yet been described for Svalbard or Franz Josef Land. Increasingly, molecular and bioinformatic analytical techniques devoted to defining biogeographic and phylogeographic patterns are being applied to studies in the polar regions (Weider et al., 1999; Marková et al., 2013). These approaches permit more accurate definition of the timing of divergence events, both between species and between populations within species, potentially allowing detailed descriptions of dispersal and colonization patterns (Allegrucci et al., 2006; Stevens, 2006; Stevens et al., 2006, 2007; McGaughan et al., 2010; Mortimer et al., 2011). Their application has led to a paradigm shift in the interpretation of the antiquity of the contemporary Antarctic terrestrial biota (Convey and Stevens, 2007; Convey et al., 2008, 2009; Vyverman et al., 2010). However, as yet these approaches have not been applied to the study of Arctic terrestrial invertebrates, and have so far generally focused on floral biogeography (Abbott and Brochmann, 2003; Brochmann et al., 2003; Alsos et al., 2007; Ávila-Jiménez, 2011).

Several dispersal vectors have been suggested for invertebrate species colonizing the polar regions. Airborne dispersal by active flight may account for many winged species. Chernov and Makarova (2008) consider the Coleoptera fauna of Svalbard to consist of flighted migratory species. Passive dispersal with air currents (anemochory) may be also responsible for many of the

species or taxa seen in the islands, for example Tardigrada, Aphidiidae, Syrphidae, Tipulidae and Lepidoptera (Elton, 1925a, 1934; Kaisila, 1973b; Pugh and McInnes, 1998; Coulson et al., 2002b). Similarly, passive dispersal by ocean currents (hydrochory), either floating on the ocean surface or rafting with floating debris of terrestrial or marine origin, such as tree trunks, seaweed rafts, or human rubbish may account for the arrival of others (Coulson et al., 2002a). Further species may hitch with migratory birds or mammals (zoochory). Lebedeva and Lebedev (2008) speculated on the possible role of birds in transporting soil microarthropods to the Arctic, although clear confirmation of the occurrence of this process is lacking. Non-parasitic mites have also been described as phoretic on larger invertebrate species such as Diptera (Coulson, 2009; Gwiazdowicz and Coulson, 2010b). Transport assisted by human processes (anthropochory) may be an increasingly common immigration route. This is especially the case with plants, where around 100 vascular plant species are now known to have been introduced to Svalbard via human activity compared to the natural flora of 164 species (Alsos et al., 2013). The effect of human-mediated dispersal on invertebrate immigration patterns has not been quantified in the High Arctic, although it is recognised as a factor far outweighing natural dispersal events in the Antarctic (Frenot et al., 2005) where it has also been highlighted as a major threat to biodiversity (Hughes and Convey, 2010, 2012; Chown et al., 2012a, 2012b; Greenslade and Convey, 2012). In the anthropogenic soils of the mining town of Barentsburg (Svalbard), 11 of the 46 identified invertebrate species (24%) were non-native (Coulson et al., 2013a, 2013b). Svalbard may be particularly vulnerable to anthropogenic introduction of alien species due to the high volume of visitors arriving both by ship and aeroplane (Ware et al., 2011). In contrast, access to Franz Josef Land and Novaya Zemlya is currently more restricted, albeit after a long history of military usage with, presumably, little or no attention to biosecurity issues.

A range of synanthropic species have also been described from the Svalbard archipelago in human settlements (Coulson, 2007b) which are, in the main, unlikely to establish in the natural environment due to the Arctic conditions. However, as is characteristic of human introductions elsewhere, and in particular in the Antarctic (Frenot et al., 2005; Greenslade et al., 2012), a proportion of such species are likely to be able to survive in the natural environment and subsequently become invasive. Furthermore, the majority of invertebrate fauna are cryptic and require specialist expertise for recognition and the probability of successful remedial extermination once establishment has occurred is likely to be low (see Hughes and Convey, 2012 for discussion of these issues in a parallel Antarctic context).

Most terrestrial invertebrate biogeographic studies carried out to date in Arctic areas are based on community assemblages and have examined groups such as Collembola (Hågvar, 2010; Ávila-Jiménez and Coulson, 2011a, Fig. 4), Tardigrada (Pugh and McInnes, 1998), or Rotifera (Gíslason, 2005). For many groups meaningful comparisons of the invertebrate communities between the archipelagoes are not possible due primarily to lack of sampling effort and taxonomic confusion. However, for some groups it is feasible to make an overall assessment of similarities (Table 2). Within data limitations it is notable that, for many groups, the species diversities of Svalbard and Novaya Zemlya are numerically similar, but that they have few, or very few, species in common indicating limited connectivity between the archipelagoes.

7. Environmental change

The archipelagoes of the Barents Sea lie in the High Arctic region that is expected to be particularly sensitive to oceanographic and




Fig. 4. Dispersal routes suggested to and within the Arctic archipelagoes in the Barents Sea. Solid arrows indicate dispersal directions for Collembola species (modified from Ávila-Jiménez and Coulson, 2011a). Undulating arrow indicates a link with the Nearctic region suggested for the Tardigrada (Pugh and McInnes, 1998).

Table 2

Similarities between the invertebrate faunas of the archipelagoes. Figures indicate: total number of species in common (total number of species in first archipelago; total number of species in second archipelago). Only species considered resident are included. Dashes indicate comparisons not possible, usually as no species of the group concerned have been recorded from Franz Josef Land.

Group		Novaya Zemlya to Svalbard	Franz Josef Land to Svalbard	Franz Josef Land to Novaya Zemlya
Rotifera	Bdelloidea	1 (2:67)	3 (3:67)	0 (0:2)
	Monogononta	45 (71:134)	16 (20:134)	15 (20:71)
Gastropoda		0 (0:1)	—	—
Nematoda	Freeliving	24 (81:95)	—	—
Annelida	Lumbricidae	0 (1:2)	—	—
Tardigrada		40 (68:92)	17 (19:92)	12 (19:68)
Acari	Mesostigmata	11 (27:29)	3 (6:29)	4 (6:27)
	Oribatida	39 (64:87)	5 (15:87)	8 (15:64)
Araneae		8 (20:14)	2 (2:14)	2 (2:20)
Collembola		20 (53:68)	12 (14:68)	8 (14:53)
Insecta	Phthiraptera	4 (7:37)	—	—
	Hemiptera	0 (1:3)	—	—
	Coleoptera	1 (28:14)	—	—
	Diptera	29 (150:122)	1 (4:122)	0 (4:150)
	Chironomidae	19 (73:66)	1 (1:66)	0 (1:73)
	Other Diptera	10 (77:56)	0 (3:56)	0 (3:77)
	Siphonaptera	1 (1:2)	—	—
	Lepidoptera	0 (14:3)	—	—
Crustacea	Cladocera	5 (8:17)	—	—
	Copepoda	1 (16:6)	—	—
	Anostraca	0 (4 : 0)	—	—
	Ostracoda	0 (5:2)	—	—
	Notostraca	1 (1:1)	—	—
	Malacostraca	0 (3:1)	—	—

climatic changes, and a strong indicator of their biological consequences (ACIA, 2005; Chapin III et al., 2005; Convey et al., 2012). Svalbard, and even Novaya Zemlya, are subject to warm North Atlantic influences from the west, and cold Arctic Ocean influences from the east, as well as lying at the boundary of the region experiencing large scale changes in winter and multi-year Arctic sea ice extent (Serreze et al., 2007). All three archipelagoes lie at the high latitudes subject to the ‘polar amplification’ of general global climate trends, although Svalbard is the only location of the three archipelagoes considered here to have a detailed publicly accessible long term meteorological record by which to confirm recent warming trends (Førland et al., 2011). Increasingly sophisticated general circulation models continue to predict considerable further warming over the next century in the high latitude polar regions (IPCC, 2007). Temperature warming is accompanied by a suite of other changes of biological relevance, including in the form and amount of precipitation, cloudiness, humidity and insolation, and the timing and frequency of freeze-thaw events. Finally, although the Arctic does not normally experience the organized formation of a seasonal ozone hole as is seen in the Antarctic, intermittent and significant depletion does occur spatially at Arctic latitudes throughout the Arctic summer, with a number of potential biological impacts identified (e.g. Rozema, 1999).

The general biological responses to environmental change in the Arctic have received considerable attention (e.g. for review see Callaghan et al., 2004a, 2004b; Chapin III et al., 2005; AMAP, 2011). However, studies on the impacts of climate change on soil animal communities in High Arctic environments are limited. Although environmental manipulation methodologies have been applied widely in the context of ITEX studies to a range of Arctic vegetation habitats, generally these studies have focussed on vegetation responses and have not addressed, or included, the soil or other elements of the invertebrate fauna. Studies of soil nematode communities at Abisko, Sweden, have indicated that while

population densities are increased, biodiversity is generally affected negatively and distinct changes in trophic structure are caused by environmental perturbations (Ruess et al., 1999a). This seems to be an indirect effect of changes in vegetation cover, plant species composition, litter quality and below-ground input by plants, which in turn will have a major impact on nutrient turnover through microorganisms and soil fauna (Ruess et al., 1999b; Sohlenius and Boström, 1999; Simmons et al., 2009). Similar initial responses to manipulations have also been reported in Antarctic studies, which also identified that caution needs to be used in separating initial, and sometimes drastic artefactual changes, in population density and diversity from those that appear to become established after longer periods of manipulation have permitted the impacted communities to stabilise (Convey and Wynn-Williams, 2002).

Webb et al. (1998), in a three year open-topped chamber manipulation at Ny-Ålesund, found very little change in soil oribatid mite community composition, although noting possible subtle changes in species relative abundances. These authors concluded that the soil microhabitat would be more buffered from short-term changes in temperature than would be the case for invertebrates of the overlying vegetation. This difference is perhaps illustrated by the striking findings of Strathdee et al. (1993), who reported an order of magnitude increase in overwintering aphid eggs within versus outside chamber manipulated vegetation, indicating a possible step change in the population dynamics of this species under realistic warming scenarios. However, as noted above, a similar response has not been observed in recent studies of natural aphid populations in areas that are thought to have warmed already by a similar amount in recent decades.

In general terms, the two most important environmental variables subject to change in Arctic (and Antarctic) terrestrial ecosystems of relevance to the invertebrate fauna are those relating to temperature and the availability of liquid water. While water may provide the primary limiting factor to the temporal activity of invertebrates in these ecosystems, temperature provides the energy required to fuel biological processes. In many instances, where climate change leads to relaxation of the constraints provided by either or both of these variables, the invertebrate biota are likely to benefit, with expectation of increased production, biomass, population size, community complexity, and colonisation (Convey, 2011; Nielsen et al., 2011; Nielsen and Wall, 2013). However, in terms of biodiversity, these positive impacts of climate change may then be outweighed by other impacts of human activities, in particular the establishment of invasive non-indigenous species.

More broadly, anthropogenic climate change poses a serious threat to freshwater ecosystems in Barents Sea region. Widely reported reductions in sea ice have been mirrored in freshwater systems. For example, an extended ice free period has resulted in higher water temperatures and lower water levels in Kongressvatnet in Svalbard (Holm et al., 2011). Elevated snow fall may increase the opacity of translucent block-ice delaying the start of primary production in the spring (Svenning et al., 2007). Recently, lakes on granitic bed rock appear to have become more acid, perhaps due to increased acid precipitation, a spring influx of low pH water during the melt and the low buffering capacity of granitic rocks (Betts-Piper et al., 2004).

It is important to recognize that increased temperature due to global warming may induce a multitude of changes in detail in the High Arctic environment, in addition to the broad generalizations described above. Included amongst these are increased snow depth, earlier snow melt and more frequent freeze/thaw cycles in winter (Christensen et al., 2007; AMAP, 2011; Wilson et al., 2013). In particular, the presence of a solid ice cover directly on the soil surface may seriously affect the Collembola and presumably other

communities (Coulson et al., 2000). Changes in local faunal composition are likely to occur under current warming scenarios, but over the short to medium term (years to decades) the Svalbard environment probably has sufficient buffer capacity to offer suitable habitats for even the most cold adapted species. In terms of biodiversity conservation, special attention should be given to monitoring the status of species which are absent from Arctic continental mainland landmasses, as these may be the first to be pushed towards extinction.

8. Conclusions and future research priorities

The archipelagoes of the Barents Sea are inhabited by diverse communities of invertebrates, despite the short period since deglaciation and the clear environmental challenges. There is an obvious imbalance in our understanding of the biodiversity of the three archipelagoes. Research in Svalbard is increasing rapidly while there are still few reports, particularly in the western literature, from Franz Josef Land and Novaya Zemlya. Our knowledge of the faunas of all three archipelagoes is relatively recent, the majority of records commencing in the early Twentieth Century.

In attempting to describe or compare the invertebrate fauna of the archipelagoes of the Barents Sea it is immediately clear from the consideration of all taxa here that great problems exist that challenge our understanding of the region. First, there is the lack of comprehensive sampling campaigns. Many locations have only been sampled on one occasion, sampling locations were often selected primarily due to logistical considerations and sampling frequently carried out by non-specialists, and often a limited range of taxa were focused on driven by the skills and interests of the particular taxonomists/ecologists associated with the sampling programme. There is a strong need for repeated sampling campaigns designed to capture seasonal and interannual variation in the Barents Sea region. For Novaya Zemlya and Franz Josef Land there has been the added problem of access to a closed military region. Hence, we often have a very prejudiced knowledge biased towards locations with relative ease of access and to particular taxa. The second hurdle to surmount is the taxonomic confusion existing in the historic literature and the current ongoing debates within particular taxa. Several invertebrate taxa present in the Arctic may belong to species groups with an intricate taxonomy and which are challenging to identify. There are multiple instances of misidentifications and synonyms in the literature. Of the 88 Tardigrade taxa currently recognised in the literature from Svalbard many originate from older reports and identifications have not been verified based on modern taxonomy (Kaczmarek et al., 2012b). Another example is given by the 87 species of oribatid mite reported from Svalbard, many of which have not recently been observed and where synonyms and misidentifications may be suspected. This situation exists with most, if not all, the taxa discussed in this article. To complicate the situation further, material from earlier sampling may no longer exist, either being lost or, as in the case of much of Thor's material (including type specimens), deliberately destroyed (Winston, 1999). Hence, re-examination using modern taxonomic principles is no longer possible and a new inventory based on fresh material lodged in appropriate museums and collections is urgently required. Furthermore, forthcoming studies should employ molecular methods such as DNA-barcoding, which have yielded promising results in recent studies of Chironomidae (Stur and Ekrem, 2011). Molecular data may prove to be valuable in the identification of dispersal routes and time-scales for the invertebrate fauna of the Barents Sea archipelagoes. Based on morphological studies, efforts should also be made in preparing good and well-illustrated identification keys accessible to non-specialists so as to increase the taxonomic value of

upcoming ecological studies and enable future monitoring programs in the Arctic.

For both the terrestrial and freshwater systems there is clearly a need to assess biodiversity in areas away from the main settlements, and in specific habitats such as warm springs, naturally nutrient rich locations and more extreme habitats. Better understanding of food webs, life history strategies and the interactions between freshwater, terrestrial and marine ecosystems in different regions of the Arctic is also required. Work is underway to develop a monitoring network for freshwater biodiversity in the Arctic under the auspices of the Arctic Council (Culp et al., 2011) and the same is required in the terrestrial environment.

Current knowledge indicates that there are relatively few species endemic either to individual archipelagoes or to the region as a whole. This most likely reflects either the young age of the communities or relatively high linkage to mainland populations, both issues that may be resolved by the application of molecular methodologies. Observed endemism levels may also be more apparent than real, and reflect the limited sampling effort in other Arctic regions. Aspects of the dissimilarity of the invertebrate faunas of the different archipelagoes are striking. In particular, it might have been expected that Novaya Zemlya and Svalbard would show greater similarity or overlap in diversity than this study has found (Table 2). Clarification of the relative importance of eastern and western sources of colonizing diversity over time and in relation with regional glacial processes for both archipelagoes is clearly required.

This extensive synthesis of Barents Sea archipelago invertebrate biodiversity provides both a benchmark for the region and the foundation for future research in several key areas. In summary, we highlight the need for:

- explicit phylogeographical studies across the entire region (and more widely in the High Arctic),
- resolution of taxonomic confusion and the development of combined molecular and morphological approaches,
- strengthening of the linkages across biological and physical disciplines (e.g. glaciology, geomorphology, geology) in order to more clearly identify potentially ice-free areas,
- integration with oceanography and climatology in the context of understanding the role currents play in the occurrence and frequency of transfer events,
- linkage with regional climate change studies, to provide baselines for the documentation of, and studies of, colonizing species (including those associated with anthropogenic influence) and their impacts,
- integration of biodiversity studies across groups to give better description of ecosystem structure and function, especially in the context of large-scale carbon and nitrogen cycles, linkages between terrestrial and marine environments, and linkages between terrestrial and freshwater environments at catchment scale.

Acknowledgements

This paper is dedicated to the memory of Torstein Solhøy who passed away at a late stage in its preparation.

This work results from the Research Council of Norway ES446370 support to S.J. Coulson for the Invertebrate Fauna of Svalbard workshop, March 2011. We are grateful to Oleksandr Holovachov for help with obtaining Russian Nematoda references and Malin Daase (Norwegian Polar Institute) and Anna Sjöblom (UNIS) for assistance with the preparation of the figures. We also express our thanks to the three anonymous reviewers who provided valuable contributions to the manuscript. Ł. Kaczmarek was

partially funded by National Science Centre (Poland) and grant number N N304 014939.

Contribution of specific expertise: *Rotifera* De Smet, W.H.: *Nematoda* Boström, S., Sohlenius, B.: *Helminths* Carlsson, A., Kuklin, V.: *Gastrotricha* Kolicka M.: *Enchytraeidae* Maraldo, K.: *Tardigrada* Kaczmarek, Ł., Zawierucha, K.: *Acari* Gwiadzowicz, D.J., Lebedeva, N., Makarova, O., Melekhina, E., Solhøy, T.: *Araneae* Aakra, K., Tanasevitch, A.: *Collembola* Babenko, A., Fjellberg, A.: *Hemiptera* Simon, J.C.: *Phthiraptera* Gustafsson, D.: *Coleoptera* Ødegaard, F.: *Diptera* Ekrem, T., Søli, G., Stur, E.: *Hymenoptera* Hansen, L.O.: *Lepidoptera* Aarvik, L.: *Siphonaptera* Pilskog, H.E.: *Still waters* Christoffersen, K.S.: *Running waters* Brittain, J.E., Füreder, L.: *Adaptation to conditions* Simon J.C., Holmstrup M., *Paleoclimates* Velle, G., *Biogeography* Ávila-Jiménez, M.L.: *Environmental change* Convey, P.: *overall input of ideas and ms writing All authors.*

Appendix A. Supplementary data

Supplementary data related to this article can be found at <http://dx.doi.org/10.1016/j.soilbio.2013.10.006>.

References

- Aagaard, K., Bergvik, T., Sendstad, E., 1975. *Pyla (Salebria) fusca* Haw. (Lep., Pyralidae) found at Spitsbergen. Norw. J. Entomol. 22, 163–164.
- Aakra, K., Haug, E., 2003. Checklist of Norwegian spiders (Arachnida: Araneae), including Svalbard and Jan Mayen. Norw. J. Entomol. 50, 109–129.
- Abbott, R.J., Brochmann, C., 2003. History and evolution of the Arctic flora: in the footsteps of Eric Hultén. Mol. Ecol. 12, 299–313.
- ACIA, 2005. Impacts of a Warming Arctic: Arctic Climate Impact Assessment. Cambridge University Press, Cambridge.
- Adams, B.J., Bardgett, R.D., Ayres, E., Wall, D.H., Aislabie, J., Bamforth, S., Bargagli, R., Cary, C., Cavacini, P., Connell, L., Convey, P., Fell, J.W., Frati, F., Hogg, I.D., Newsham, K.K., O'Donnell, A.G., Russell, N., Seppelt, R.D., Stevens, M.I., 2006. Diversity and distribution of Victoria Land biota. Soil. Biol. Biochem. 38, 3003–3018.
- Albon, S.D., Stien, A., Irvine, R.J., Langvatn, R., Ropstad, E., Halvorsen, O., 2002. The role of parasites in the dynamics of a reindeer population. Proc. Royal Soc. Lond. Ser. B-biol. Sci. 269, 1625–1632.
- Aleksandrova, V.D., 1977. The Arctic and Antarctic: Their Division into Geobotanical Areas (D. Love, Trans, English Translation 1980). Cambridge University Press.
- Aleksandrova, V.D., 1983. Vegetation of the Soviet Polar Deserts (D. Love, Trans, English Translation 1988). Cambridge University Press.
- Alendal, E., Andersen, T., Fjeldså, Å., 1980. New records of *Apamea maillardi* (Geyer, 1832) (Lep., Noctuidae) in Adventdalen, Spitsbergen. Fauna. Norv. Ser. B 27, 78.
- Alexander, C.P., 1922. The Crane-flies (Superfamily Tipuloidea, order Diptera). In: Holtendahl, O. (Ed.), Norw. Nov. Zemlya Exp. 1921. No. 5. Det norske viden-skaps-akademii, Kristiania, pp. 1–26.
- Allegrucci, G., Carchini, G., Todisco, V., Convey, P., Sbordoni, V., 2006. A molecular phylogeny of Antarctic Chironomidae and its implications for biogeographical history. Polar Biol. 29, 320–326.
- Alsos, I.G., Eidesen, P.B., Ehrlich, D., Skrede, I., Westergaard, K., Jacobsen, G.H., Landvik, J.Y., Taberlet, P., Brochmann, C., 2007. Frequent long-distance plant colonization in the changing arctic. Science 316, 1601–1608.
- Alsos, I.G., Arnesen, G., Sandbakk, B.E., Elven, R., 2013. Svalbard Flora (online). Available from: <http://www.svalbardflora.net> (accessed January 2013).
- AMAP, 2011. Snow, Water, Ice and Permafrost in the Arctic (SWIPA): Climate Change and the Cryosphere. Arctic Monitoring and Assessment Programme (AMAP), Oslo.
- Amréen, H., 1964a. Ecological studies of zooplankton populations in some ponds on Spitsbergen. Zool. Bidr. fran Upps. 36, 161–191.
- Amréen, H., 1964b. Temporal variation of the rotifer *Keratella quadrata* (Müll.) in some ponds on Spitsbergen. Zool. Bidr. franUpps. 36, 193–208.
- Amréen, H., 1964c. Ecological and taxonomical studies on zooplankton from Spitsbergen. Zool. Bidr. fran Upps. 36, 209–276.
- Andersson, T., Forman, S.L., Ingólfsson, Ó., Manley, W.F., 2000. Stratigraphic and Morphologic constraints on the Weichselian glacial history of northern Prins Karls Forland, western Svalbard. Geogr. Ann.: Ser. A. Phys. Geogr. 82, 455–470.
- Andrássy, I., 1998. Nematodes in the sixth continent. J. Nemat. Morphol. System. 1, 107–186.
- Asbak, K., Aars, J., Derocher, A.E., Wiig, O., Oksanen, A., Born, E.W., Dietz, R., Sonne, C., Godfrid, J., Kapel, C.M.O., 2010. Serosurvey for *Trichinella* in polar bears (*Ursus maritimus*) from Svalbard and the Barents Sea. Vet. Parasitol. 172, 256–263.
- Aurivillius, C.W.S., 1883a. Eine Anguillulide aus der Schneeflora Spitzbergens. Bihang till K. Svenska vet. akad. handlingar 8, 3–15.
- Aurivillius, C., 1883b. Insecta a viris Doctissimis Nordenskiöld illum ducem sequentibus in insulis Waigatsch et Novaja Semlia anno 1875 collecta. Lepidoptera. Entomol. Tidskrift 4, 191–194.
- Ávila-Jiménez, M.L., 2011. High Arctic Invertebrate Biogeography. Patterns and Colonization Processes since the Last Glacial Maximum. Ph.D. thesis. University of Bergen, Bergen, Norway.
- Ávila-Jiménez, M.L., Coulson, S.J., 2011a. A Holarctic biogeographical analysis of the Collembola (Arthropoda, Hexapoda) unravels recent post-glacial colonization patterns. Insects 2, 273–296.
- Ávila-Jiménez, M.L., Coulson, S.J., 2011b. Can snow depth predict the distribution of the high Arctic aphid *Acyrtosiphon svalbardicum* (Hemiptera: Aphididae) on Spitsbergen? BMC Ecol. 11, 25.
- Ávila-Jiménez, M.L., Coulson, S.J., Solhøy, T., Sjöblom, A., 2010. Overwintering of Arctic arthropods: the case of the invertebrate fauna of Svalbard now and in the future. Polar Res. 29, 127–137.
- Ávila-Jiménez, M.L., Gwiadzowicz, D.J., Coulson, S.J., 2011. On the gamasid (Acar: Parasitiformes) mite fauna of Svalbard: a revised checklist of a high Arctic archipelago. Zootaxa 3091, 33–41.
- Babenko, A., Bulavintsev, V., 1993. Fauna and populations of Collembola on the Novaya Zemlya Archipelago. Rus. Entomol. J. 2, 3–19.
- Babenko, A., Fjellberg, A., 2006. Collembola Septentrionale. A Catalogue of Springtails of the Arctic Regions. KMK Scientific Press Ltd., Moscow.
- Baer, J.G., 1962. Cestoda. Zool. Iceland 2, 1–63.
- Balsamo, M., d'Hondt, J.-L., Kisielewski, J., Pierboni, L., 2008. Global diversity of gastrotrichs (Gastrotricha) in fresh waters. Hydrobiologi 595, 85–91.
- Baraniak, E., 2007. Taxonomic revision of the genus *Plutella* Schrank, 1802 (Lepidoptera: Plutellidae) from the Palaearctic region with notes on its phylogeny. Polskie Pismo Entomol. 76 (Suppl.), 1–122.
- Bardgett, R., 2005. The Biology of Soil. Oxford University Press, Oxford.
- Bayartgotokh, B., Schatz, H., Ekrem, T., 2011. Distribution of the soil mites of Svalbard with redescriptions of three known species (Acar: Oribatida). Int. J. Acarol. 37, 467–484.
- Bell, E., 2012. Life at Extremes. CABI Publishing, Wallingford.
- Bengtsson, S.-A., Fjellberg, A., Solhøy, T., 1974. Abundance of tundra arthropods in Spitsbergen. Entomol. Scand. 5, 137–142.
- Bengtsson, S.A., Fjellberg, A., Solhøy, T., 1975. *Amara quenselii* new record (Coleoptera: Carabidae) new to Svalbard, Norway. Norw J Zool 22, 81–82.
- Bengtsson, B.Å., Johansson, R., 2011. Nationalnyckeln till Sveriges Flora och Fauna. In: Fjärilar: Bronsmalar-rullvingemalar. Lepidoptera: Roeslerstammiidae-Lyonetiidae. ArtDatabanken, SLU, Uppsala.
- Bennike, O., Hedenas, L., 1995. Early Holocene land floras and faunas from Edgeøya, eastern Svalbard. Polar Res. 14, 205–214.
- Bernardová, A., Košnar, J., 2012. What do Holocene sediments in Petuniabukta, Spitsbergen reveal? Pol. Polar Res. 33 (4), 329–345.
- Bertolani, R., Rebecchi, L., 1993. A revision of the *Macrobiotus hufelandi* group (Tardigrada, Macrobiotidae), with some observations on the taxonomic characters of eutardigrades. Zool. Scr. 22, 127–152.
- Bertolani, R., Rebecchi, L., Giovannini, I., Cesari, M., 2011. DNA barcoding and integrative taxonomy of *Macrobiotus hufelandi* C.A.S. Schulze 1834, the first tardigrade species to be described, and some related species. Zootaxa 2997, 19–36.
- Bertram, G.C.L., Lack, D., 1938. Notes on the animal ecology of Bear Island. J. Animal Ecol. 7, 27–52.
- Betts-Piper, A.M., Zeeb, B.A., Smol, J.P., 2004. Distribution and autecology of chrysophyte cyst from high Arctic Svalbard lakes: preliminary evidence of recent environmental change. J. Paleolimnol. 31, 467–481.
- Beyens, L., Chardez, D., 1987. Evidence from testate amoebae for changes in some local hydrological conditions between c. 5000 BP and c. 3800 BP on Edgeøya (Svalbard). Polar Res. 5, 165–169.
- Beyens, L., Chardez, D., 1995. An annotated list of testate amoebae observed in the Arctic between the longitudes 27° E and 168° W. Arch. Protistenkd. 146, 219–233.
- Binda, M., Pilato, G., Dastych, H., 1980. Descrizione di una nuova specie di Eutardigrado, *Doryphoribus macrodon*. Animalia 7, 23–27.
- Birkemoe, T., Dozsa-Farkas, K., 1994. New records of Enchytraeidae (Oligochaeta) from Spitsbergen, Svalbard. Fauna Norv. Ser. A 15, 35–44.
- Birkemoe, T., Leinaas, H.P., 1999. Reproductive biology of the Arctic collembolan *Hypogastrura tulbergi*. Ecography 22, 31–39.
- Birkemoe, T., Leinaas, H.P., 2001. Growth and development in a high Arctic Collembola: adaptive variation in local populations living in contrasting thermal environments. Ecol. Entomol. 26, 100–105.
- Birkemoe, T., Sømme, L., 1998. Population dynamics of two collembolan species in an Arctic tundra. Pedobiologia 42, 131–145.
- Birkemoe, T., Coulson, S.J., Sømme, L., 2000. Life cycles and population dynamics of enchytraeids (Oligochaeta) from the High Arctic. Can. J. Zool. 78, 2079–2086.
- Birks, H.H., 1991. Holocene vegetational history and climatic change in west Spitsbergen: plant macrofossils from Skardtjørna, an Arctic lake. Holocene 1, 209–218.
- Biserov, V.I., 1996. Tardigrada of the Novaya Zemlya archipelago, collected by the marine arctic complex expedition in 1994. Arthropoda Selecta 5, 151–157.
- Biserov, V.I., 1999. A review of the Tardigrada from Novaya Zemlya with description of three new species and an evaluation of the state of the environment in the region. Zool. Anz. 238, 169–182.
- Block, W.C., 1966. Some arctic Oribatei. Acarologia 8, 161–162.

- Block, W., Convey, P., 1995. The biology, life cycle and ecophysiology of the Antarctic mite *Alaskozetes antarcticus* (Michael). *J. Zool.* 236, 431–449.
- Bogen, J., Bonsnes, T.E., 2003. Erosion and sediment transport in High Arctic rivers. *Svalbard. Polar Res.* 22, 175–189.
- Boheman, C.H., 1865. Bidrag til kännedom om Spetsbergens insektafauna. *Ov. K. Svenska Vet.-Akad. Forh.* B 22, 563–577.
- Boström, S., 1987. A new terrestrial nematode species (Rhabditida: Cephalobidae) from Spitzbergen. *Polar Biol.* 7, 375–378.
- Boström, S., 1988. A scanning electron microscope study of some species of terrestrial nematodes from Spitzbergen. *Nematologica* 33, 366–374.
- Boström, S., 1989. The taxonomic position of some teratoccephalid nematodes – a scanning electron microscope study. *Rev. Nématol.* 12, 181–190.
- Brinck-Lindroth, G., Smit, F.G.A.M., 2007. The Fleas (Siphonaptera) of Fennoscandia and Denmark. Brill Academic Pub, Leiden-Boston.
- Brittain, J.E., Milner, A.M., 2001. Ecology of glacier-fed rivers: current status and concepts. *Freshw. Biol.* 46, 1571–1578.
- Brittain, J.E., Gislason, G.M., Ponomorev, V.I., Bogen, J., Brørs, S., Jensen, A.J., Khoklova, L.G., Kochanov, S.K., Kokovkin, A.V., Melvold, K., Olafsson, J.S., Pettersson, L.-E., Stenina, A.S., 2009. Arctic rivers. In: Tockner, K., Uehlinger, U., Robinson, C.T. (Eds.), *Rivers of Europe*. Academic Press, London, pp. 337–379.
- Brochmann, C., Gabrielsen, T.M., Nordal, I., Landvik, J.Y., Elven, R., 2003. Glacial survival or tabula rasa? The history of the North Atlantic biota revisited. *Taxon* 52, 417–450.
- Brooks, S.J., Birks, H.J.B., 2004. The dynamics of Chironomidae (Insecta: Diptera) assemblages in response to environmental change during the past 700 years on Svalbard. *J. Paleolimnol.* 31, 483–498.
- Brown, J.M., 1936. Collembola from west Spitsbergen. *Entomol. Month. Mag.* 72, 62–65.
- Brundin, L., 1940. Studien über die Atheta Untergattung *Oreostiba* Ganglb. (Col. Staphylinidae). *Entomol. Tidskrift* 61, 56–130.
- Brussaard, L., Pulleman, M.M., Ouédraogo, E., Mando, A., Six, J., 2007. Soil fauna and soil function in the fabric of the food web. *Pedobiologia* 50, 447–462.
- Bryce, D., 1897. Contributions to the non-marine fauna of Spitsbergen.— Part II. Report on the Rotifera. *Proc. Zool. Soc. Lond.* 52, 793–799.
- Bryce, D., 1922. On some Rotifera from Spitsbergen. The Oxford University expedition to Spitsbergen, 1921. *J. Quekett Microsc. Club Ser.* 2 (14), 305–332.
- Bulavintsev, V.I., 1999. The Eurasian high arctic. *Priroda* 7, 54–65 (in Russian).
- Bulavintsev, V.I., Babenko, A.B., 1983. Soil-dwelling invertebrates in the eastern sector of Franz Josef land. *Zool. Z.* 62, 1114–1116 (in Russian).
- Bye, K., 1985. Cestodes of reindeer (*Rangifer tarandus platyrhynchus* Vrolik) on the Arctic islands of Svalbard. *Can. J. Zool.* 63, 2885–2887.
- Byzova, Yu.B., Uvarov, A.V., Petrova, A.D., 1995. Seasonal changes in communities of soil invertebrates in tundra ecosystems of Hornsund, Spitsbergen. *Polish Polar Res.* 16, 245–266.
- Bøyum, A., Kjemsø, J., 1978. Physiography of lake Linnévatn western Spitsbergen. *Verh. Internat. Verein. Limnol.* 20, 609–614.
- Callaghan, T.V., Bjørn, L.O., Chernov, Y., Chapin, T., Christensen, T.R., Huntley, B., Ims, R.A., Jonansson, M., Jolly, D., Janasson, S., Matveyeva, N., Panikov, N., Oechel, W., Shaver, G., Elster, J., Henttonen, H., Laine, K., Taulavuori, K., Taulavuori, E., Zöckler, C., 2004a. Climate change and UV-B impacts on Arctic tundra and polar desert ecosystems: biodiversity, distributions and adaptations of Arctic species in the context of environmental change. *Ambio* 33, 404–417.
- Callaghan, T.V., Bjørn, L.O., Chernov, Y., Chapin, T., Christensen, T.R., Huntley, B., Ims, R.A., Johansson, M., Jolly, D., Jonasson, S., Matveyeva, N., Panikov, N., Oechel, W., Shaver, G., 2004b. Climate change and UV-B impacts on Arctic tundra and polar desert ecosystems: effects on the structure of Arctic ecosystems in the short- and long-term perspectives. *Ambio* 33, 436–447.
- Carlsson, A.M., Wilson, K., Irvine, R.J., Piercy, S.B., Halvorsen, O., Coulson, S.J., Stien, A., Albon, S.D., 2012. Disease transmission in an extreme environment: nematode parasites infect reindeer during the Arctic winter. *Int. J. Parasitol.* 42, 89–795.
- Carlsson, A.M., Irvine, R.J., Wilson, K., Coulson, S.J., 2013. Adaptations to the Arctic: low-temperature development and cold tolerance in the free-living stages of a parasitic nematode from Svalbard. *Polar Biol.* 36, 997–1005.
- Carpenter, G.H., 1900. Collembola from Franz-Josef Land. *The Scientific Proc. Roy. Dublin Soc.* 9, 271–278.
- Carpenter, G.H., 1927. Further record of Collembola from Spitzbergen. *Proc. Roy. Irish Acad.* 37, 193–200.
- Carpenter, G.H., Phillips, J.K.C., 1922. The Collembola of Spitsbergen and Bear island. *Proc. Roy. Irish Acad.* 36, 11–21.
- Castella, E., Adalsteinsson, H., Brittain, J.E., Gislason, G.M., Lehmann, A., Lencioni, V., Lods-Crozet, B., Maiolini, B., Milner, A.M., Olafsson, J.S., Saltveit, S.J., Snook, D.L., 2001. Macrobenthic invertebrate richness and composition along a latitudinal gradient of European glacier-fed streams. *Freshwater Biol.* 46, 1811–1831.
- Chapin III, F.S., Berman, M., Callaghan, T.V., Convey, P., Crepin, A.-S., Danell, K., Ducklow, H., Forbes, B., Kofinas, G., McGuire, A.D., Nuttall, M., Virginia, R., Young, O., Zimov, S., Christensen, T., Godduhn, A., Murphy, E.J., Wall, D., Zockler, C., 2005. Polar systems. Chapter 25. In: Hassan, R., Scholes, R., Ash, N. (Eds.), *Millennium Ecosystem Assessment: Conditions and Trends*. Island Press, Washington, pp. 717–744.
- Chaubet, B., Derocles, S., Hullé, M., Le Ralec, A., Outreman, Y., Simon, J.C., Tomanović, Z., 2013. Two new species of aphid parasitoids (Hymenoptera, Braconidae, Aphidiinae) from the high arctic (Spitsbergen, Svalbard). *Zool. Anz.* 252, 34–40.
- Chernov, Y.I., Makarova, O.I., 2008. Beetles (Coleoptera) in high arctic. In: Penev, L., Erwin, T., Assmann, T. (Eds.), *Back to the Roots and Back to the Future? Towards a New Synthesis between Taxonomic, Ecological and Biogeographical Approaches in Carabidology*. Proceedings of the XIII European Carabidologists Meeting, Blagoevgrad, August 20–24, 2007. Pensoft Publishers, Moscow, pp. 213–246.
- Chown, S.L., Huiskes, A.H.L., Gremmen, N.J.M., Lee, J.E., Terauds, A., Crosbie, K., Frenot, Y., 2012a. Continent-wide risk assessment for the establishment of nonindigenous species in Antarctica. *Proc Natl Acad Sci U S A* 109, 4938–4943.
- Chown, S.L., Lee, J.E., Hughes, K.A., Barnes, J., Barrett, P.J., Bergstrom, D.M., Convey, P., Cowan, D.A., Crosbie, K., Dyer, G., Frenot, Y., Grant, S.M., Herr, D., Kennicutt, M.C., Lamers, M., Murray, A., Possingham, H.P., Reid, K., Riddle, M., Ryan, P.G., Sanson, L., Shaw, J.D., Sparrow, M.D., Summerhayes, C., Terauds, A., Wall, D.H., 2012b. Challenges to the future conservation of the Antarctic. *Science* 337, 158–159.
- Christensen, B., Doza-Farkas, K., 2006. Invasion of terrestrial enchytraeids into two postglacial tundras: north-eastern Greenland and the Arctic Archipelago of Canada (Enchytraeidae, Oligochaeta). *Polar Biol.* 29, 454–466.
- Christensen, J.H., Hewitson, B., Busuioc, A., Chen, A., Gao, X., Held, I., Jones, R., Kolli, R.K., Kwon, W.-T., Laprise, R., Magaña, V., Mearns, L., Menédez, C.G., Räisänen, J., Rinke, A., Sarr, A., Whetton, P., 2007. Regional climate projections. In: Solomon, S., Qin, D., Manning, M., Chen, Z., Marquis, M., Averyt, K.B., Tignor, M., Miller, H.L. (Eds.), *Climate Change 2007: the Physical Science Basis, Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge and New York, pp. 847–940.
- Christoffersen, K.S., Jeppesen, L., Moorhead, D.L., Tranvik, L., 2008. Food web relationships and community structures in high-latitude lakes. In: Vincent, W., Laybourn-Parry, J. (Eds.), *High Latitude Lake and River Ecosystems*. Oxford University Press, Oxford, pp. 269–291.
- Claxton, S.K., 1998. A revision of the genus *Minibiotus* (Tardigrada: Macrobiotidae) with descriptions of eleven new species from Australia. *Rec. Aust. Mus.* 50, 125–160.
- Clay, T., 1976. Geographical distribution of the avian lice (Phthiraptera): a review. *J. Bombay Natural Hist. Soc.* 71, 536–547.
- Convey, P., 2005. Recent lepidopteran records from sub-Antarctic South Georgia. *Polar Biol.* 28, 108–110.
- Convey, P., 2007. Influences on and origins of terrestrial biodiversity of the sub-Antarctic islands. *Papers Proc. Royal Soc. Tasmania* 141, 83–93.
- Convey, P., 2011. Antarctic terrestrial biodiversity in a changing world. *Polar Biol.* 34, 1629–1641.
- Convey, P., 2013. Antarctic ecosystems. In: Levin, S.A. (Ed.), *Encyclopedia of Biodiversity*, Second ed., vol. 1 San Diego, pp. 179–188.
- Convey, P., Smith, R.I.L., 1997. The terrestrial arthropod fauna and its habitats in northern Marguerite Bay and Alexander Island, maritime Antarctic. *Antarct. Sci.* 9, 12–26.
- Convey, P., Stevens, M.I., 2007. Antarctic biodiversity. *Science* 317, 1877–1878.
- Convey, P., Worland, M.R., 2000. Survival of freezing by free-living Antarctic soil nematodes. *Cryo-letters* 21, 327–332.
- Convey, P., Wynn-Williams, D.D., 2002. Antarctic soil nematode response to artificial climate amelioration. *Eur. J. Soil Biol.* 37, 255–259.
- Convey, P., Block, W., Peat, H.J., 2003. Soil arthropods as indicators of water stress in Antarctic terrestrial habitats? *Glob. Change Biol.* 9, 1718–1730.
- Convey, P., Gibson, J., Hillenbrand, C.D., Hodgson, D.A., Pugh, P.J.A., Smellie, J.L., Stevens, M.I., 2008. Antarctic terrestrial life – challenging the history of the frozen continent? *Biol. Rev.* 83, 103–117.
- Convey, P., Stevens, M.I., Hodgson, D.A., Smellie, J.L., Hillenbrand, C.-D., Barnes, D.K.A., Clarke, A., Pugh, P.J.A., Linse, K., Cary, S.C., 2009. Exploring biological constraints on the glacial history of Antarctica. *Quat. Sci. Rev.* 28, 3035–3048.
- Convey, P., Aitken, S., di Prisco, G., Gill, M.J., Coulson, S.J., Barry, T., Jónsdóttir, I.J., Dang, P.T., Hik, D., Kulkarni, T., Lewis, G., 2012. The impacts of climate change on circumpolar biodiversity. *Biodiversity* 13, 1–10.
- Cooper, E.J., 2011. Polar desert vegetation and plant recruitment in Murchisonfjord, Nordaustlandet, Svalbard. *Geogr. Ann. Ser. A* 93, 243–252.
- Cooper, E.J., Dullinger, S., Semenchuk, P., 2011. Late snow melt delays plant development and results in lower reproductive success in the High Arctic. *Plant Sci.* 180, 157–167.
- Coulson, S.J., 2007a. The terrestrial and freshwater invertebrate fauna of the High Arctic archipelago of Svalbard. *Zootaxa* 1448, 41–58.
- Coulson, S.J., 2007b. On the occurrence of *Oryzaephilus mercator* (Fauvel, 1889) (Coleoptera: Silvanidae) in Svalbard, Norway. *Norw. J. Entomol.* 54, 21–22.
- Coulson, S.J., 2009. Association of the soil mite *Diapterobates notatus* (Acari, Oribatidae) with *Cynomya mortuorum* (Linnaeus, 1761) (Caliphoridae, Calliphorinae): implications for the dispersal of oribatid mites. *Int. J. Acarol.* 35, 175–177.
- Coulson, S.J., 2013a. The terrestrial invertebrate fauna of the Svalbard archipelago in a changing world: history of research and challenges. *Can. Entomol.* 145, 131–146.
- Coulson, S.J., 2013b. Checklist of the Terrestrial and Freshwater Invertebrate Fauna of Svalbard [online]. Available from: http://www.unis.no/35_STAFF/staff_webpages/biology/steve_coulson/default.htm [Accessed January 2013].
- Coulson, S.J., Birkemoe, T., 2000. Long term cold tolerance in Arctic invertebrates: recovery after four years at below -20°C. *Can. J. Zool.* 78, 2055–2058.
- Coulson, S.J., Refseth, D., 2004. The terrestrial and freshwater invertebrate fauna of Svalbard (and Jan Mayen). In: Prestrud, P., Strøm, H., Goldman, H.V. (Eds.),

- A Catalogue of the Terrestrial and Marine Animals of Svalbard, Skrifter, 201. Norwegian Polar Institute, Tromsø, pp. 57–122.
- Coulson, S.J., Hodkinson, I.D., Strathdee, A.T., Bale, J.S., Block, W., Worland, M.R., Webb, N.R., 1993. Simulated climate change: the interaction between vegetation type and microhabitat temperatures at Ny Ålesund, Svalbard. *Polar Biol.* 13, 67–70.
- Coulson, S.J., Hodkinson, I.D., Strathdee, A.T., Block, W., Webb, N.R., Bale, J.S., Worland, M.R., 1995. Thermal environments of Arctic soil organisms during winter. *Arct. Alp. Res.* 27, 365–371.
- Coulson, S.J., Leinaas, H.P., Ims, R.A., Søvik, G., 2000. Experimental manipulation of the winter surface ice layer: the effect on a High Arctic soil microarthropod community. *Ecography* 23, 299–306.
- Coulson, S.J., Hodkinson, I.D., Webb, N.R., Harrison, J.A., 2002a. Survival of terrestrial soil-dwelling arthropods on land and in seawater: implications for trans-oceanic dispersal. *Funct. Ecol.* 16, 353–356.
- Coulson, S.J., Hodkinson, I.D., Webb, N.R., Mikkola, K., Harrison, J.A., Pedgley, D.E., 2002b. Aerial colonization of high Arctic islands by invertebrates: the diamondback moth *Plutella xylostella* (Lepidoptera: Yponomeutidae) as a potential indicator species. *Divers. Distrib.* 8, 327–334.
- Coulson, S.J., Hodkinson, I.D., Webb, N.R., 2003a. Microscale distributional patterns in high Arctic soil microarthropods communities: the influence of plant species within the vegetation mosaic. *Ecography* 26, 801–809.
- Coulson, S.J., Hodkinson, I.D., Webb, N.R., Convey, P., 2003b. A high-Arctic population of *Pyla fusca* (Lepidoptera, Pyralidae) on Svalbard? *Polar Biol.* 26, 283–285.
- Coulson, S.J., Moe, B., Monson, F., Gabrielsen, G.W., 2009. The invertebrate fauna of High Arctic seabird nests: the microarthropod community inhabiting nests on Spitsbergen, Svalbard. *Polar Biol.* 32, 1041–1046.
- Coulson, S.J., Ávila-Jiménez, M.L., Fjellberg, A., Snazell, R., Gwiazdowicz, D.J., 2011. On the Collembola, Araneae and Gamasida from the Kinnvika region of Nordaustlandet, Svalbard. *Geogr. Ann.* 93, 253–257.
- Coulson, S.J., Fjellberg, A., Gwiazdowicz, D.J., Lebedeva, N.V., Melekhina, E.N., Solhøy, T., Erséus, C., Maraldo, K., Miko, L., Schatz, H., Schmelz, R.M., Søli, G., Stur, E., 2013a. Introduction of invertebrates into the High Arctic via imported soils: the case of Barentsburg in the Svalbard. *Biol. Invas.* 15, 1–5.
- Coulson, S.J., Fjellberg, A., Gwiazdowicz, D.J., Lebedeva, N.V., Melekhina, E.N., Solhøy, T., Erséus, C., Maraldo, K., Miko, L., Schatz, H., Schmelz, R.M., Søli, G., Stur, E., 2013b. The invertebrate fauna of the anthropogenic soils in the High Arctic settlement of Barentsburg: Svalbard. *Polar Res.* 32, 19273. <http://dx.doi.org/10.3402/polar.v32i0.19273>.
- Culp, J., Gantner, N., Gill, N., Reist, J., Wrona, F., 2011. Development of an Arctic Freshwater Biodiversity Monitoring Plan: Framework Document. In: CAFF Monitoring Series Report, Vol. 4. CAFF International Secretariat, Akureyri.
- Cyrich, D., Krumpál, M., 1991. Incidence and taxonomy of *Mioctenopsylla arctica arctica* Rothschild 1922 (Siphonaptera) in the Spitsbergen. *Biológia (Bratislava)* 46, 545–549.
- Dampf, A., 1911. Land-Arthropoden von der Bären-Insel und Spitzbergen gesammelt in den Jahren 1907 und 1908. Aphaniptera. In: Koenig, A. (Ed.), Avifauna Spitzbergensis, Forschungsreisen nach der Bären-Insel und dem Spitzbergen Archipel, mit ihren faunistischen und floristischen Ergebnissen, pp. 276–279. Printed privately, Bonn.
- Danks, H.V., 1971. Overwintering of some north temperate and Arctic Chironomidae. I. Winter environment. *Can. Entomol.* 103, 589–604.
- Danks, H.V., 1992. Long life cycles in insects. *Can. Entomol.* 124, 167–187.
- Dastych, H., 1983. Two new Eutardigrada species from west Spitsbergen and the Tatra Mts. *Bull. Soc. Amis Sci. Lett. Poznań* 23, 195–200.
- Dastych, H., 1985. West Spitsbergen Tardigrada. *Acta Zool. Cracvien.* 28, 169–214.
- Davidson, R.K., Romig, T., Jenkins, E., Tryland, M., Robertson, L.J., 2012. The impact of globalization on the distribution of *Echinococcus multilocularis*. *Trends Parasitol.* 28, 239–247.
- De Jonckheere, J.F., 2006. Isolation and molecular identification of free-living amoebae of the genus *Naegleria* from Arctic and sub-Antarctic regions. *Eur. J. Protistol.* 42, 115–123.
- De Smet, W.H., 1988. Rotifers from Bjørnøya (Svalbard), with the description of *Cephalodella evabroedi* n. sp. and *Synchaeta lakowitziana arctica* n. subsp. *Fauna Norv. Ser. A* 9, 1–18.
- De Smet, W.H., 1990. Notes on the monogonont rotifers from submerged mosses collected on Hopen (Svalbard). *Fauna Norv. Ser. A* 11, 1–8.
- De Smet, W.H., 1993. Report on rotifers from Barentsøya, Svalbard (78°30'N). *Fauna Norv. Ser. A* 14, 1–26.
- De Smet, W.H., 1995. Description of *Encentrum dieteri* sp. nov. (Rotifera, Dicranophoridae) from the High Arctic, with redescription of *E. bidentatum* (Lie-Petersen, 1906) and *E. murraiyae* Bryce, 1922. *Belgian J. Zool.* 125, 349–361.
- De Smet, W.H., Van Rompu, E.A., 1994. Rotifera and Tardigrada from some cryoconite holes on a Spitsbergen (Svalbard) glacier. *Belgian J. Zool.* 124, 27–37.
- De Smet, W.H., Van Rompu, E.A., Beyens, L., 1987. Rotifera, Gastrotricha en Tardigrada uit Shetland, de Faroëren en Spitsbergen (Rotifera, Gastrotricha et Tardigrada provenant du Shetland, les îles Féroé et du Spitsberg). *Natuurwet Tijdschr.* 69, 81–102.
- De Smet, W.H., Van Rompu, E.A., Beyens, L., 1988. Contribution to the rotifers and aquatic Tardigrada of Edgeøya (Svalbard). *Fauna Norv. Ser. A* 9, 19–30.
- Denlinger, D.L., Lee, R., 2010. Low Temperature Biology of Insects. Cambridge University Press, Cambridge.
- Desamps, S., 2013. Winter temperature affects the prevalence of ticks in an arctic seabird. *PLoS ONE* 8, e65374.
- Dollery, R., Hodkinson, I.D., Jónsdóttir, I.S., 2006. Impact of warming and timing of snow melt on soil microarthropod assemblages associated with *Dryas*-dominated plant communities on Svalbard. *Ecography* 29, 111–119.
- Dunton, K., 1992. Arctic biogeography: the paradox of the marine benthic fauna and flora. *Trends Ecol. Evol.* 7, 183–189.
- Eastop, V.F., Blackman, R.L., 2005. Some new synonyms in Aphididae (Hemiptera: Sternorrhyncha). *Zootaxa* 1089, 1–36.
- Edwards, F.W., 1935. Diptera from Bear Island. *Ann. Mag. Natural Hist.* 15, 531–543.
- Ehrenberg, G.C., 1874. Das unsichtbar wirkende Leben der Nordpolarzone am Lande und in den Meerestieflgründen bei 300 mal verstärkter Sehkraft, nach Materialien der Germania erläutert. In: Hartlaub, G., Lindeman, M. (Eds.), Die zweite deutsche Nordpolarfahrt in den Jahren 1869 und 1870 unter Führung des Kapitän Karl Koldewey, Leipzig, pp. 437–467.
- Eidesen, P.B., Ehrlich, D., Bakkestuen, V., Alsos, I.G., Gilg, O., Taberlet, P., Brochmann, C., 2013. Genetic roadmap of the Arctic: plant dispersal highways, traffic barriers and capitals of diversity. *New Phytol.* <http://dx.doi.org/10.1111/nph.12412>. Online First.
- Ellis-Evans, J.C., Galchenko, V., Laybourn-Parry, J., Mylnikov, A.P., Petz, W., 2001. Environmental characteristics and microbial plankton activity of freshwater environments at Konbgsfjorden, Spitsbergen (Svalbard). *Arch. Hydrobiol.* 152, 609–632.
- Elton, C.S., 1925a. The dispersal of insects to Spitsbergen. *Trans. Entomol. Soc. Lond.* 73, 289–299.
- Elton, C.S., 1925b. Coleoptera and Lepidoptera from Spitsbergen. Results of the Oxford University expedition to Spitsbergen, 1924. *Ann. Mag. Natural Hist.* 16, 357–359.
- Elton, C.S., 1934. Aphids and hover-flies in North-East Land (Spitsbergen) in 1924: an additional note. Greenland and Spitsbergen Papers. Scientific results of the Oxford University Expedition to Greenland in 1928 together with some further results of the Oxford Expeditions to Spitsbergen in 1921, 1923 and 1924. *Proc. Entomol. Soc. Lond.* 23, 76–77.
- Elvebakke, A., Hertel, H., 1996. Lichens. In: Elvebakke, A., Prestrud, P. (Eds.), A Catalogue of Svalbard Plants, Fungi, Algae and Cyanobacteria, Skrifter, vol. 198. Norwegian Polar Institute, Oslo, pp. 271–360.
- Elven, R., Elvebakke, A., 1996. Vascular plants. In: Elvebakke, A., Prestrud, P. (Eds.), A Catalogue of Svalbard Plants, Fungi, Algae and Cyanobacteria, Skrifter, vol. 198. Norwegian Polar Institute, Oslo, pp. 9–56.
- European Commission, 2012. Research and Innovation Infrastructures [online] Available from: http://ec.europa.eu/research/infrastructures/index_en.cfm?pg=esfri [accessed May 2012].
- Eveleigh, E.S., Threlfall, W., 1976. Population dynamics of lice (Mallophaga) on auks (Alcidae) from Newfoundland. *Can. J. Zool.* 54, 1694–1711.
- Evenset, A., Christensen, G.N., Kallenborn, R., 2005. Selected chlorobornanes, polychlorinated naphthalenes and brominated flame retardants in Bjørnøya freshwater biota. *Environ. Pollut.* 136, 419–430.
- Fadnes, K., 2010. Climate History of the Arctic: Temperature Reconstruction of an Arctic Lake on Spitsbergen Based on Chironomids. M.Sc.thesis. Department of Biology, University of Bergen, Bergen.
- Fauna Europaea, 2011. Fauna Europaea Version 2.4. Web Service available online at: <http://www.faunaeur.org> (Accessed at January 10 2012).
- Ferris, G.F., 1923. Mallophaga. In: Holtendahl, O. (Ed.), Norw. Nov. Zemlya Exp. 1921. No. 8. Det norske videnskaps-akademii, Kristiania, pp. 11–12.
- Fjellberg, A., 1975. *Vertagopus pseudocinereus* n.sp. A new species of Collembola (Isotomidae) from North Norway. *Entomol. Scand.* 6, 212–214.
- Fjellberg, A., 1983. Three species of staphylinid Coleoptera new to Spitsbergen, Norway. *Fauna. Norv. Ser. B* 30, 110–111.
- Fjellberg, A., 1994. The Collembola of the Norwegian Arctic Islands. In: Meddelelser, vol. 133. Norwegian Polar Institute, Oslo.
- Fontaneto, D., Herniou, E.A., Boschetti, C., Caprioli, M., Melone, G., Ricci, C., Barracough, T.G., 2007. Independently evolving species in asexual bdelloid rotifers. *PLoS Biol.* 5, 1–8.
- Fontaneto, D., Kaya, M., Herniou, E.A., Barracough, T.G., 2009. Extreme levels of hidden diversity in microscopic animals (Rotifera) revealed by DNA taxonomy. *Mol. Phylogenet. Evol.* 53, 182–189.
- Fontoura, P., Pilato, G., 2007. *Diphascon (Diphascon) faiälense* sp. nov. a new species of Tardigrada (Eutardigrada, Hypsibidae) from the Azores and a key to the species of the *D. pingue* group. *Zootaxa* 1589, 47–55.
- Forman, S.L., Lubinski, D., Weihe, R.R., 2000. The Holocene occurrence of reindeer on Franz Josef land, Russia. *Holocene* 10, 763–768.
- Forsslund, K.H., 1957. Notizen über die Tierwelt des nordschwedischen Waldgebietes. *Statens Skogsförskörs. Meddel* 34, 1–283.
- Forsslund, K.H., 1964. *Liochthonius muscorum* n. sp. und *L. lapponicus* (Träg.). *Entomol. Tidskrift* 85, 236–239.
- Freckman, D.W., Virginia, R.A., 1997. Low-diversity Antarctic soil nematode communities: distribution and response to disturbance. *Ecology* 78, 363–369.
- Frenot, Y., Chown, S.L., Whinam, J., Selkirk, P.M., Convey, P., Skotnicki, M., Bergstrom, D.M., 2005. Biological invasions in the Antarctic: extent, impacts and implications. *Biol. Rev.* 80, 45–72.
- Friese, H., 1923. Apidae in Hymenoptera. In: Holtendahl, O. (Ed.), Norw. Nov. Zemlya Exp. 1921. No. 14. Det norske videnskaps-akademii, Kristiania, pp. 1–10.
- Frisvoll, A.A., Elvebakke, A., 1996. Bryophytes. In: Elvebakke, A., Prestrud, P. (Eds.), A Catalogue of Svalbard Plants, Fungi, Algae and Cyanobacteria, Skrifter, vol. 198. Norwegian Polar Institute, Oslo, pp. 57–172.

- Fuglei, E., Stien, A., Yoccoz, N.G., Ims, R.A., Eide, N.E., Prestrud, P., Oksanen, A., 2008. Spatial distribution of *Echinococcus multilocularis*, Svalbard, Norway. *Emerg. Infect. Dis.* 14, 73–75.
- Füreder, L., Brittain, J.E., 2006. High Arctic Stream Food Webs – what Remains at the Limits of Benthic Life?. Annual meeting North American Bentholological Society, Anchorage, Alaska.
- Førland, E.J., Benestad, R., Hanssen-Bauer, I., Haugen, J.E., Skaugen, T.E., 2011. Temperature and precipitation development at Svalbard 1900–2100. *Adv. Meteorol.* Article ID 893790.
- Gagarin, V.G., 1997a. *Aphelenchoides robustus* sp. n., a new freshwater nematode from archipelago Novaya Zemlya (Nematoda: Aphelenchoididae). *Zoosyst. Rossica* 5, 209–211.
- Gagarin, V.G., 1997b. New species of freshwater nematodes of the order Monhysterida from the Novaya Zemlya archipelago (Nematoda). *Zoosyst. Rossica* 6, 21–30.
- Gagarin, V.G., 1997c. Fauna svobodnozhivushchih vodoemov archipelaga Novaja Zemlya. *Biol. Vnudrennikh Vod* 1997, 33–40 (in Russian).
- Gagarin, V.G., 1999. Nekotorye dannye o svobodnozhivushchih nematodach vodoemov archipelaga Novaja Zemlya i ostrova Vajgatch. *Biol. Vnudrennikh Vod*, 32–41 (in Russian).
- Gagarin, V.G., 2000. New species of nematodes from waterbodies of Russian Arctica. *Biol. Vnudrennikh Vod*, 9–15 (in Russian).
- Gagarin, V.G., 2001. Review of free-living nematode fauna from fresh waters of the Russian Arctic and Subarctic. *Biol. Vnudrennikh Vod*, 32–37 (in Russian).
- Galaktionov, K.V., 1996. Life cycles and distribution of seabird helminths in Arctic and sub-Arctic regions. *Bull. Scand. Soc. Parasitol.* 6, 31–49.
- Galaktionov, K.V., Bustness, J.O., 1999. Distribution patterns of marine bird digenetic larvae in periwinkles along the southern Barents Sea coast. *Dis. Aquat. Org.* 37, 221–230.
- Galaktionov, K.V., Marasaev, S.F., 1992. Parasitological Investigations at Hooker Island. In: Meddelelser, vol. 120. Norwegian Polar Institute, Tromsø, pp. 51–54.
- Gataullin, V., Mangerud, J., Svendsen, J.I., 2001. The extent of the Late Weichselian ice sheet in the southeastern Barents Sea. *Glob. Planet Change* 31, 453–474.
- Ghilarov, A.M., 1967. The zooplankton of Arctic rock pools. *Oikos* 18, 82–95.
- Giebel, C.G.A., 1874. Insecta Epizoica. In: Die auf Saugethieren und Vogeln schmarotzenden Insecten nach Chr. L. Nitzsch's Nachlass bearbeitet. O. Wiegand, Leipzig.
- Gillespie, M., Hodgkinson, I.D., Cooper, E.J., Bird, J.M., Jónsdóttir, I.S., 2007. Life history and host-plant relationships of the rare endemic Arctic *Acyrtosiphon calvulus* in a changing environment. *Entomol. Exp. Appl.* 123, 229–237.
- Gíslason, G.M., 2005. Origin of freshwater fauna of the North-Atlantic islands: present distribution in relation to climate and possible migration routes. *Verh. Internat. Verein. Limnol.* 29, 198–203.
- Gokhman, V.V., 1988. The runoff regime of rivers on Spitsbergen. *Polar Geogr. Geol.* 12, 285–296.
- Gorbunow, G.P., 1929. Vorläufiger Bericht über die in den Jahren 1923, 1924, 1925 und 1927 ausgeführten Untersuchungen der süßen und brackigen Gewässer auf Nowaja Semlja, vol. 40. Transactions of the Institute for Exploration of the North, Moscow, pp. 147–154 (in Russian with German summary).
- Greenslade, P., 1995. Collembola from the Scotia arc and Antarctic Peninsula including descriptions of two new species and notes on biogeography. *Polskie Pismo Entomol.* 64, 305–319.
- Greenslade, P., 2006. The Invertebrates of Macquarie Island. Australian Antarctic Division, Hobart.
- Greenslade, P., Convey, P., 2012. Exotic Collembola on subantarctic islands: pathways, origins and biology. *Biol. Invas.* 14, 405–417.
- Greenslade, P., Potapov, M., Russel, D., Convey, P., 2012. Lessons from History – Global Collembola on Deception Island. *J. Insect. Sci.* 12, Article 111.
- Gressitt, J.L., 1970. Subantarctic entomology and biogeography. *Pac Insects Monogr.* 23, 295–374.
- Guilizzoni, P., Marchetto, A., Lami, A., Brauer, A., Vigliotti, L., Musazzi, S., Langone, L., Manca, M., Lucchini, F., Calanchi, N., Dinelli, E., Mordenti, A., 2006. Records of environmental and climatic changes during the late Holocene from Svalbard: palaeolimnology of Kongressvatnet. *J. Paleolimnol.* 36, 325–351.
- Gwiazdowicz, D.J., Coulson, S.J., 2010a. First record of *Thinoseius spinosus* (Acari, Euphidiidae) from the High Arctic Island of Spitsbergen (Svalbard) including a key to deutonymphs of genus *Thinoseius*. *Int. J. Acarol.* 36, 233–236.
- Gwiazdowicz, D.J., Coulson, S.J., 2010b. Phoresis of the deutonymphs of *Thinoseius spinosus* (Acari, Euphidiidae) on the calliphorid dipteran *Protophormia terraenovae* on the High Arctic island of Spitsbergen (Svalbard) including a key to deutonymphs of *Thinoseius*. *Internat. J. Acarol.* 36, 233–236.
- Gwiazdowicz, D.J., Coulson, S.J., 2011. High arctic gamasid mites (Acari, Mesostigmata): community composition on Spitsbergen, Svalbard. *Polar Res.* 30, 1–7.
- Gwiazdowicz, D.J., Gulvik, M.E., 2008. Mesostigmatid mites (Acari, Mesostigmata) in Svalbard. In: Proceedings of the 32th International Polar Symposium, Wrocław, pp. 32–34.
- Gwiazdowicz, D.J., Rakowski, R., 2009. Redescription of *Proctolaelaps parvanalis* (Thor, 1930) (Acari: Ascidae) from Spitsbergen. *Entomol. Fenn.* 20, 281–286.
- Gwiazdowicz, D.J., Coulson, S.J., Ávila-Jiménez, M.L., 2009. First records of *Zercon andrei* Sellnick, 1958 and *Zerconopsis muestairi* (Schweizer, 1949) (Acari, Mesostigmata) from Björnøya, Svalbard. *Norw. J. Entomol.* 56, 117–119.
- Gwiazdowicz, D.J., Teodorowicz, E., Coulson, S.J., 2011a. Redescription of *Zercon solenites* Haarlov, 1942 (Acari, Zerconidae) with a key to the Svalbard species of the genus *Zercon*. *Int. J. Acarol.* 37, 135–148.
- Gwiazdowicz, D.J., Teodorowicz, E., Coulson, S.J., 2011b. Redescription of *Arctoseius haarovi* Lindquist, 1963 (Acari: Ascidae) from Spitsbergen. *Entomol. Fenn.* 22, 140–148.
- Gwiazdowicz, D.J., Coulson, S.J., Grytnes, J.A., Pilskog, H.E., 2012a. The bird ectoparasite *Dermanyssus hirundinis* (Acari, Mesostigmata) in the High Arctic: a new parasitic mite to Spitsbergen, Svalbard. *Acta Parasitol.* 57, 378–384.
- Gwiazdowicz, D.J., Solhøy, T., Coulson, S.J., Lebedeva, N., Melekhina, E., 2012b. First record of *Vulgarogamasus immanis* (Acari, Mesostigmata) in Svalbard. *Polish Polar Res.* 33, 35–39.
- Hackman, W., Nyholm, E.S., 1968. Notes on the arthropod fauna of Spitsbergen II. 9. Mallophaga from Spitsbergen and Bear Island. *Ann. Entomol. Fenn.* 34, 75–82.
- Hagen, J.O., Lefauconnier, B., 1995. Reconstructed runoff from the high Arctic basin Bayelva based on mass-balance measurements. *Nord. Hydrol.* 26, 285–296.
- Hågvar, S., 1971. Some Observations on Coleoptera, Hymenoptera and Siphonaptera in Svalbard 1968. *Norsk Polarinstitutt Arbok* 1969. Norwegian Polar Institute, Oslo, pp. 101–106.
- Hågvar, S., 2010. Primary succession of springtails (Collembola) in a Norwegian glacier foreland. *Arctic Antarct. Alpine Res.* 42, 422–429.
- Hald, M., Dokken, T., Hagen, S., 1996. Palaeoceanography on the European Arctic margin during the last deglaciation. In: Andrews, J.T., Austin, W.E.N., Bergsten, H., Jennings, A.E. (Eds.), Late Quaternary Palaeoceanography of the North Atlantic Margins, vol. 111. Geological Society Special Publication, London, pp. 275–287.
- Halvorsen, O., 1986. Epidemiology of reindeer parasites. *Parasitol. Today* 2, 334–339.
- Halvorsen, O., Bye, K., 1999. Parasites, biodiversity, and population dynamics in an ecosystem in the High Arctic. *Vet. Parasitol.* 84, 205–227.
- Halvorsen, O., Gullestad, N., 1976. Freshwater crustacean in some areas of Svalbard. *Arch. Hydrobiol.* 78, 383–395.
- Hammar, J., 2000. Cannibals and parasites: conflicting regulators of bimodality in high latitude Arctic char, *Salvelinus alpinus*. *Oikos* 88, 33–47.
- Hammer, M., 1946. The Zoology of east Greenland. *Oribatids. Medd. om Grønland* 122, 1–39.
- Hammer, Ø., Jamtveit, B., Benning, L.G., Dysthe, D.K., 2005. Evolution of fluid chemistry during travertine formation in the Troll thermal springs, Svalbard, Norway. *Geofluids* 5, 140–150.
- Hanssen, O., Lid, J., 1932. Flowering plants of Franz Josef Land collected on the Norwegian scientific expedition 1930. *Skrifter om Svalbard og Ishavet* 39, 1–42.
- Hanssen-Bauer, I., Førland, E.J., 1998. Long-term trends in precipitation and temperature in the Norwegian Arctic: can they be explained by changes in atmospheric circulation patterns? *Clim. Res.* 10, 143–153.
- Harriman, V.B., Alisauskas, R.T., 2010. Of fleas and geese: the impact of an increasing nest ectoparasite on reproductive success. *J. Avian Biol.* 41, 573–579.
- Harrison, P.M., Rothery, P., Block, W., 1991. Drying processes in the Antarctic collembolan *Cryptopygus antarcticus* (Willem). *J. Insect Physiol.* 37, 883–890.
- Hauge, E., Somme, L., 1997. Records of spiders (Araneae) from Nordaustlandet and Sjuøyane, Svalbard. *Fauna Norv. Ser. A* 18, 17–20.
- Heikinheimo, O., 1968. Notes on the arthropod fauna of Spitsbergen. II: The aphid fauna of Spitsbergen. *Ann. Entomol. Fenn.* 34, 82–93.
- Henttonen, H., Fuglei, E., Gower, C.N., Haukisalmi, V., Ims, R.A., Niemimaa, J., Yoccoz, N.G., 2001. *Echinococcus multilocularis* on Svalbard: introduction of an intermediate host has enabled the local life-cycle. *Parasitology* 12, 547–552.
- Hertzberg, K., Yoccoz, N.G., Ims, R.A., Leinaas, H.P., 2000. The effects of spatial habitat configuration on recruitment, growth and population structure in Arctic Collembola. *Oecologia* 124, 382–390.
- Hessen, D.O., Blomqvist, P., Dahl-Hansen, G., Drakare, S., Lindstrom, E.S., 2004. Production and food web interactions of Arctic freshwater plankton and responses to increased DOC. *Arch. Hydrobiol.* 159, 289–307.
- Hille Ris Lambers, D., 1952. The aphid fauna of Greenland. *Medd. om Grønland* 136, 1–33.
- Hille Ris Lambers, D., 1960. Additions to the aphid fauna of Greenland. *Medd. om Grønland* 159, 1–18.
- Hirschmann, W., 1966. Gangsystematik der Parasitiformes. Teil 15. *Gänge von Litoral milben und neue Litoral milbenarten. Acarologie* 9, 25–44.
- Hirvenoja, M., 1967. Chironomidae and Culicidae (Dipt.) from Spitsbergen. *Ann. Entomol. Fenn.* 33, 52–61.
- Hisdal, V., 1985. Geography of Svalbard. Norwegian Polar Institute, Oslo, Norway.
- Hodgkinson, I.D., 2004. *Aptinothrips rufus* Haliday (Thysanoptera) and *Apamea zeta* (Treitschke) (Lepidoptera) on Svalbard. *Norw. J. Entomol.* 51, 127–128.
- Hodgkinson, I.D., 2013. Terrestrial and freshwater invertebrates. In: Meltofte, H. (Ed.), Arctic Biodiversity Assessment. Status and Trends in Arctic biodiversity. Conservation of Arctic Flora and Fauna, Akureyri, pp. 194–223.
- Hodgkinson, I.D., Bird, J.M., 2004. Anoxia tolerance in High Arctic terrestrial microarthropods. *Ecol. Entomol.* 29, 506–509.
- Hodgkinson, I.D., Coulson, S.J., Webb, N.R., Block, W., Strathdee, A.T., Bale, J.S., Worland, M.R., 1996. Temperature and the biomass of flying midges (Diptera: Chironomidae) in a high Arctic ecosystem. *Oikos* 75, 241–248.
- Hodgkinson, I.D., Coulson, S.J., Harrison, J., 2001. What a wonderful web they weave: spiders, nutrient capture and early ecosystem development in the high Arctic – some counter-intuitive ideas on community assembly. *Oikos* 95, 349–352.
- Hodgkinson, I.D., Coulson, S.J., Bird, J., Webb, N.R., 2002. Discovery of the rare alate morph of *Acyrtosiphon svalbardicum* Heikinheimo (Homoptera: Aphididae): description and implications for species ecology. *Norw. J. Entomol.* 49, 77–80.

- Hodkinson, I.D., Coulson, S.J., Webb, N.R., 2003. Community assembly on proglacial chronosequences in the high Arctic: vegetation and soil development in north west Svalbard. *J. Ecol.* 91, 651–653.
- Hodkinson, I.D., Coulson, S.J., Webb, N.R., 2004. Invertebrate community assembly along proglacial chronosequences in the high Arctic. *J. Animal Ecol.* 73, 556–568.
- Hodson, A., Anesio, A.M., Tranter, M., Fountain, A., Osborn, M., Priscu, J., Laybourn-Parry, J., Sattler, B., 2008. Glacial ecosystems. *Ecol. Monogr.* 78, 41–67.
- Holm, Å., 1958. The spiders of the Isfjord Region of Spitsbergen. *Zool. Bidr. från Upps.* 33, 29–67.
- Holm, T.M., Koinig, K.A., Andersen, T., Donali, E., Hormes, A., Klaveness, D., Psenner, R., 2011. Rapid physicochemical changes in the high Arctic Lake Kongressvatn caused by recent climate change. *Aq. Sci.: Res. Across Boundaries*, 1–11.
- Holmgren, A.E., 1869. Bidrag till kannedomen om Beeren Eilands och Spetsbergens insekt fauna. *K. Svenska Vet. Akad. Handl.* 8, 1–56.
- Holmgren, A.E., 1883. Insecta a viris Doctissimis Nordenškiöld illum ducem sequentibus in insulis Waigatsch et Novaja Semlia anno 1875 collecta. *Hymenoptera et Diptera. Entomol. Tidskr.* 4, 140–190.
- Holmstrup, M., Sømme, L., 1998. Dehydration and cold hardiness in the Arctic Collembolan *Onychiurus arcticus* Tullberg, 1876. *J. Comp. Physiol.* 168 B, 197–203.
- Holmstrup, M., Westh, P., 1994. Dehydration of earthworm cocoons exposed to cold – a novel cold-hardiness mechanism. *J. Comp. Physiol. B* 164, 312–315.
- Holmstrup, M., Bayley, M., Ramlov, H., 2002. Supercool or dehydrate? an experimental analysis of overwintering strategies in small permeable Arctic invertebrates. *Proc. Natl. Acad. Sci. U. S. A.* 99, 5716–5720.
- Hughes, K.A., Convey, P., 2010. The protection of Antarctic terrestrial ecosystems from inter- and intra-continental transfer of non-indigenous species by human activities: a review of current systems and practices. *Glob. Environ. Change* 20, 96–112.
- Hughes, K.A., Convey, P., 2012. Determining the native/non-native status of newly discovered terrestrial and freshwater species in Antarctica – current knowledge, methodology and management action. *J. Environ. Manag.* 93, 52–66.
- Hull, J.E., 1922. On some land-mites (Acaria) from Spitsbergen and Bear Island. Results of the Oxford University expedition to Spitsbergen, 1921. No. 23. *Ann Mag. Nat. Hist. Lond.* 10, 621–623.
- Hullé, M., Bonhomme, J., Maurice, D., Simon, J.C., 2008. Is the life cycle of high Arctic aphids adapted to climate change? *Polar Biol.* 31, 1037–1042.
- Humphreys, P.N., 1975. Wet-feather associated with *Holomenopon leucoxanthum* in a duck. *Vet Rec.* 97, 96–97.
- Hunter, J.E., Colwell, M.A., 1994. Phthiraptera infestation of five shorebird species. *Wilson Bull.* 106, 400–403.
- Husmann, S., Jacobi, H.U., Meijering, M.P.D., Reise, B., 1978. Distribution and ecology of Svalbarðs Cladocera. *Verh. Internat. Verein. Theor. Angew. Limnol.* 20, 2452–2456.
- Idelson, M.S., 1925. Zur Erforschung der Rotatorienfauna der Gewässer auf Nowaja Zemlya, 12. *Transact. Oceanogr. Inst.*, Moscow, pp. 79–99 (in Russian with German summary).
- Ims, R.A., Leinaas, H.P., Coulson, S.J., 2004. Spatial and temporal variation in patch occupancy and population density in a model system of the Arctic Collembola species assemblages. *Oikos* 105, 89–100.
- Ingólfsson, O., Landvik, J.Y., 2013. The Svalbard - Barents Sea ice-sheet - historical, current and future perspectives. *Quat. Sci. Rev.* 64, 33e60.
- IPCC, 2007. The Physical Science Basis. In: Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change: Summary for Policy Makers. WMO, UNEP, Geneva.
- Irvine, R.J., Stien, A., Halvorsen, O., Langvatn, R., Albon, S.D., 2000. Life-history strategies and population dynamics of abomasal nematodes in Svalbard reindeer (*Rangifer tarandus platyrhynchus*). *Parasitology* 120, 297–311.
- Irvine, R.J., Stien, A., Dallas, J.F., Halvorsen, O., Langvatn, R., Albon, S.D., 2001. Contrasting regulation of fecundity in two abomasal nematodes of Svalbard reindeer (*Rangifer tarandus platyrhynchus*). *Parasitology* 122, 673–681.
- Jacobson, G., 1898. Insecta Novaja Semljensia. Mém. l'Acad. Imp. Sci. St. Pétersbourg VIII, 1899.
- Jalava, J., Miller, W.E., 1998. Boreal Olethreutini 1. (Lepidoptera: Tortricidae): new synonymies and Holarctic records. *Entomol Fenn* 9, 137–142.
- Jamtveit, B., Hammer, Ø., Andersson, C., Dysthe, D.K., Helmann, J., Fogel, M.L., 2006. Travertines from the Troll thermal springs, Svalbard. *Norw. J. Geol.* 86, 387–395.
- Janiec, K., 1996. The comparison of freshwater invertebrates of Spitsbergen (Arctic) and King George Island (Antarctic). *Polish Polar Res.* 17, 173–202.
- Janiec, K., Salwicka, K., 1996. Monogonont composition in different freshwater habitats on Spitsbergen (Arctic) and King George Island (Antarctica). In: Proceedings of the International Symposium on Environmental Research in the Arctic: 19–21 July 1995, Tokyo, Japan, vol. 51. Memoirs of National Institute of Polar Research, pp. 91–98.
- Jeppesen, E., Christoffersen, K., Landkildehus, F., Lauridsen, T., Amsinck, S.L., Riget, F., Søndergaard, M., 2001. Fish and crustaceans in northeast Greenland lakes with special emphasis on interactions between Arctic charr (*Salvelinus alpinus*), *Lepidurus arcticus* and benthic chydorids. *Hydrobiologia* 442, 329–337.
- Johansson, C., Miller, W.R., Linder, E.T., Adams, B.J., Boreliz-Alvarado, E., 2013. Tardigrades of Alaska: distributional patterns, diversity and species richness. *Polar Res.* 32, 18793. <http://dx.doi.org/10.3402/polar.v32i0.18793>.
- Jong, Y. de, 2011. Coleoptera and Hymenoptera in Fauna Europaea. <http://www faunaeur.org>.
- Jónsdóttir, I.S., 2005. Terrestrial ecosystems on Svalbard: heterogeneity, complexity and fragility from an Arctic island perspective. *Proc. R. Irish Acad.* 105, 155–165.
- Jørgensen, I., Eie, J.A., 1993. NINA report. The Distribution of Zooplankton, Zoobenthos and Fish in Lakes and Ponds of the Mosel Peninsula, Svalbard, vol. 45. Norwegian Institute of Nature Research, Trondheim, pp. 1–25.
- Kaczmarek, Ł., Michałczyk, Ł., Eggemont, H., 2008. *Dactylobiotus luci*, a new freshwater tardigrade (Eutardigrada: Macrobiotidae) from the Rwenzori Mountains, Uganda. *Afr. Zool.* 43, 150–155.
- Kaczmarek, Ł., Gołdyn, B., Prokop, Z.M., Michałczyk, Ł., 2011. New records of Tardigrada from Bulgaria with the description of *Macrobiotus binieki* sp. nov. (Eutardigrada: Macrobiotidae) and a key to the species of the *harmsworthi* group. *Zootaxa* 2781, 29–39.
- Kaczmarek, Ł., Schabetsberger, R., Litwin, M., Michałczyk, Ł., 2012a. A new freshwater eutardigrade from Fiji and Vanuatu (Oceania), with remarks on the genus *Dactylobiotus*. *N. Z. J. Zool.* 39, 311–318.
- Kaczmarek, Ł., Zawierucha, K., Smykla, J., Michałczyk, Ł., 2012b. Tardigrada of the Revdalén (Spitsbergen) with the descriptions of two new species: *Bryodelphax parvuspolaris* (Heterotardigrada) and *Isohypsistius coulsoni* (Eutardigrada). *Polar Biol.* 35, 1013–1026.
- Kaisila, J., 1973a. The Anoplura and Siphonaptera of Spitsbergen. *Ann. Entomol. Fenn.* 39, 63–66.
- Kaisila, J., 1973b. Notes on the arthropod fauna of Spitsbergen. III. 15. The Lepidoptera of Spitsbergen. *Ann. Entomol. Fenn.* 39, 60–63.
- Kangas, E., 1967. Notes on the arthropod fauna of Spitsbergen. I: 5. Identification of the Coleoptera collected by the Finnish Spitsbergen expeditions. *Ann. Entomol. Fenn.* 33, 41–43.
- Kangas, E., 1973. Notes on the arthropod fauna of Spitsbergen. I: 2. On the Coleoptera fauna of Spitsbergen. *Ann. Entomol. Fenn.* 39, 68–70.
- Karpainen, E., 1967. Notes on the arthropod fauna of Spitsbergen. Data on the Oribatids (Acaria) from Spitsbergen. *Ann. Entomol. Fenn.* 33, 18–26.
- Karpainen, E., Krivolutsky, D.A., 1982. List of oribatid mites (Acarina, Oribatei) of northern palaeartic region. I. Europe. *Acta Ent. Fenn.* 41, 18.
- Kaya, M., De Smet, W.H., Fontaneto, D., 2010. Survey of moss-dwelling bdelloid rotifers from middle Arctic Spitsbergen (Svalbard). *Polar Biol.* 33, 833–842.
- Kennedy, C.R., 1978. The parasite fauna of resident char *Salvelinus alpinus* from Arctic islands, with special reference to Bear Island. *J. Fish Biol.* 13, 457–466.
- Kieffer, J.J., 1922. Chironomides de la Nouvelle-Zemble. In: Holtendahl, O. (Ed.), Norw. Nov. Zemlya Exp. 1921. No. 2. Det norske videnskaps-akademii, Kristiania, pp. 1–24.
- Kieffer, J.J., 1923. Serphidae in Hymenoptera. In: Holtendahl, O. (Ed.), Norw. Nov. Zemlya Exp. 1921. No. 14. Det norske videnskaps-akademii, Kristiania, p. 11.
- Killingtveit, Å., Pettersson, L.-E., Sand, K., 2003. Water balance investigations in Svalbard. *Polar Res.* 22, 161–174.
- Kim, G.H., Klochkova, T.A., Han, J.W., Kang, S.-H., Chol, H.G., Chung, K.W., Kim, S.J., 2011. Freshwater and terrestrial algae from Ny-Ålesund and Blomstrandhalvøya Island (Svalbard). *Arctic* 64, 25–31.
- Klekowski, R.Z., Opaliński, K.W., 1986. Matter and energy flow in Spitsbergen ornithogenic tundra. *Polar Res.* 4, 187–197.
- Klekowski, R.Z., Opaliński, K.W., 1989. Oxygen consumption in Tardigrada from Spitsbergen. *Polar Biol.* 9, 299–304.
- Klemetsen, A., Grotnes, P.E., Holthe, H., Kristoffersen, K., 1985. Bear island (Barents sea) charr. *Nord. J. Freshw. Res.* 62, 98–119.
- Koch, L., 1879. Arachniden aus Sibirien und Novaja Semlja eingesammelt von der schwedischen Expedition im Jahre 1875. *K. Svenska Vet. Akad. Handl.* 16.
- Koch, K.D., Meijering, M.P.D., 1985. On the distribution and ecology of Cyclopidae on Bear Island (74°30'N, 19°E). *Verh. Internat. Verein. Limnol.* 12, 3144–3148.
- Krantz, G.W., Walter, D.E., 2009. A Manual of Acarology. Texas Tech University Press, Lubbock.
- Krawczyk, W.E., Pettersson, L.-E., 2007. Chemical denudation rates and carbon dioxide drawdown in an ice-free polar karst catchment: Londonelva, Svalbard. *Permafrost. Periglacial Proces.* 18, 337–350.
- Krivolutsky, D.A., Kalyakin, V.N., 1993. Soil microfauna in environmental control on Novaya Zemlya. In: Boyarskii, P.V. (Ed.), Novaya Zemlya, vol. 2, pp. 125–131. Moscow.
- Krivolutsky, D.A., Drozdov, N.N., Lebedeva, N.V., Kaljakin, V.M., 2003. Geography of Soil Microarthropods of Arctic Islands. In: Ser. 5. Geographiya 6. Vestnik Moskovskogo Gosudarstvennogo Universiteta, pp. 33–40 (in Russian with English summary).
- Krumpá, M., Cyprich, D., Zejda, J., Ambros, M., 1991. The occurrence of field vole (*Microtus arvalis* Pallas 1778) and its acarofauna on Spitsbergen (Svalbard). *Biologia* 46, 881–885.
- Kubíček, F., Terek, J., 1991. Zooplankton Svalbárdu (Špicbergy). *Biologia* 46, 873–879.
- Kuklin, V.V., 2000. The comparative ecological analysis of the helminthofauna of seabirds of Novaya Zemlya and East Murman (east Barents Sea). *Dokl. Akad. Nauk.* 371, 139–141 (in Russian).
- Kuklin, V.V., 2001. Helminthofauna of seabirds of Archangelskaya Bay (Northern island of Novaya Zemlya). *Parasitologia* 35, 124–134 (in Russian).
- Kuklin, V.V., Kuklina, M.M., 2005. Helminths of Birds of the Barents Sea: Fauna, Ecology, Impact on the Hosts. Kol'skiy Nauchnyi Zentri Rossiyskoy Akademii Nauk. Apatity (in Russian).
- Kuklin, V.V., Galkin, A.K., Marasaeva, S.F., Marasaeva, E.F., 2004. The characteristics of the helminthofauna of sea birds of the Svalbard Archipelago. *Dokl. Biol. Sci.* 395, 124–126.

- Kuzmin, L.L., 1976. Free-living nematodes in the tundra of western Taimyr. *Oikos* 27, 501–505.
- Laasonen, E.M., 1985. Huippuvuorten perhoset. *Baptia* 10, 69–72.
- Lack, D., 1933. Trichoptera, Lepidoptera, and Coleoptera from Bear Island. *Ann. Mag. Nat. Hist.* 12, 205–210.
- Lagisz, M., 1999. Fauna Niesporczaków Tardigrada Na Nunatakach Spitsbergenu Jako Przykład Fauny Wyspowej. Jagiellonian University, Kraków. M.Sc. thesis.
- Lakka, H.-K., 2013. The Ecology of a Freshwater Crustacean:—*Lepidurus Arcticus* (Branchiopoda; Notostraca) in a High Arctic Region. The University of Helsinki, Helsinki. M.Sc. thesis.
- Landvik, J.Y., Hjort, C., Mangerud, J., Moller, F., Salvigsen, O., 1995. The Quaternary record of eastern Svalbard – an overview. *Polar Res.* 14, 95–103.
- Landvik, J.Y., Brook, E.J., Gualtieri, L., Raisbeck, G., Salvigsen, O., Yiou, F., 2003. Northwest Svalbard during the last glaciation: ice-free areas existed. *Geology* 31, 905–908.
- Lauritzen, S.E., Bottrell, S., 1994. Microbiological activity in thermoglacial karst springs, south Spitsbergen. *Geomicrobiol. J.* 12, 161–173.
- Laybourn-Parry, J., Marshall, W.A., 2003. Photosynthesis mixotrophy and microbial plankton dynamics in two high Arctic lakes during summer. *Polar Biol.* 26, 17–524.
- Lebedeva, N.V., Krivolutsky, D.A., 2003. Birds spread soil microarthropods to Arctic islands. *Dokl. Biol. Sci.* 391, 329–332.
- Lebedeva, N.V., Lebedev, V.D., 2008. Transport of Oribatid mites to the polar areas by birds. In: Integrative Acarology. Proceedings of the 6th European Congress, pp. 359–367.
- Lebedeva, N.V., Lebedev, V.D., Melekhina, E.N., 2006. New data on the oribatid mite (Oribatei) fauna of Svalbard. *Dokl. Biol. Sci.* 407, 182–186.
- Lehman, S.J., Forman, S.L., 1992. Late Weichselian glacier retreat in Kongsfjorden, west Spitsbergen, Svalbard. *Quat. Res.* 37, 139–154.
- Leinaas, H.P., Ambrose, W.G., 1992. Utilization of different foraging habitats by the Purple Sandpiper *Calidris maritima* on a Spitsbergen beach. *Fauna Norv. Serm. C* 15, 85–91.
- Leinaas, H.P., Ambrose, W.G., 1999. Decision between small and large prey: reduced energy acquisition by pre-migratory purple sandpipers, *Calidris maritima*, on Svalbard. *Polar Biol.* 22, 264–270.
- Lenz, F., Thienemann, A., 1922. Chironomidenlarven aus Nowaja Semlja. In: Holtendahl, O. (Ed.), Norw. Nov. Zemlya Exp. 1921. No. 3. Det norske viden-skaps-akademii, Kristiania, pp. 1–8.
- Lewis, R.E., Stone, E., 2001. *Psittopsylla mexicana*, a new genus and species of bird flea from Chihuahua, Mexico (Siphonaptera: Ceratophyllidae: Ceratophyllinae). *J. N. Y. Entomol. Soc.* 109, 360–366.
- Lindquist, E.E., Makarova, O.L., 2011. Two new circumpolar mite species of the genus *Arctoseius Thor* (Parasitiformes, Mesostigmata, Ascidae). *Entomol. Rev.* 91, 1054–1072.
- Linnaniemi, W.M., 1935a. Beitrag zur Kenntnis der Collembolenfauna Spitsbergens. *Ann. Entomol. Fenn.* 1, 137–141.
- Linnaniemi, W.M., 1935b. Collembolen aus Spitsbergen, Insel Hopen, Kong Karls Land und Jan Mayen, eingesammelt von norwegischen arktischen Expeditionen. *Norsk Entomol. Tidskrift* 3, 379–381.
- Löbl, I., Smetana, A., 2004. Catalogue of Palaearctic Coleoptera. In: Hydropiloidea, Histeroidea, Staphyloidea, vol. 2. Apollo Books, Stenstrup.
- Lods-Crozet, B.V., Lencioni, V., Olafsson, J.S., Snook, D.L., Velle, G., Brittain, J.E., Castella, E., Rossaro, B., 2001. Chironomid (Diptera: Chironomidae) communities in six European glacier-fed streams. *Freshw. Biol.* 46, 1791–1809.
- Lods-Crozet, B.V., Lencioni, V., Brittain, J.E., Marziali, L., Rossaro, B., 2007. Contrasting chironomid assemblages in two high Arctic streams on Svalbard. *Arch. Hydrobiol.* 170, 211–222.
- Loof, P.A.A., 1971. Freeliving and plant parasitic nematodes from Spitzbergen, collected by Mr. H. van Rossem. *Meded Landbouwhogesch Wagening* 71, 1–86.
- Loof, P.A.A., 1975. Dorylaimoidea from some sub-Antarctic islands. *Nematologica* 21, 219–255.
- Lubbock, J., 1898. On some Spitsbergen Collembola. *J. Linnean Soc.* 26, 616–619.
- Lubinski, D.J., Forman, S.L., Miller, G.H., 1999. Holocene glacier and climate fluctuations on Franz Josef Land, Arctic Russia, 80 degrees N. *Quat. Sci. Rev.* 18, 85–108.
- Luoto, T.P., Nevalainen, L., Kubischa, F., Kultti, S., Knudsen, K.L., Salonen, V.P., 2011. Late Quaternary ecological turnover in high arctic lake Einstaken, Nordaustlandet, Svalbard (80°N). *Geogr. Ann. Ser. A Phys. Geogr.* 93, 337–354.
- Maiolini, B., Lencioni, V., Brittain, J.E., 2006. Invertebrate drift in a high Arctic stream under natural and experimental light conditions: preliminary results. *Verh. Internat. Verein. Theor. Angew. Limnol.* 29, 2201–2204.
- Makarchenko, E.A., Makarchenko, M.A., Vekhov, N.V., 1998. A preliminary data on chironomid fauna (Diptera, Chironomidae) of the Archipelago Novaya Zemlya. In: Boyarsky, P.V. (Ed.), Novaya Zemlya. Nature. History. Archaeology. Culture. Book 1. Nature. Russian Res. Inst. Cultural and Natural Heritage, Moscow, pp. 262–267 (In Russian).
- Makarova, O.L., 1999. Mesostigmatic mites (Parasitiformes, Mesostigmata) of polar deserts in Severnaya Zemlya. *Entomol. Rev.* 79, 982–990.
- Makarova, O.L., 2000a. To the study of mites of the genus *Arctoseius Thor* (Parasitiformes, Ascidae) from the Far North. 1. Morphometric analysis of the taxonomic features of Arctic species and description of *Arctoseius tschernovi* sp. n. and *A. miranalensis* sp. n. *Entomol. Rev.* 80 (Suppl. 1), 112–130.
- Makarova, O.L., 2000b. To the study of mites of the genus *Arctoseius Thor* (Parasitiformes, Ascidae) from the Far North: 2. Description of *A. productus* sp. n. and *A. babenkoi* sp. n. and key to the High Arctic species. *Entomol. Rev.* 80 (Suppl. 1), 131–142.
- Makarova, O.L., 2000c. To the study of mites of the genus *Arctoseius Thor* (Parasitiformes, Ascidae) from the Far North: 3. Ranges and ecological preferences of species. *Entomol. Rev.* 80 (Suppl. 1), 143–150.
- Makarova, O.L., 2009. The fauna of free-living gamasid mites (Parasitiformes, Mesostigmata) in the northern taiga: an analysis of the zonal specificity. *Entomol. Rev.* 89, 993–1009.
- Makarova, O.L., 2011. Review of gamasid mites (Parasitiformes, Mesostigmata) in the taiga of Pechoro-Ilychskiy Reserve (Northern Cis-Ural) and the analysis of assemblages in spruce forests. *Entomol. Rev.* 91, 915–931.
- Makarova, O.L., 2013a. Gamasid mites (Parasitiformes, Mesostigmata) of the Arctic: taxonomical structure of the fauna. *Zool. Zhurnal, Moscow* (in press) (in Russian).
- Makarova, O.L., 2013b. Gamasid mites (Parasitiformes, Mesostigmata) of the European Arctic and their distribution patterns. *Entomol. Rev.* 93, 113–133.
- Makarova, O.L., Böcher, J., 2009. Diversity and geographical ranges of Greenland mites (Acaria: Oribatida and Mesostigmata). In: Golovatch, S.I., Makarova, O.L., Babenko, A.B., Penev, L.D. (Eds.), Species and Communities in Extreme Environments. Pensoft Publishers and KMK Scientific Press, Sofia–Moscow, pp. 165–186.
- Mäklin, F.W., 1881. Coleoptera insamlade under den Nordenskiöldskas expeditionen 1875 på några öar vid Norges nordvestkust, på Novaja Semlja och ön Waigatsch samt vid Jenissej i Sibirien. *Kongliga Svenska Vetenskaps-akad. Handlingar* 18, 1–48.
- Mangerud, J., Bolstad, M., Elgersma, A., Helliesen, D., Landvik, J.Y., Lonne, I., Lycke, A.K., Salvigsen, O., Sandahl, T., Svendsen, J.I., 1992. The last glacial maximum on Spitsbergen, Svalbard. *Quat. Res.* 38, 1–31.
- Mangerud, J., Kaufman, D., Hansen, J., Svendsen, J.I., 2008. Ice-free conditions in Novaya Zemlya 35,000 to 30,000 cal years BP, as indicated by radiocarbon ages and amino acid racemization evidence from marine molluscs. *Polar Res.* 27, 187–208.
- Maraldo, K., Holmstrup, M., 2010. Enchytraeids in a changing climate: a mini-review. *Pedobiologia* 53, 161–167.
- Marcus, E., 1928. Spinnentiere oder Arachoides. IV Bärtierchen (Tardigrada). In: Tierwelt Deutschlands und der angrenzenden Meeresteile, vol. 12. G. Fischer, Jena, pp. 1–230.
- Markham, A.H., 1881. A Polar Reconnaissance Being the Voyage of the "Ibsjörn" to Novaja-semjla in 1879. C. Kegan Paul & Co, London.
- Markov, G.S., 1937. Die Parasiten von *Uria lomvia* auf der Insel Nowaja Semlja. *Trav. Soc. Nat., Leningrad. Section Zool.* 66, 456–466.
- Markov, G.S., 1941. Parasitic worms of Bezymiannaya Bay (Novaya Zemlya). *Doklady Acad. Nauk SSSR* 30, 573–576 (in Russian).
- Marková, S., Dufresne, F., Manca, M., Kotík, P., 2013. Mitochondrial capture Misleads about ecological speciation in the *Daphnia pulex* complex. *Plos One* 8, e69497. <http://dx.doi.org/10.1371/journal.pone.0069497>.
- Marziali, L., Gozzini, M., Rossaro, B., Lencioni, V., 2009. Drift patterns of Chironomidae (Insecta, Diptera) in an Arctic stream (Svalbard Islands): an experimental approach. *Studi Tridentino Sci. Nat.* 84, 87–96.
- Maslen, N.R., Convey, P., 2006. Nematode diversity and distribution in the southern maritime Antarctic – clues to history? *Soil Biol. Biochem.* 38, 3141–3151.
- Maucci, W., 1996. Tardigrada of the Arctic tundra with descriptions of two new species. *Zool. J. Linn. Soc.* 116, 185–204.
- McGaughran, A., Torricelli, G., Carapelli, A., Frati, F., Stevens, M.I., Convey, P., Hogg, I.D., 2010. Contrasting phylogeographical patterns for springtails reflect different evolutionary histories between the Antarctic Peninsula and continental Antarctica. *J. Biogeogr.* 37, 103–119.
- McInnes, S.J., 1994. Zoogeographic distribution of terrestrial and freshwater tardigrades from current literature. *J. Nat. Hist.* 28, 257–352.
- Mehl, R., 1992. Fleas (Siphonaptera) from seabirds and their nests in mainland Norway and Spitzbergen. *Natl. Inst. Public Health Ann.* 15, 3–15.
- Mehl, R., Traavik, T., 1983. The tick *Ixodes uriae* (Acaria, Ixodidae) in seabird colonies in Norway. *Fauna Norv. Ser. B* 30, 94–107.
- Mehl, R., Bang, C., Kjos-Hanssen, B., Lie, H., 1982. Mallophaga from Svalbard. *Fauna Norv. Ser. B* 29, 19–23.
- Mehlum, F., 1984. Concentration of sea birds along the face of glaciers and outlets of rivers in Svalabrd. *Fauna* 37, 156–160 (in Norwegian with English summary).
- Melekhina, E.N., 2011. Taxonomic diversity and areology of Oribatid mites (Oribatei) of the European North of Russia. *Izvest. Komi Nauchnogo Centra* 2, 30–37.
- Melekhina, E.N., Zinovjeva, A.N., 2012. The first data on oribatid mites (Acaria: Oribatida) of Pay – Khoy ridge (Yugor peninsula). *Izvest. Komi Nauchnogo Centra* 2, 42–50.
- Menzel, R., 1920. Über freilebende Nematoden aus der Arktis. In: *Festschrift zur Feier des 60. Geburtstages (27. Mai 1920) von Friedrich Zschocke*, Prof., Basel, vol. 17. Prof. Basel, pp. 1–15.
- Michaelsen, W., 1900. Oligochaeta. (Das Tierreich 10). Friedländer, Berlin.
- Michalczyk, Ł., Kaczmarek, Ł., 2006. Revision of the *Echiniscus bigranulatus* group with a description of a new species *Echiniscus madonnae* (Tardigrada: Heterotardigrada: Echiniscidae) from South America. *Zootaxa* 1154, 1–26.
- Michalczyk, Ł., Weinicz, W., Frohme, M., Kaczmarek, Ł., 2012a. Redescriptions of three *Milnesium* Doyère, 1840 taxa (Tardigrada: Eutardigrada: Milnesiidae), including the nominal species for the genus. *Zootaxa* 3154, 1–20.
- Michalczyk, Ł., Weinicz, W., Frohme, M., Kaczmarek, Ł., 2012b. Corrigenda of *Zootaxa*, 3154: 1–20 Redescriptions of three *Milnesium* Doyere, 1840 taxa

- (Tardigrada: Eutardigrada: Milnesiidae), including the nominal species for the genus. *Zootaxa* 3393, 66–68.
- Milner, A.M., Brittain, J.E., Castella, E., Petts, G.E., 2001. Trends of macroinvertebrate community structure in glacier-fed rivers in relation to environmental conditions: a synthesis. *Freshw. Biol.* 46, 1833–1847.
- Ministry of Justice and the Police, 2009. Statens Meldning Number 22, Svalbard (2008–2009), Norwegian Government Stoltenberg II. Department Public Publications, Oslo.
- Mjöberg, E., 1910. Studien über Mallophagen und Anopluren. *Arkiv Zool.* 6, 1–296.
- Morten, K.J., 1923. Plecoptera. In: Holtendahl, O. (Ed.), Norw. Nov. Zemlya Exp. 1921. No. 16. Det norske videnskaps-akademi, Kristiania, pp. 1–6.
- Mortimer, E., van Vuuren, B.J., Lee, J.E., Marshall, D.J., Convey, P., Chowan, S.L., 2011. Mite dispersal among the Southern Ocean Islands and Antarctica before the last glacial maximum. *Proc. R. Soc. B Biol. Sci.* 278, 1247–1255.
- Münster, T.H., 1925. Coleoptera. In: Holtendahl, O. (Ed.), Norw. Nov. Zemlya Exp. 1921. No. 30. Det norske videnskaps-akademi, Kristiania, pp. 1–15.
- Münster, T.H., 1935. Northern *Olophrum* (Col. Staph.). *Nor Entomol Tidsskr* 4, 22–30.
- Murray, J., 1907. Arctic Tardigrada, collected by W.S. Bruce. *Trans. R. Soc. Edinb.* 45, 669–681.
- Murray, J., 1908. Arctic rotifers collected by Dr William S. Bruce. *Proc. R. Phys. Soc. Edinb.* 17, 121–127.
- Nathorst, A.G., 1884. Redögörelse för den tillsammans med G. der Geer år 1882 företagna geologiska expedition til Spetsbergen. P.A. Norstedt & Söner, Stockholm.
- Nentwig, W., Blick, T., Gloor, D., Hänggi, A., Kropf, C., 2012. Spiders of Europe. www.araneae.unibe.ch.
- Nevalainen, L., Van Damme, K., Luoto, T., Salonen, V.P., 2012. Fossil remains of an unknown *Alona* species (Chydoridae, Aloninae) from a high Arctic lake in Nordaustlandet (Svalbard) in relation to glaciation and Holocene environmental history. *Polar Biol.* 35, 325–333.
- Niedbala, W., 1971. Oribatei (Acari) of Spitsbergen. *Bull. Polish Acad. Sci.* 11, 737–742.
- Nielsen, U.N., Wall, D.H., 2013. The future of soil invertebrate communities in Polar Regions: comparable climate change responses in the Arctic and Antarctic? *Ecol. Lett.* 16, 409–419.
- Nielsen, U.N., Wall, D.H., Adams, B.J., Virginia, R.A., 2011. Antarctic nematode communities: observed and predicted responses to climate change. *Polar Biol.* 34, 1701–1711.
- Norton, R.A., Behan-Pelletier, V.M., 2009. Oribatida. In: Krantz, G., Walter, D.E. (Eds.), *A Manual of Acarology*, Third Ed. Texas Tech University Press, Lubbock.
- Norwegian Meteorological Institute, 2013. Web Service available online at: <http://retro.met.no/observasjoner/svalbard/index.html>.
- Nurminen, M., 1965. Enchytraeid and lumbricid records (Oligochaeta) from Spitsbergen. Enchytraeids (Oligochaeta) from northern Norway and western Lapland. Preliminary notes on the Enchytraeids (Oligochaeta) of the Ahvenanmaa Island, South Finland. *Ann. Zool. Fenn.* 2, 1–17.
- Nurminen, M., 1966. Further notes on the Enchytraeids (Oligochaeta) of Spitsbergen. *Ann. Zool. Fenn.* 3, 68–69.
- Nurminen, M., 1967. Faunistic notes on north-European enchytraeids (Oligochaeta). *Ann. Zool. Fenn.* 4, 567–587.
- Odasz, A.M., 1994. Nitrate reductase activity in vegetation below an Arctic bird cliff, Svalbard, Norway. *J. Veg. Sci.* 5, 913–920.
- Ødegaard, F., Tømmerås, B., 2000. Compost heaps refuges and stepping-stones for alien arthropod species in Northern Europe. *Divers. Distrib.* 6, 45–59.
- Odhner, T., 1905. Die Trematoden des arktischen Gebietes. Römer and Shaudinn. *Fauna Arctica* 4, 291–372.
- Oedekoven, M.A., Joern, A., 2000. Plant quality and spider predation affects affects grasshoppers (Acrididae): food quality dependent compensatory mortality. *Ecology* 81, 66–77.
- Økland, F., 1928. Land- and Süßwasserfauna von Nowaja Semlja. In: Norw. Nov. Zemlya Exp. 1921. No. 42. Det norske videnskaps-akademi, Kristiania, pp. 1–125.
- Olofsson, O., 1918. Studien über die Süßwasserfauna Spitzbergens. Beitrag zur Systematik, Biologie und Tiergeographie der Crustaceen und Rotatorien. *Zool. Bidr. från Upps.* 6, 183–648.
- Opaliński, K.W., Klekowski, R.Z., 1989. Oxygen consumption in *Macrotrachela muscosa* and *Trichotria truncata* (Rotatoria) from the High Arctic. *Polar Res.* 7, 133–137.
- Opaliński, K.W., Klekowski, R.Z., 1992. Metabolic adaptations to temperature in Spitsbergen invertebrates. In: Opaliński, K.W., Klekowski, R.Z. (Eds.), *Landscape, Life World and Man in High Arctic*. Institute of Ecology, Polish Academy of Sciences, Warszawa, pp. 299–306.
- Opravilova, V., 1989. Some information on testate amoebae from Spitsbergen. *Fauna Norv. Ser. A* 10, 33–37.
- Ossiannilsson, F., 1958. *Acyrtosiphon calvulus*, n.sp. a new aphid (Hemiptera, Homoptera.) from Spitzbergen. *Entomol Tidskr* 79, 66–69.
- Ott, D., Rall, B.C., Brose, U., 2012. Climate change effects on macrofaunal litter decomposition: the interplay of temperature, body masses and stoichiometry. *Philos. Trans. Royal Soc. Lond. Ser. B* 367, 3025–3032.
- Parry, W.E., 1828. Narrative of an Attempt to Reach the North Pole in Boats Fitted for Purpose, and Attached to His Majesty's Ship Hecla. J. Murray, London.
- Paus, A., Velle, G., Larsen, J., Nesje, A., Lie, O., 2006. Lateglacial nunataks in central Scandinavia: Biostratigraphical evidence for ice thickness from Lake Flafattjonn, Tynset, Norway. *Quat. Sci. Rev.* 25, 1228–1246.
- Pedersen, P.G., Holmstrup, M., 2003. Freeze or dehydrate: only two options for the survival of subzero temperatures in the arctic enchytraeid *Fridericia ratzeli*. *J. Comp. Physiol. B* 173, 601–609.
- Pejler, B., 1974. Planktonic Rotatoria from Bjørnøya, vol. 69. Norsk Polar Institutt Årbok 1972.
- Petersen, H., Luxton, M., 1982. A comparative analysis of soil fauna populations and their role in decomposition processes. *Oikos* 39, 287–388.
- Petrova, A.D., Makarova, O.L., 1991. *Arctoseius tajimyricus* – a new species of gamasid mites (Mesostigmata, Aceosejidae) phoretic on trichocerid flies (Diptera). *Entomol. Rev.* 70, 139–142.
- Pienitz, R., Doran, P.T., Lamoureux, S.C., 2008. Origin and geomorphology of lakes in the polar regions. In: Vincent, W., Laybourn-Parry, J. (Eds.), *High Latitude Lake and River Ecosystems*. Oxford University Press, Oxford, pp. 25–41.
- Pilato, G., Binda, M.G., 1987. *Isohypsibus ceciliae*, nuova specie di Eutardigrado (Hypsibiidae) dello Spitsbergen. *Animalia* 14, 125–129.
- Pilato, G., Bertolani, R., Binda, M.G., 1982. Studio degli *Isohypsibus* del gruppo *elegans* (Eutardigrada, Hypsibiidae) con descrizione di due nuove specie. *Animalia* 9, 185–198.
- Pilskog, H.E., 2011. The Invertebrate Fauna of Svalbard Bird Nests: Ecology and as Facilitating Colonisation of an Arctic Archipelago. M.Sc. The University of Bergen, Bergen.
- Platnick, N.I., 2012. The World Spider Catalog, Version 12.5. American Museum of Natural History online at <http://research.amnh.org/iz/spiders/catalog>.
- Pomorski, R.J., Skarzynski, D., 1995. Springtails (Collembola) collected in Chupa Inlet region (N. Karelia, Russia). *Acta Univ. Wratislaviensis* 1744 (Prace Zool. 29), 46–57.
- Pomorski, R.J., Skarzynski, D., 2001. Springtails (Collembola) collected in Chupa Inlet region (N. Karelia, Russia). *Acta Univ. Wratislaviensis* 1744 (Prace Zool. 29), 47–57.
- Poppius, B.R., 1910. Die Coleopteren des arktischen Gebietes. *Fauna Arctica* 5, 291–447.
- Post, E., Forchhammer, M.C., Bret-Harte, M.S., Callaghan, T.V., Christensen, T.R., Elberling, B., Fox, A.D., Gilg, O., Hik, D.S., Høye, T.T., Ims, R.A., Jeppesen, E., Klein, D.R., Madsen, J., McGuire, A.D., Rysgaard, S., Schindler, D.E., Stirling, I., Tamstorf, M.P., Tyler, N.J.C., van der Wal, R., Welker, J., Wookey, P.A., Schmidt, N.M., Aastrup, P., 2009. Ecological dynamics across the Arctic associated with recent climate change. *Science* 325, 1355–1358.
- Price, R.D., Hellenthal, R.A., Palma, R.L., Johnson, K.P., Clayton, D.H., 2003. The Chewing Lice: World Checklist and Biological Overview. Illinois Natural History Survey. Special Publication 24.
- Primicerio, R., Klemetsen, A., 1999. Zooplankton seasonal dynamics in the neighbouring lakes Takvatn and Lombola (Northern Norway). *Hydrobiologia* 411, 19–29.
- Procter, D.L.C., 1977. Nematode densities and production on Trueland Lowland. In: Bliss, L.C. (Ed.), *Trueland Lowland, Devon Island, Canada: A High Arctic Ecosystem*. University of Alberta Press, Edmonton, pp. 347–361.
- Prowse, T.D., Wrona, F.J., Reist, J.D., Hobbs, J.E., Lévesque, L.M.J., Vincent, W.F., 2006. General features of the Arctic relevant to climate change in freshwater ecosystems. *Ambio* 35, 330–338.
- Pugh, P.J.A., 1994. Non-indigenous Acari of Antarctica and the sub-Antarctic islands. *Zool. J. Linn. Soc.* 110, 207–217.
- Pugh, P.J.A., McInnes, S.J., 1998. The origin of Arctic terrestrial and freshwater tardigrades. *Polar Biol.* 19, 177–182.
- Rautio, M., Dufresne, F., Laurion, I., Bonilla, S., Warwick, S.V., Christoffersen, K.S., 2011. Shallow freshwater ecosystems of the circumpolar Arctic. *EcoScience* 18, 204–222.
- Rebel, H., 1923. Lepidoptera von Novaja Semlja. In: Holtendahl, O. (Ed.), Norw. Nov. Zemlya Exp. 1921. No. 7. Det norske videnskaps-akademi, Kristiania, p. 15.
- Rebel, H., 1925. Ueber das Auffinden einer Noctuide auf Spitzbergen. *Nor Entomol Tidsskr* 2, 53.
- Retowsky, L., 1935. Süßwasserfauna von Nowaja Semlja und Franz-Joseph Land, 14. *Transactions of the Arctic Institute, Leningrad*, pp. 3–78 (in Russian with German summary).
- Richard, J., 1898. Sur la faune des eaux douces explorées en 1898 pendant la campagne du yacht Princesse-Alice (Lofoten, Spitsberg, îles Beeren, Hope, de Barents et Færøer). *Mém. Soc. zool. France* 11, 326–338.
- Richard, K.J., Convey, P., Block, W., 1994. The terrestrial arthropod fauna of the Byers peninsula, south Shetland islands. *Polar Biol.* 14, 371–379.
- Richards, W.R., 1963. The Aphidiidae of the Canadian Arctic (Homoptera). *Can. Entomol.* 95, 449–464.
- Richters, F., 1903. Nordische Tardigraden. *Zool. Anz.* 27, 168–172.
- Richters, F., 1904. Arktische Tardigraden. *Fauna Arctica* 3, 283–286.
- Richters, F., 1911. Faune des mousses. Tardigrades. In: Duc d'Orléans. *Campagne arctique de 1907. Imprimerie scientifique Charles Bulens, Bruxelles*.
- Rothschild, N.C., 1922. Siphonaptera. In: Holtendahl, O. (Ed.), Norw. Nov. Zemlya Exp. 1921. No. 4. Det norske videnskaps-akademi, Kristiania, pp. 1–9.
- Rozema, J., 1999. Stratospheric Ozone Depletion, the Effects of Enhanced UV-b Radiation on Terrestrial Ecosystems. Backhuys, Leiden.
- Ruess, L., Michelsen, A., Jonasson, S., 1999a. Simulated climate change in subarctic soils: responses in nematode species composition and dominance structure. *Nematology* 1, 513–526.
- Ruess, L., Michelsen, A., Schmidt, I.K., Jonasson, S., 1999b. Simulated climate change affecting microorganisms, nematode density and biodiversity in subarctic soils. *Plant Soil* 212, 63–73.
- Sack, P., 1923. Dipteren aus Nowaja Semlja. In: Holtendahl, O. (Ed.), Norw. Nov. Zemlya Exp. 1921. No. 15. Det norske videnskaps-akademi, Kristiania, pp. 1–10.

- Sahlberg, J.R., 1897. Staphylinidae in Novaja Semlja a G. Jacobson et in Insulis Novo-Sibiricis a Dr. A. Bunge et Bar. Ed. Toll Collectae. Annuaire du Musée Zoologique de l'Académie Impériale des Sciences de St. Pétersbourg 2, pp. 365–368.
- Sahlberg, J., 1901. Coleoptera in den Polargegenden. In: Medd. Soc. Fauna et Flora Fennica 27. Helsingfors.
- Samchysyna, L., Hansson, L.A., Christoffersen, K., 2008. Patterns in the distribution of Arctic freshwater zooplankton related to glaciation history. *Polar Biol.* 31, 1427–1435.
- Schäffer, C., 1895. Verzeichnis der von den Herren Dr. Küenthal und Dr. Walter auf Spitzbergen gesammelten Collembolen. In: *Zoologische Jahrbücher*, vol. 8. Abteilung für Systematik, Geographie und Biologie der Tiere, pp. 128–130.
- Schäffer, C., 1900. Die arktischen und subarktischen Collembola. *Fauna Arctica* 1, 147–216.
- Schneeweiss, G.M., Schönswetter, P., 2011. A re-appraisal of nunatak survival in Arctic-alpine phylogeography. *Mol. Ecol.* 20, 190–192.
- Schött, H., 1899. Fyra Collembolformer från Franz Josephs land. In: *Förhandlingar 15 Skandinaviska Naturforskarmötet* 7–12 Juli 1898, Stockholm.
- Schött, H., 1923. Collembola. In: Holtendahl, O. (Ed.), *Norw. Nov. Zemlya Exp.* 1921. No. 12. Det norske videnskaps-akademii, Kristiania, pp. 1–14.
- Scott, F.L.S., 1899. Report on the marine and freshwater Crustacea from Franz-Josef land collected by Mr. William S. Bruce of the Jackson-Harmsworth expedition. *J. Linnean Soc. Lond. Zool.* 27, 60–126.
- Scourfield, D.J., 1897. Contributions to the non-marine fauna of Spitsbergen. Part I. Preliminary notes, and reports on the Rhipozoda, Tardigrada, Entomostraca. *Proc. Zool. Soc. Lond.*, 784–792.
- Segers, H., 2008. Global diversity of rotifers (Rotifera) in freshwater. *Hydrobiologia* 595, 49–59.
- Sendstad, E., Bergvik, T., Hegstad, A., 1976. *Plusia interrogationis* l. (Lepidoptera, Noctuidae) found at Svalbard. *Norw. J. Entomol.* 23, 91–92.
- Seniczak, S., Plichta, W., 1978. Structural dependence of moss mite populations (Acarı, Oribatei) on patchiness of vegetation in moss-lichen-tundra at the north coast of Hornsund, West Spitsbergen. *Pedobiologia* 18, 145–152.
- Seniczak, A., Solhøy, T., Seniczak, S., 2006. Oribatid mites (Acarı: Oribatida) in the glacier foreland at Hardangerjokulen (Norway). *Biol. Lett.* 43, 231–235.
- Serebryanny, L., Andreev, A., Malyasova, E., Tarasov, P., Romanenko, F., 1998. Late-glacial and early-Holocene environments of Novaya Zemlya and Kara Sea region of the Russian Arctic. *Holocene* 8, 323–330.
- Serreze, M.C., Holland, M.M., Stroeve, J., 2007. Perspectives on the Arctic's Shrinking sea-ice cover. *Science* 315, 1533–1536.
- Serreze, M.C., Barrett, A.P., Cassano, J.J., 2011. Circulation and surface controls on the lower tropospheric air temperature field of the Arctic. *J. Geophys. Res.* 116, Article ID D07104.
- Simmons, B.L., Wall, D.H., Adams, B.J., Ayres, E.J., Barrett, E., Virginia, R.A., 2009. Long-term experimental warming reduces soil nematode populations in the McMurdo Dry Valleys, Antarctica. *Soil Biol. Biochem.* 41, 2052–2060.
- Simon, J.C., Bonhomme, J., Blackman, R.L., Hullé, M., 2008. Winged morph of the high Arctic aphid *Acyrtosiphon svalbardicum* (Hemiptera: Aphididae): abundance, reproductive status, and ecological significance. *Can. Entomol.* 140, 385–387.
- Skogseth, R., Haugin, P.M., Jakobsen, M., 2005. Watermass transformations in Storfjorden. *Cont. Shelf Res.* 25, 667–695.
- Skorikow, A.S., 1900. Zoologische Ergebnisse der Russischen Expedition nach Spitzbergen im Jahre 1899. In: Collembola. *EZhegodnik Zoologicheskogo Muzeya Imperatorskoi Akademii Nauk, St.-Petersburg*, vol. 5, pp. 190–209.
- Skou, P., 1984. Nordens Måler. Håndbog over de danske og fennoskandiske arter af Drepanidae og Geometridae (Lepidoptera). In: *Danmarks Dyreliv*, vol. 2. Fauna Bøger og Apollo Bøger, København og Svendborg.
- Skrede, I., Eidense, P., Piñeiro Portela, R., Brochman, C., 2006. Refugia, differentiation and postglacial migration in Arctic-alpine Eurasia, exemplified by the mountain avens (*Dryas octopetala* L.). *Mol. Ecol.* 15, 1827–1840.
- Skrýabin, K.I., 1926. Study of helminthological collection gathered by G.Y. Sedov, expeditions to North Pole. *Trudy GIEV* 4, 114–122 (in Russian).
- Smol, J.P., Douglas, M.S.V., 2007. Crossing the final ecological threshold in high Arctic ponds. *Proc. Natl. Acad. Sci. U. S. A.* 104, 12395–12397.
- Smykla, J., Kaczmarek, Ł., Huzarska, K., Michalczyk, Ł., 2011. The first record of a rare marine tardigrade, *Halobiotus crispae* Kristensen, 1982 (Eutardigrada: Hypsistidae), from the Svalbard Archipelago. *Polar Biol.* 34, 1243–1247.
- Sobecka, E., Piasecki, W., 1993. Parasitic fauna of Arctic charr, *Salvelinus alpinus* (L., 1758) from the Hornsund region (Spitsbergen). *Acta Ichyl. Piscatoria* 23, 99–106.
- Sohlenius, B., Boström, S., 1999. Effects of global warming on nematode diversity in a Swedish tundra soil - a soil transplantation experiment. *Nematology* 1, 695–709.
- Sohlenius, B., Boström, S., Ekebom, A., 1997. Metazoan microfauna in an ombrotrophic mire at Abisko, northern Sweden. *Eur. J. Soil Biol.* 33, 31–39.
- Solhøy, T., 1976. *Camisia foveolata* Hammer, 1955 (Acarı, Oribatei) found in Norway and on Svalbard. *Norw. J. Entomol.* 23, 89.
- Speight, M.R., Hunter, M.D., Watt, A.D., 1999. Ecology of Insects. Concepts and Applications. Blackwell Science Ltd, Oxford.
- Stach, J., 1962. On the Collembola from Spitsbergen. *Acta Zool. Cracoviensis* 7, 1–20.
- Steiner, G., 1916. Freilebende Nematoden von Novaja-Semlja. *Zool. Anzeiger* 47, 50–74.
- Stempniewicz, L., Zwolicki, A., Iliszko, L., Zmudzynska, K., Wojtun, B., 2006. Impact of plankton- and fish-eating seabird colonies on the Arctic tundra ecosystem—a comparison. *J. Ornithol.* 147, 257–258.
- Stephenson, J., 1922. The Oligochaeta of the Oxford University Spitsbergen expedition. *Proc. Zool. Soc. Lond.*, 1109–1138.
- Stephenson, J., 1924. On some Oligochaete worms from Spitsbergen. Results of the Merton College expedition to Spitsbergen, 1923. *vol. 3. Ann. Mag. Nat. Hist.* 13, 210–216.
- Stephenson, J., 1925. The Oligochaeta of Spitsbergen and Bear Island: some additions and summary. *Proc. Zool. Soc. Lond.*, 1293–1322.
- Stevens, R.D., 2006. Historical processes enhance patterns of diversity along latitudinal gradients. *Proc. Royal Soc. B Biol. Sci.* 273, 2283–2289.
- Stevens, M.I., Greenslade, P., Hogg, I.D., Sunnucks, P., 2006. Southern Hemisphere Springtails: could any have survived glaciation of Antarctica? *Mol. Biol. Evol.* 23, 874–882.
- Stevens, M.I., Frati, F., McGaughan, A., Spisanti, G., Hogg, I.D., 2007. Phylogeographic structure suggests multiple glacial refugia in Northern Victoria Land for the endemic Antarctic springtail *Desoria klovstadi* (Collembola, Isotomidae). *Zool Scripta* 36, 201–212.
- Stien, A., Irvine, R.J., Ropstad, E., Halvorsen, O., Langvatn, R., Albon, S.D., 2002. The impact of gastrointestinal nematodes on wild reindeer: experimental and cross-sectional studies. *J. Animal Ecol.* 71, 937–945.
- Stien, A., Voutilainen, L., Haukisalmi, V., Fuglei, E., Mork, T., Yoccoz, N.G., Ims, R.A., Henttonen, H., 2010. Intestinal parasites of the Arctic fox in relation to the abundance and distribution of intermediate hosts. *Parasitology* 137, 149–157.
- Stöp-Bowitz, C., 1969. A contribution to our knowledge of the systematics and zoogeography of Norwegian earthworms (Annelida Oligochaeta: Lumbricidae). *Nytt Mag. Zool.* 17, 169–276.
- Strand, A., 1942. Die Käferfauna von Svalbard. In: *Meddeleser*, 54. Norwegian Polar Institute, Oslo.
- Strand, A., 1969. *Simplocaria metallica* Sturm new to Svalbard (Coleoptera, Byrrhidae). *Norsk Entomol. Tidsskrift* 16, 23–24.
- Strathdee, A.T., Bale, J.S., 1995. Factors limiting the distribution of *Acyrtosiphon svalbardicum* (Hemiptera: Aphididae) on Spitsbergen. *Polar Biol.* 15, 375–380.
- Strathdee, A.T., Bale, J.S., 1998. Life on the edge: insect ecology in Arctic environments. *Annu. Rev. Entomol.* 43, 85–106.
- Strathdee, A.T., Bale, J.S., Block, W.C., Webb, N.R., Hodkinson, I.D., Coulson, S.J., 1993. Extreme adaptive life-cycle in a high Arctic aphid, *Acyrtosiphon svalbardicum*. *Ecol. Entomol.* 18, 254–258.
- Strathdee, A.T., Bale, J.S., Strathdee, F.C., Block, W., Coulson, S.J., Hodkinson, I.D., Webb, N.R., 1995. Climatic severity and the response to warming of Arctic aphids. *Glob. Change Biol.* 1, 23–28.
- Strecker, A.L., Milne, R., Arnott, S.E., 2008. Dispersal limitation and climate-related environmental gradients structure microcrustacean composition in freshwater lakes, Ellesmere Island, Canada. *Can. J. Fish. Aquat. Sci.* 65, 1905–1918.
- Stur, E., Ekrem, T., 2011. Exploring unknown life stages of Arctic Tanytarsini (Diptera: Chironomidae) with DNA barcoding. *Zootaxa* 2743, 27–39.
- Styczynski, B., Rakusa-Suszczewski, S., 1963. Tendipedidae of selected water habitats of Hornsund region (Spitsbergen). *Pol. Arch. Hydrobiol.* 11, 327–341.
- Suatoni, E., Vicario, S., Rice, S., Snell, T., Caccone, A., 2006. An analysis of species boundaries and biogeographic patterns in a cryptic species complex: the rotifer *Brachionus plicatilis*. *Mol. Phylogenet. Evol.* 41, 86–98.
- Summerhayes, V.S., Elton, C.S., 1923. Contributions to the ecology of Spitsbergen and Bear Island. *J. Ecol.* 11, 214–286.
- Summerhayes, V.S., Elton, C.S., 1928. Further contributions to the ecology of Spitsbergen. *J. Ecol.* 16, 193–268.
- Svendsen, J.I., Mangerud, J., 1997. Holocene glacial and climatic variations on Spitsbergen, Svalbard. *Holocene* 7, 45–57.
- Svenning, M.A., Gullestad, N., 2002. Adaptations to stochastic environmental variations: the effects of seasonal temperatures on the migratory window of Svalbard Arctic charr. *Environ. Biol. Fishes* 64, 165–174.
- Svenning, M.-A., Klemetsen, A., Olsen, T., 2007. Habitat and food choice of Arctic charr in Linnévatn on Spitsbergen, Svalbard: the first year-round investigation in a High Arctic lake. *Ecol. Freshw. Fish* 16, 70–77.
- Sywula, T., Namiotko, T., Sell, J., Witkowski, A., Zajaczkowski, M., 1994. Crustacean species new to Spitsbergen with notes on the polymorphism and the subfossil preservation of *Cytherissa lacustris* (G.O. Sars). *Polar Res.* 13, 233–235.
- Sæther, O.A., Spies, M., 2012. Fauna Europaea: Chironomidae. In: de Jong, H., Pape, T. (Eds.), *Nematocera* (accessed 03.06.12.) <http://www.faunaeur.org>.
- Sømme, S., 1934. Contributions to the Biology of Norwegian Fish Food Animals. I. *Lepidurus arcticus*. Avhand. Norske Videnskaps-Akademie Oslo 6, pp. 1–36.
- Sømme, L., 1979. Insektsliv på Svalbard. *Fauna* 32, 137–144.
- Sømme, L., 1999. The physiology of cold hardiness in terrestrial arthropods. *Eur. J. Entomol.* 96, 1–10.
- Sømme, L., Birkemoe, T., 1997. Cold tolerance and dehydration in Enchytraeidae from Svalbard. *J. Comp. Physiol. B* 167, 264–269.
- Sørensen, J.G., Holmstrup, M., 2011. Cryoprotective dehydration is widespread in Arctic springtails. *J. Insect Physiol.* 57, 1147–1153.
- Sørensen, L.I., Holmstrup, M., Maraldo, K., Christensen, S., Christensen, B., 2006. Soil fauna communities and microbial respiration in high Arctic tundra soils at Zackenberg, Northeast Greenland. *Polar Biol.* 29, 189–195.
- Søvik, G., 2004. The biology and life history of Arctic populations of the littoral mite *Ameronothrus lineatus* (Acarı: Oribatida). *Exp. Appl. Acarol.* 34, 3–20.
- Tambs-Lyche, H., 1967. Notes on the distribution of some Arctic spiders. *Astarte* 28, 1–13.
- Tanasevitch, A., 2012. Linyphiid Spiders of the World. <http://www.andtan.newmail.ru/list/index.htm>.

- Taskaeva, A.A., 2009. Springtail (Collembola) assemblages in Floodlands of the taiga zone of the Republic of Komi. Entomol. Rev. 89, 965–974.
- Taskaeva, A.A., Nakul, G.L., 2010. Collembolan populations in colonial nesting sites of seagulls in Kolokolkova Bay. In: Taskaev, A.I., Kosolapov, D.A., Valuyskikh, O.E. (Eds.), Relevant Questions of Biology and Ecology. Komi Science Center UB RAS, Syktyvkar, pp. 120–122 (in Russian).
- Taylor, A.L., 1981. Adventitious molt in red knot possibly caused by *Actornithophilus* (Mallophaga: Menoponidae). J. Field Ornithol. 52, 241.
- Thomas, D.N., Fogg, G.E., Convey, P., Fritsen, C.H., Gili, J.M., Gradinger, R., Laybourn-Parry, J., Reid, K., Walton, D.W.H., 2008. The Biology of Polar Regions. Oxford University Press, Oxford.
- Thomasson, K., 1958. Zur Planktonkunde Spitzbergens. Hydrobiologia 12, 226–236.
- Thomasson, K., 1961. Zur Planktonkunde Spitzbergens 2. Hydrobiologia 18, 192–198.
- Thor, S., 1930. Beiträge zur Kenntnis der invertebraten Fauna von Svalbard. In: Skrifter om Svalbard og Ishavet, vol. 27. Norwegian Polar Institute, Oslo.
- Thor, S., 1934. Neue Beiträge zur Kenntnis der invertebraten Fauna von Svalbard. Zool. Anz. 107, 114–139.
- Thorell, T., 1871. Om Arachnider från Spitzbergen och Beeren-Eiland. Övers. Kongl. Vet. Akad. Förh. 28, 683–702.
- Timmermann, G., 1957. Studien zu einer vergleichenden Parasitologie der Charadriiformes oder Regeleifervögel, Teil 1: Mallophaga. Parasitol. Schriftreihe 8, 1–204.
- Tkach, N.V., Röser, M., Hoffman, M., 2008. Range size variation and diversity distribution in the vascular plant flora of the Eurasian Arctic. Org. Diver. Evolut. 8, 251–266.
- Todaro, M.A., Hummon, W.D., 2012. Ubiquity of microscopic animals? Evidence from the morphological approach in species identification. Gastrotricha. In: Fontaneto, D. (Ed.), Biogeography of Microscopic Organisms: Is Everything Small Everywhere? Cambridge University Press, Cambridge, pp. 245–249.
- Todaro, M.A., Dal Zotto, M., Jondelius, U., Hochberg, R., Hummon, W.D., Kåneby, T., Rocha, C.E.F., 2012. Gastrotricha: a marine Sister for a freshwater Puzzle. PLoS ONE 7, 1–11. Article Number: e31740.
- Trägårdh, I., 1901. Revision der von Thorell aus Grönland, Spitzbergen und der Bären Insel und von L. Koch aus Siberien und Novaja Semlja beschriebenen Acariden. Zool. Anz. 25, 56–62.
- Trägårdh, I., 1904. Monographie der arktischen Acariden. Fauna Arctica 4, 1–78.
- Trägårdh, I., 1928. Acari (excl. Sarcoptidae). In: Holtendahl, O. (Ed.), Norw. Nov. Zemlya Exp. 1921. No. 40. Det norske videnskaps-akademii, Kristiania, pp. 1–11.
- Treaty of Spitsbergen, February 9, 1920. Treaty Concerning the Archipelago of Spitsbergen, Signed at Paris. In: League of Nations Treaty Series. Available from: <http://www.lovdata.no/traktater/texte/tre-19200209-001.html> (accessed 01.02.13.).
- Triplet, F., Christe, P., Möller, A.P., 2002. The importance of host spatial distribution for parasite specialization and speciation: a comparative study of bird fleas (Siphonaptera: Ceratophyllidae). J. Animal Ecol. 71, 735–748.
- Trouessart, E., 1895. Révision des Acariens des régions Arctiques et description d'espèces nouvelles. Mém. Soc. nat. sci. natur. math. Cherbourg, 183–206.
- Tullberg, T., 1876. Collembola borealia. Nordiska Collembola beskrifna af Tycho Tullberg. Översikt af Kongl. Vetenskaps Akad. Förfhandlinger B 33, 23–42.
- Tumanov, D.V., 2006. Five new species of the genus *Milnesium* (Tardigrada, Eutardigrada, Milnesiidae). Zootaxa 1122, 1–23.
- Turetsky, M.R., Bond-Lamberty, B., Euskirchen, E., Talbot, J., Frolking, S., McGuire, A.D., Tuittila, E.S., 2012. The resilience and functional role of moss in boreal and Arctic ecosystems. New Phytol. 196, 49–67.
- Ude, H., 1902. Die arktischen Enchytraiden und Lumbriciden sowie die geographische Verbreitung dieser Familien. Fauna Arctica 2, 1–34.
- Ullrich, W.G., Campbell, J.M., 1974. A revision of the apterus-group of the genus *Tachinus* Gravenhorst (Coleoptera: Staphylinidae). Can. Entomol. 106, 627–644.
- Ulmer, G., 1925. Ephemeropteren und Trichopteren von Novaya Zemlya. In: Holtendahl, O. (Ed.), Norw. Nov. Zemlya Exp. 1921, No. 29. Det norske videnskaps-akademii, Kristiania, p. 4.
- Usher, M.B., Booth, R.G., 1984. Arthropod communities in a maritime Antarctic moss-turf habitat: three-dimensional distribution of mites and Collembola. J. Animal Ecol. 53, 427–441.
- Usher, M.B., Booth, R.G., 1986. Arthropod communities in a maritime Antarctic moss-turf habitat: multiple scales of pattern in the mites and Collembola. J. Animal Ecol. 55, 155–170.
- Uspenskiy, S.M., Govorukha, L.S., Belikov, S.Y., Bulavintsev, V.I., 1987. Proposed protected zones in the Franz-Josef Land area. Polar Geog. Geol. 11, 210–220.
- Valdecasas, A.G., Camacho, A.J., Pelaez, M.L., 2006. Do small animals have a biogeography? Exp. Appl. Acarol. 40, 133–144.
- Valpas, A., 1967. Collemboles of Spitsbergen. Ann. Entomol. Fennici 33, 28–40.
- van der Wal, R., Hessen, D.O., 2009. Analogous aquatic and terrestrial food webs in the high Arctic: the structuring force of a harsh climate. Persp. Plant Ecol. Evolut. Syst. 11, 231–240.
- van der Wal, R., Irvine, J., Stien, A., Shepherd, N., Albon, S.D., 2000. Faecal avoidance and the risk of infection by nematodes in a natural population of reindeer. Oecologia 124, 19–25.
- Van Donk, E., Faafeng, B.A., de Lange, H.J., Hessen, D.O., 2001. Differential sensitivity to natural ultraviolet radiation among phytoplankton species in Arctic lakes (Spitsbergen, Norway). Plant Ecol. 154, 249–259.
- Van Geest, G.J., Hessen, D.O., Spijrenburg, P., Dahl-Hansen, G.A.P., Christensen, G., Faerovig, P.J., Brehm, M., Loonen, M.J.J.E., Van Donk, E., 2007. Goose-mediated nutrient enrichment and planktonic grazer control in Arctic freshwater ponds. Oecologia 153, 653–662.
- Van Helldingen, P., 1984. World distribution of Linyphiidae. In: Eberhard, W.G., Lubin, Y.D., Robinson, B. (Eds.), Proceedings of the 9th Congress of Arachnology. Smithsonian Institution Press, Washington, pp. 121–126.
- Van Rompu, E.A., De Smet, W.H., 1988. Some aquatic Tardigrada from Bjørnøya (Svalbard). Fauna Norv. Ser. A 9, 31–36.
- Van Rompu, E.A., De Smet, W.H., 1991. Contribution to the freshwater Tardigrada from Barentsøya, Svalbard (78° 30'N). Fauna Norv. Ser. A 12, 29–39.
- Van Rompu, E.A., De Smet, W.H., 1994. Freshwater tardigrades from Hopen, Svalbard (76° 31'N). Fauna Norv. Ser. A 17, 1–9.
- Vekhoff, N.V., 1997. Large branchiopod Crustacea (Anostraca, Notostraca, Spinicaudata) of the Barents region of Russia. Hydrobiologia 359, 69–74.
- Velichko, A.A., 2002. Dynamics of Terrestrial Landscape Components and Inner Marine Basins of Northern Eurasia during the Last 130 000 Years. Atlas-monograph, GEOS, Moscow.
- Velle, G., Kongshavn, K., Birks, H.J.B., 2011. Minimizing the edge-effect in environmental reconstructions by trimming the calibration set: chironomid-inferred temperatures from Spitsbergen. Holocene 21, 417–430.
- Vestby, S.E., 1983. En Regional Limnologisk Undersøkelse Av Små Innsjøer Ved Svea, Svalbard. Institute for Marine Biology and Limnology, University of Oslo, Oslo. M.Sc. thesis.
- Vicente, F., Bertolani, R., 2013. Considerations on the taxonomy of the Phylum Tardigrada. Zootaxa 3626, 245–248.
- Vincent, W.F., MacIntyre, S., Spigel, R.H., Laurion, I., 2008. The physical limnology of high-latitude lakes. In: Vincent, W.F., Laybourn-Parry, J. (Eds.), Polar Lakes and Rivers. Limnology of Arctic and Antarctic Aquatic Ecosystems. Oxford University Press, Oxford, pp. 65–82.
- von Goes, A., 1862. Om Tardigrader Anguillulae m. f. Spetsbergen. Ofv. K. Svenska Vet.-Akad. Forh. 19, p. 18.
- Vyverman, W., Verleyen, E., Wilmette, A., Hodgson, D.A., Willem, A., Peeters, K., Van de Vijver, B., De Wever, A., Leliaert, F., Sabbe, K., 2010. Evidence for widespread endemism among Antarctic micro-organisms. Polar Sci. 4, 103–113.
- Ware, C., Bergstrom, D.M., Müller, E., Alsos, I.G., 2011. Humans introduce viable seeds to the Arctic on footwear. Biol. Invas. 14, 567–577.
- Waterston, J., 1922a. On the Mallophaga of the Spitsbergen expedition. Trans. Entomol. Soc. Lond. 70, 251–253.
- Waterston, J., 1922b. Results of the Oxford University expedition to Spitsbergen, 1921. No. 25. Hymenoptera Parasitica: Ichneumonidae. Ann. Mag. Nat. Hist. 11, 31–33.
- Webb, N.R., Coulson, S.J., Hodgkinson, I.D., Block, W., Bale, J.S., Strathdee, A.T., 1998. The effects of experimental temperature elevation on populations of cryptostigmatic mites in high Arctic soils. Pedobiologia 42, 298–308.
- Weider, L.J., Hobæk, A., 2000. Phylogeography and Arctic biodiversity: a review. Ann. Zool. Fenn. 37, 217–231.
- Weider, L.J., Hobæk, A., Colbourne, J.K., Crease, T.J., Dufresne, F., Herbert, P.D.N., 1999. Holartic phylogeography of an asexual species complex—I. Mitochondrial DNA variation in Arctic *Daphnia*. Evolution 53, 777–792.
- Weigmann, G., 2006. Hornmilben (Oribatida). Goecke and Evers, Keltern.
- Westergaard, K.B., Alsos, I.G., Popp, M., Engelskjøn, T., Flatberg, K.I., Brochmann, C., 2011. Glacial survival may matter after all: nunatak signatures in the rare European populations of two west-Arctic species. Mol. Ecol. 20, 376–393.
- Weinitz, W., Grohme, M.A., Kaczmarek, L., Schill, R.O., Frohme, M., 2011. Anhydrosis in tardigrades-The last decade. J. Insect Physiol. 57 (5), 577–583.
- Wharton, D.A., 1986. A Functional Biology of Nematodes. Croom Helm, London.
- Wharton, D.A., 2002. Life at the Limits. Cambridge University Press, Cambridge.
- Williams, B.L., Griffiths, B.S., 1989. Enhanced nutrient mineralization and leaching from decomposing sitka spruce litter by enchytraeid worms. Soil Biol. Biochem. 21, 183–188.
- Wilson, R.R., Bartsch, A., Joly, K., Reynolds, J.H., Orlando, A., Loya, W.M., 2013. Frequency, timing, extent, and size of winter thaw-refreeze events in Alaska 2001–2008 detected by remotely sensed microwave backscatter data. Polar Biol. 36, 419–426.
- Winston, J.E., 1999. Describing Species. In: Practical Taxonomic Procedure for Biologists. Columbia University Press, New York.
- Wohlfarth, B., Lemdahl, G., Olsson, S., Persson, T., Snowball, I., Ising, J., Jones, V., 1995. Early Holocene environment on Bjørnøya (Svalbard) inferred from multidisciplinary lake sediment studies. Polar Res. 14, 253–275.
- Woods, P.J., 2011. Ecological Diversity in the Polymorphic Fish Arctic Charr (*Salvelinus alpinus*). Ph.D. thesis, University of Washington.
- Woodward, G., Dybkjær, J.B., Ólafsson, J.S., Gíslason, G.M., Hannesdóttir, Friberg, N., 2010. Sentinel systems on the razor's edge: effects of warming on Arctic geothermal stream ecosystems. Glob. Change Biol. 16, 1979–1991.
- Worland, M.R., Grubor-Lajsic, G., Montiel, P.O., 1998. Partial desiccation induced by sub-zero temperatures as a component of the survival strategy of the Arctic collembolan *Onychiurus arcticus* (Tullberg). J. Insect Physiol. 44, 211–219.
- Weglarska, B., 1965. Die Tardigraden (Tardigrada) Spitzbergens. Acta Zool Cracov 11, 43–51.
- Yu, D.S., van Achterberg, K., Horstmann, K., 2005. World Ichneumonidea. In: Taxonomy, Biology, Morphology and Distribution. CD/DVD. Taxapad, Vancouver.
- Yunakov, N.N., Korotyaev, B.A., 2007. A review of the weevil subgenus *Metaphyllobius* Smirnov (Coleoptera, Curculionidae, Entiminae) from eastern Europe and Siberia. Entomol. Rev. 87, 1045–1059.
- Zachariassen, K.E., 1985. Physiology of cold tolerance in insects. Physiol. Rev. 65, 799–832.

- Zawierucha, K., Coulson, S.J., Michalczyk, Ł., Kaczmarek, Ł., 2013. Current knowledge on the Tardigrada of Svalbard with the first records of water bears from Nordaustlandet (High Arctic). *Polar Res.*. In press.
- Zawisza, E., Szeroczyńska, K., 2011. Cladocera species composition in lakes in the area of the Hornsund Fjord (Southern Spitsbergen) – preliminary results. *Knowl. Manag. Aq. Ecosyst.* 402, 1–9.
- Zeeberg, J., 2002. Climate and Glacial History of Novaya Zemlya Archipelago, Russian Arctic, with Notes on the Region's History or Exploration. Rozenberg Publishers, Amsterdam.
- Zeeberg, J., Forman, S.J., 2001. Changes in glacier extent on north Novaya Zemlya in the twentieth century. *Holocene* 11, 161–175.
- Zilli, A., Varga, Z., Ronkay, G., Ronkay, L., 2009. Apameini I. The Witt Catalogue. A Taxonomic Atlas of the Eurasian and North African Noctuoidea, vol. 3. Heterocera Press, Budapest.
- Zmudczyńska, K., Zwolicki, A., Barcikowski, M., Barcikowski, A., Stempniewicz, L., 2009. Spectral characteristics of the Arctic ornithogenic tundra vegetation in Hornsund area, SW Spitsbergen. *Polish Polar Res.* 30, 249–262.
- Zmudczyńska, K., Olejniczak, I., Zwolicki, A., Iliszko, L., Convey, P., Stempniewicz, L., 2012. The influence of allochthonous nutrients delivered by colonial seabirds on soil collembolan communities on Spitsbergen. *Polar Biol.* 35, 1233–1245.
- Zschokke, F., 1903. Die arktischen Cestoden. *Fauna Arctica* 3, 1–32.
- Zschokke, F., 1926. Collembolen aus Spitzbergen. *Rev. D'Hydrobiol.* 3, 127–128.
- Zwolicki, A., Zmudczyńska-Skarbek, K.M., Iliszko, L., Stempniewicz, L., 2013. Guano deposition and nutrient enrichment in the vicinity of planktivorous and piscivorous seabird colonies in Spitsbergen. *Polar Biol.* 36, 363–372.