Key to scoring ICD-11 and DSM-5-TR criteria for PGD				
Diagnostic criteria	Symptoms	Symptom Present = 1 Not present = 0	ICD-11 criteria	DSM-5-TR criteria
<b>Criterion A:</b> Experienced the death of a loved one.	A. A close relative died or a person?		Present	Present
Criterion B: Time criterion.	B1. $\geq$ 6 months B2. $\geq$ 12 months		≥ 6 months	≥ 12 months
<b>Criterion C:</b> The presence of one or more of the following intrusive symptoms which emerged following the death.	C1. Yearning *  C2. Preoccupation*		≥ 1 symptom	≥ 1 symptom
	-			
Criterion D: Negative changes in cognition, mood and/or behavior that appeared or got worse after the death	D1. Sadness*  D2. Sense of guilt  D3. Self-blame  D4. Anger*  D5. Disbelief/denial *  D6. Lack of acceptance  D7. Loss of identity/part of self died*  D8. Absence of positive emotions/mood  D9. Emotional numbness  D10. Reduced social activity *  D11. Loneliness  D12. Avoidance  D13. Meaninglessness  D14. Bitterness  Total score in Criterion D  (ICD-11)  Total score in Criterion D  (DSM-5-TR)  Clinical assessment of if the		≥ 1 symptom	≥ 3 symptoms  Present
	client is overall afflicted by intense emotional pain			
Criterion E: The loss results in clinically significant impairment or reduction in the ability to function socially, occupational, or in other important areas	E. Functional impairment (general) E1. Work/studies/Daily activities E2. Social E3. Family/home life E4. other areas		Present	Present
Criterion F: The grief reaction clearly exceeds social, cultural, or religious norms within the individual's culture and context.	F1. The social network F2. One's own perspective		Present	Present
Criterion G: Symptom duration	F3. Interviewer's assessment Symptom duration		Duration: ≥6 months	Duration: ≥1 months
The symptoms are not better accoun	1 nted for by another diagnosis, pi	hysiological react		monuis
Identified Prolonged Grief Disord criteria?  If yes, mark a cross in the appropria	_			